



Standard Test Method for Determination of Individual Components in Spark Ignition Engine Fuels by 100–Metre Capillary (with Precolumn) High- Resolution Gas Chromatography¹

This standard is issued under the fixed designation D6730; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This test method covers the determination of individual hydrocarbon components of spark-ignition engine fuels and their mixtures containing oxygenate blends (MTBE, ETBE, ethanol, and so forth) with boiling ranges up to 225 °C. Other light liquid hydrocarbon mixtures typically encountered in petroleum refining operations, such as blending stocks (naphthas, reformates, alkylates, and so forth) may also be analyzed; however, statistical data was obtained only with blended spark-ignition engine fuels.

1.2 Based on the cooperative study results, individual component concentrations and precision are determined in the range from 0.01 % to approximately 30 % by mass. The test method may be applicable to higher and lower concentrations for the individual components; however, the user must verify the accuracy if the test method is used for components with concentrations outside the specified ranges.

1.3 This test method also determines methanol, ethanol, *t*-butanol, methyl *t*-butyl ether (MTBE), ethyl *t*-butyl ether (ETBE), and *t*-amyl methyl ether (TAME) in spark ignition engine fuels in the concentration range from 1 % to 30 % by mass. However, the cooperative study data provided insufficient statistical data for obtaining a precision statement for these compounds.

1.4 Although a majority of the individual hydrocarbons present are determined, some co-elution of compounds is encountered. If this test method is utilized to estimate bulk hydrocarbon group-type composition (PONA), the user of such data should be cautioned that some error will be encountered due to co-elution and a lack of identification of all components present. Samples containing significant amounts of naphthenic

(for example, virgin naphthas) constituents above *n*-octane may reflect significant errors in PONA-type groupings. Based on the gasoline samples in the interlaboratory cooperative study, this test method is applicable to samples containing less than 25 % by mass of olefins. However, some interfering co-elution with the olefins above C₇ is possible, particularly if blending components or their higher boiling cuts such as those derived from fluid catalytic cracking (FCC) are analyzed, and the total olefin content may not be accurate. **Annex A1** of this test method compares results of the test method with other test methods for selected components, including olefins, and several group types for several interlaboratory cooperative study samples. Although benzene, toluene, and several oxygenates are determined, when doubtful as to the analytical results of these components, confirmatory analyses can be obtained by using the specific test methods listed in the reference section.

1.4.1 Total olefins in the samples may be obtained or confirmed, or both, if necessary, by Test Method **D1319** (percent by volume) or other test methods, such as those based on multidimensional PONA-type of instruments.

1.5 If water is or is suspected of being present, its concentration may be determined, if desired, by the use of Test Method **D1744** or equivalent. Other compounds containing oxygen, sulfur, nitrogen, and so forth, may also be present, and may co-elute with the hydrocarbons. If determination of these specific compounds is required, it is recommended that test methods for these specific materials be used, such as Test Methods **D4815** and **D5599** for oxygenates, and Test Method **D5623** for sulfur compounds, or equivalent.

1.6 The values stated in SI units are to be regarded as standard. No other units of measurement are included in this standard.

1.7 *This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.*

¹ This test method is under the jurisdiction of ASTM Committee **D02** on Petroleum Products, Liquid Fuels, and Lubricants and is the direct responsibility of Subcommittee **D02.04.0L** on Gas Chromatography Methods.

Current edition approved April 1, 2016. Published May 2016. Originally approved in 2001. Last previous edition approved in 2011 as D6730–01 (2011). DOI: 10.1520/D6730-01R16.

2. Referenced Documents

2.1 ASTM Standards:²

- D1319** Test Method for Hydrocarbon Types in Liquid Petroleum Products by Fluorescent Indicator Adsorption
- D1744** Test Method for Determination of Water in Liquid Petroleum Products by Karl Fischer Reagent (Withdrawn 2016)³
- D3700** Practice for Obtaining LPG Samples Using a Floating Piston Cylinder
- D4057** Practice for Manual Sampling of Petroleum and Petroleum Products
- D4177** Practice for Automatic Sampling of Petroleum and Petroleum Products
- D4307** Practice for Preparation of Liquid Blends for Use as Analytical Standards
- D4626** Practice for Calculation of Gas Chromatographic Response Factors
- D4815** Test Method for Determination of MTBE, ETBE, TAME, DIPE, tertiary-Amyl Alcohol and C₁ to C₄ Alcohols in Gasoline by Gas Chromatography
- D5580** Test Method for Determination of Benzene, Toluene, Ethylbenzene, *p/m*-Xylene, *o*-Xylene, C₉ and Heavier Aromatics, and Total Aromatics in Finished Gasoline by Gas Chromatography
- D5599** Test Method for Determination of Oxygenates in Gasoline by Gas Chromatography and Oxygen Selective Flame Ionization Detection
- D5623** Test Method for Sulfur Compounds in Light Petroleum Liquids by Gas Chromatography and Sulfur Selective Detection
- E355** Practice for Gas Chromatography Terms and Relationships
- E594** Practice for Testing Flame Ionization Detectors Used in Gas or Supercritical Fluid Chromatography
- E1510** Practice for Installing Fused Silica Open Tubular Capillary Columns in Gas Chromatographs

3. Terminology

3.1 *Definitions*—This test method makes reference to many common gas chromatographic procedures, terms, and relationships. Detailed definitions can be found in Practice **E355**.

4. Summary of Test Method

4.1 A representative sample of the petroleum liquid is introduced into a gas chromatograph equipped with an open tubular (capillary) column coated with a methyl silicone liquid phase, modified with a capillary precolumn. Helium carrier gas transports the vaporized sample through the column, in which it is partitioned into individual components which are sensed with a flame ionization detector as they elute from the end of the column. The detector signal is presented on a strip chart recorder or digitally, or both, by way of an integrator or

integrating computer. Each eluting component is identified by comparing its retention time to that established by analyzing reference standards or samples under identical conditions. The concentration of each component in percent by mass is determined by normalization of the peak areas after correction with detector response factors. Unknown components are reported as a total unknown percent by mass.

5. Significance and Use

5.1 Knowledge of the individual component composition (speciation) of gasoline fuels and blending stocks is useful for refinery quality control and product specification. Process control and product specification compliance for many individual hydrocarbons can be determined through the use of this test method.

5.2 This test method is adopted from earlier development and enhancement.^{4,5,6,7} The chromatographic operating conditions and column tuning process, included in this test method, were developed to provide and enhance the separation and subsequent determination of many individual components not obtained with previous single-column analyses. The column temperature program profile is selected to afford the maximum resolution of possible co-eluting components, especially where these are of two different compound types (for example, a paraffin and a naphthene).

5.3 Although a majority of the individual hydrocarbons present in petroleum distillates are determined, some co-elution of compounds is encountered. If this test method is utilized to determine bulk hydrocarbon group-type composition (PONA), the user of such data should be cautioned that some error will be encountered due to co-elution and a lack of identification of all components present. Samples containing significant amounts of olefinic or naphthenic, or both, constituents above octane may reflect significant errors in PONA-type groupings.

5.4 If water is or is suspected of being present, its concentration is determined by the use of Test Method **D1744**. Other compounds containing oxygen, sulfur, nitrogen, and so forth may also be present, and may co-elute with the hydrocarbons. When known co-elution exists, these are noted in the test method data tables. If determination of these specific compounds is required, it is recommended that test methods for these specific materials be used, such as Test Method **D4815** and **D5599** for oxygenates, Test Method **D5580** for aromatics, and Test Method **D5623** for sulfur compounds.

⁴ Johansen, N.G., and Etre, L.S., "Retention Index Values of Hydrocarbons on Open Tubular Columns Coated with Methyl Silicone Liquid Phases," *Chromatographia*, Vol 5, No. 10, October 1982.

⁵ Johansen, N.G., Etre, L.S., and Miller, R.L., "Quantitative Analysis of Hydrocarbons by Structural Group Type in Gasolines and Distillates. Part 1," *Journal of Chromatography*, Vol 256, 1983, pp. 393–417.

⁶ Kopp, V.R., Bones, C.J., Doerr, D.G., Ho, S.P., and Schubert, A.J., "Heavy Hydrocarbon/Volatility Study: Fuel Blending and Analysis for the Auto/Oil Air Quality Improvement Research Program," SAE Paper No. 930143, March 1993.

⁷ Schubert, A.J. and Johansen, N.J., "Cooperative Study to Evaluate a Standard Test Method for the Speciation of Gasolines by Capillary Gas Chromatography," SAE Paper No. 930144, March 1993.

² For referenced ASTM standards, visit the ASTM website, www.astm.org, or contact ASTM Customer Service at service@astm.org. For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.

³ The last approved version of this historical standard is referenced on www.astm.org.

6. Apparatus

6.1 *Gas Chromatograph*—Instrumentation capable of column oven temperature programming, from subambient (5 °C) to at least 200 °C, in 0.1 °C/min or less rate increments, is required. Multi-step column oven temperature programming is required, consisting of an initial hold time, an initial temperature program followed by an isothermal temperature hold and another programmed temperature rise. A heated flash vaporizing injector designed to provide a linear sample split injection (that is, 200:1) is required for proper sample introduction. The associated carrier gas controls must be of sufficient precision to provide reproducible column flows and split ratios in order to maintain analytical integrity. A hydrogen flame ionization detector, with associated gas controls and electronics, designed for optimum response with open tubular columns, shall conform to the specifications as described in Practice E594, as well as having an operating temperature range of up to at least 250 °C.

6.2 *Sample Introduction*—Manual or automatic liquid sample injection to the splitting injector may be employed. Automated injections are highly recommended. Micro-syringes, auto-syringe samplers, or valves capable of 0.1 µL to 0.5 µL injections are suitable. It should be noted that some syringes and improper injection techniques as well as inadequate splitter design could result in sample fractionation. This must be determined in accordance with Section 10.

6.3 *Electronic Integrator*—Any electronic integration device used for quantitating these analyses shall meet or exceed these minimum requirements:

- 6.3.1 Capacity to handle 400 or more peaks per analysis.
- 6.3.2 Normalized area percent calculation with response factors.
- 6.3.3 Noise and spike rejection.
- 6.3.4 Accurate area determination of fast (1 s to 2 s) peaks (10 Hz or greater sampling rate).
- 6.3.5 Maintain peak detection sensitivity for narrow and broad peaks.
- 6.3.6 Positive and negative sloping baseline correction.
- 6.3.7 Perpendicular drop and tangent skimming as needed.
- 6.3.8 Display of baseline used to ensure correct peak area determination.

6.4 *Open Tubular Column*—The column used for this test method consists of a primary (100 m) analytical column and a precolumn. The ability to provide the required component separations is dependent on the precise control of the column selectivity, which is typically slightly more than that exhibited by current commercially available columns. Some older columns, and columns that have a sample residue from repeated use without conditioning, may exhibit the required polarity. Until adequate columns are commercially available, the currently used methyl silicone columns can be modified or *tuned* to meet the method column specifications. See Section 11 for a description of the column performance specifications and Annex A1 for a description of the column modification procedure.

6.4.1 The primary gas chromatographic column used for this test method will meet the following specifications.

Material	fused silica
Length	100 m
Internal diameter	0.25 mm
Liquid phase	methyl silicone
Film thickness	0.50 µm
Theoretical plates, n, pentane at 35 °C	~ 400 000 to 500 000
Retention factor, k, pentane at 35 °C	0.45 to 0.50
Resolution, R, <i>t</i> -butanol and 2-methylbutene-2 at 35 °C	3.25 to 5.25
Peak symmetry, <i>t</i> -butanol at 35 °C	> 1.0 to < 5.0

6.4.2 *Precolumn*—A variable length (1 m to 4 m) of 5 % phenyl/95 % dimethylpolysiloxane fused silica open tubular column (0.25 mm inside diameter) is added to the front (injector) end of the 100 m column, as described in Annex A1.

7. Reagents and Materials

7.1 *Carrier Gas*—Helium, 99.999 % pure. (**Warning**—Helium, air, nitrogen, compressed gas under pressure.)

7.2 *Oxidant*—Air, 99.999 % pure. (**Warning**—see 7.1.)

7.3 *Detector Makeup Gas*—Nitrogen, 99.999 % pure. (**Warning**—see 7.1.)

7.4 *Fuel Gas*—Hydrogen, 99.999 % pure. (**Warning**—Hydrogen, flammable gas under high pressure.)

7.5 *Reference Standards:*

7.5.1 *Purity of Reagents*—Reagent grade chemicals shall be used in all tests. Unless otherwise indicated, it is intended that all reagents conform to the specifications of the Committee on Analytical Reagents of the American Chemical Society⁸ where such specifications are available. Other grades may be used, provided it is first ascertained that the reagent is of sufficiently high purity to permit its use without lessening the accuracy of the determination.

7.5.2 *Methanol*—(**Warning**—These materials are flammable and may be harmful or fatal, if ingested or inhaled.)

7.5.3 *Ethanol*—Only absolute ethanol of 99.5 minimum percent meets the requirements of this test method. (**Warning**—see 7.5.2.)

7.5.4 *Hydrocarbon and Other Component References*—Individual and mixed component reference materials are commercially available and may be used to establish qualitative and quantitative calibration. (**Warning**—see 7.5.2.)

7.5.5 *System and Column Evaluation Mixture*—A quantitatively prepared mixture, complying with Practice D4307, of individual hydrocarbons and oxygenates of interest is used for system and column evaluation (see Table 1). (**Warning**—see 7.5.2.) Fig. 1 is a chromatogram of the recommended mixture in Table 1.

8. Sampling

8.1 Hydrocarbon liquids with Reid vapor pressures of 110 kPa (16 psi) or less may be sampled either into a floating piston cylinder or into an open container (Practices D4057 and D4177). If the sample as received does not meet the upper

⁸ *Reagent Chemicals, American Chemical Society Specifications*, American Chemical Society, Washington, DC. For suggestions on the testing of reagents not listed by the American Chemical Society, see *Analar Standards for Laboratory Chemicals*, BDH Ltd., Poole, Dorset, U.K., and the United States Pharmacopeia and National Formulary, U.S. Pharmacopeial Convention, Inc. (USPC), Rockville, MD.

TABLE 1 System and Column Evaluation Mixture

	%
Ethanol	8.00
<i>n</i> -pentane	2.00
<i>t</i> -butanol	0.50
2-methylbutene-2	2.50
2,3-dimethylbutane	0.50
Methyl- <i>t</i> -butyl ether	10.00
<i>n</i> -hexane	2.00
1-methylcyclopentene	0.50
Benzene	1.00
Cyclohexane	28.90
3-ethylpentane	0.20
1,2 <i>t</i> -dimethylcyclopentane	0.50
<i>n</i> -heptane	2.00
2,3,3-trimethylpentane	0.50
Toluene	7.00
<i>n</i> -octane	2.00
Ethylbenzene	25.00
<i>p</i> -xylene	1.00
2,3-dimethylheptane	0.20
<i>n</i> -nonane	2.00
5-methylnonane	0.20
1-methyl-2-ethylbenzene	0.50
<i>n</i> -decane	1.00
<i>n</i> -undecane	0.50
1,2,3,5-tetramethylbenzene	0.25
Naphthalene	0.50
<i>n</i> -dodecane	0.25
1-methylnaphthalene	0.25
<i>n</i> -tridecane	0.25

boiling range requirements of 1.1, it may be necessary to extend the analysis time and raise the upper column temperature of this test method to ensure complete elution of higher boiling range sample material from the column.

8.1.1 *Piston Cylinder Sampling*—Refer to Practice **D3700** for instructions on transferring a representative sample of a hydrocarbon fluid from a source into a floating piston cylinder. Add inert gas to the ballast side of the floating piston cylinder to achieve a pressure of 350 kPa (45 psi) above the vapor pressure of the sample.

8.1.2 *Open Container Sampling*—Refer to Practice **D4057** for instructions on manual sampling from bulk storage into open containers. Stopper the container immediately after taking a sample.

8.2 Preserve the sample by cooling to approximately 4 °C and maintaining that temperature prior to analysis.

8.3 Transfer an aliquot of the cooled sample to a precooled septum vial and seal immediately.

8.4 Obtain the test specimen for analysis directly from the sealed septum vial, for either manual or automatic injection.

9. Preparation of Apparatus

9.1 Install the 100 m column and, if required, a precolumn according to the manufacturer's or supplier's instructions and **Annex A1**. See Practice **E1510/8** for recommended installation procedures.

9.2 Determine the required length of the precolumn in accordance with **Annex A1**. Adjust the operating conditions of the gas chromatograph to those listed in **Table 2** or as determined by Section **12** and **Annex A1**.

9.3 During setup and, when not performing analyses, it is advisable to turn off the cryogenic operation and set the column oven temperature at 35 °C. Attach the column outlet to the flame ionization detector inlet and check for leaks throughout the system. If leaks are found, tighten or replace fittings before proceeding.

9.4 Confirm or adjust, or both, the column carrier gas flow rate by making injections of methane or natural gas. *The methane retention time shall be 7.00 min ± 0.02 min with the column oven temperature at 35 °C*, which results in an average linear velocity of 24 cm/s, as determined using **Eq 1**. This will result in a methane retention time of 6.53 min at 5 °C. Raising or lowering the carrier gas pressure to the injector makes flow rate adjustment. A starting point of 277 kPa (40 psig) helium pressure is recommended, although columns requiring as high as 332 kPa (48 psig) helium have been encountered.

$$\text{average linear gas velocity: } u_{\text{ave}} \text{ (cm/s)} = \text{column length (cm)} / t_{M(s)} \quad (1)$$

9.5 After final adjustment of the carrier gas flow rate, note the carrier gas inlet pressure. Measure and, if necessary, readjust the injector split flow rate to give the specified or desired split ratio. Calculate the column outlet flow rate using **9.5.1** and the split ratio using **9.5.2**.

9.5.1 *Column Carrier Gas Flow Rate (at outlet):*

9.5.1.1 $P = (\text{head pressure (psig)} + \text{ambient pressure}) / \text{ambient pressure}$.

9.5.1.2 $j = \text{compressibility factor} = 3/2((P^2-1)/(P^3-1))$.

9.5.1.3 $u_o = u_{\text{ave}} j$ = column outlet velocity.

9.5.1.4 $A_c = \text{pi}(r)^2$ = column cross-sectional area (cm²), where r = column internal radius (cm).

9.5.1.5 Flow rate (cm³/min) = $u_o \times A_c \times 60$.

9.5.2 *Injection Split Ratio*—(Split flow rate + column flow rate)/column flow rate.

9.5.3 *Example*—Using a 100 m × 0.25 mm capillary column:

9.5.3.1 $U_{\text{ave}} = 100 \times 100/6.98 \times 60 = 23.88 \text{ cm/s}$.

9.5.3.2 $P = 40 \text{ psig} + 12.0/12.0 = 4.33$.

9.5.3.3 $j = 3/2((18.778-1)/(81.370-1)) = 0.33$

9.5.3.4 $u_o = 23.88/0.33 = 71.96 \text{ cm/s}$.

9.5.3.5 $A_c = \text{pi}(0.025/2)^2 = 4.9 \times 10^{-4} \text{ cm}^2$.

9.5.3.6 Flow rate = $71.96 \times 4.9 \times 10^{-4} \times 60 = 2.12 \text{ cm}^3/\text{min}$.

9.5.3.7 Split Ratio = $(192 + 2.12)/2.12 = 91.6:1$.

9.6 Make a blank analysis (no sample injection) run to ensure proper instrument operation and further condition the column and instrumentation. If stray peaks or a rising baseline signal is observed, the column oven shall be kept at the upper temperature until the baseline becomes steady and returns to within approximately 5 % of the starting temperature detector signal.

9.7 After any extended conditioning period, or if the instrument has been shut down, it is advisable to repeat **9.4**, **9.5**, and **9.6** to ensure proper carrier gas flows are being used and the column is clean.

10. Split Injection Linearity

10.1 Splitting injector linearity must be established to determine proper quantitative parameters and limits. The split




FIG. 1 DHA Speciation Analysis—System and Column Evaluation Mixture (7.5.5)

ratio used is dependent upon the split linearity characteristics of the particular injector and the sample retention factor of the

column. The retention factor of a particular column for a sample component is proportional to the amount of liquid

TABLE 2 GC Operating Conditions

Column Temperature Program	
Initial temperature	5 °C
Initial time	10 min.
First program rate	5.0 °/min
First hold temperature	50 °C
First hold time	to the elution of ethylbenzene (~50 min)
Second program rate	1.5 °/min
Final temperature	200 °C
Final hold time	5 min
Injector	
Temperature	250 °C
Split ratio	150:1
Sample size	0.1 µL – 0.2 µL
Detector	
Type	flame ionization
Temperature	250 °C
Use manufacturers recommended detector gas flows or:	
Fuel gas	hydrogen at 30 mL/min
Oxidant	air at 300 mL/min
Make-up gas, where required	nitrogen at 20 mL/min
Carrier Gas	
Type	helium
Pressure	~ 277 kPa (40 psig)
Average linear velocity	24 cm/s at 35 °C

phase (loading or film thickness) and the ratio of the column temperature to the component boiling point (vapor pressure). Overloading of the column may cause loss of resolution for some components and, since overloaded peaks are skewed, variance in retention times. This can lead to erroneous component identification. During column evaluations and split linearity studies, be aware of any peaks that may appear *front skewed*, indicating column overload. Note the component size and avoid conditions leading to this problem during actual analyses.

10.2 Set the injector temperature and split ratio to the following values and, for each set of conditions inject the listed quantities of the system and column evaluation mixture (7.5.5), using the operating conditions listed in Table 2 or as determined in Section 12.

injector temperature: 250 °C < $\begin{matrix} \text{split: 100:1} \\ \text{split: 200:1} \end{matrix}$ > sample: 0.2 µL, 0.5 µL, 1.0 µL

injector temperature: 300 °C < $\begin{matrix} \text{split: 100:1} \\ \text{split: 200:1} \end{matrix}$ > sample: 0.2 µL, 0.5 µL, 1.0 µL

10.3 Compare the calculated concentrations to the known standard concentrations after calculating the corrected area normalization using the response factors from 13.2 and Table A1.1.

$$\% \text{ relative error} = \quad (2)$$

$$100 \times (\text{concentration determined} - \text{concentration known}) / \text{concentration known}$$

10.4 Report and use only those combinations of conditions from 10.2 that result in 3 % or less relative error. This is the splitter linearity range.

11. Column Evaluation

11.1 In order to establish that a column will perform as required, the following specifications shall be determined for new column acceptability and are useful for periodic evaluation of column deterioration. These specification determina-

tions can be made with or without a precolumn, since the precolumn will have little effect on their values. See Annex A1, Fig. A1.1, for examples of these determinations. After performing the steps in Sections 9 and 10, analyze the column performance mixture (7.5.5) at 35 °C isothermal, at least through heptane. The remainder of the analysis may be ignored, but the remaining components must be eluted from the column prior to performing another analysis. Setting the column temperature to 220 °C for an additional 20 min will be sufficient.

11.2 Calculate the retention factor (k) for pentane at 35 °C:

$$k = (t_R - t_M) / t_M \quad (3)$$

where:

t_M = gas holdup time (methane), and
 t_R = retention time for pentane, min.

11.2.1 The retention factor must be between 0.45 and 0.50 for proper application of this test method.

11.3 Calculate the column efficiency using the pentane peak:

$$n = 5.545 (t_R / w_{1/2h})^2 \quad (4)$$

where:

n = column efficiency (theoretical plates),
 t_R = retention time of pentane, and
 $w_{1/2h}$ = peak width at half height.

11.3.1 The column efficiency must be at least 400 000 plates for proper application of this test method.

11.4 The selectivity of apparently identical columns toward hydrocarbons may vary regarding oxygenated compounds; either due to extraneous materials in the liquid phase, or due to activity of the column wall surface. The addition of a precolumn has little if any affect on the selectivity toward oxygenates (see Annex A1, Fig. A1.4). The relative resolution of oxygenates is inherent to the quality of the primary 100 m column, and is specified by the resolution of *t*-butanol from 2-methylbutene-2 at 35 °C. Calculate the resolution:

$$R = 2(t_{R2-M-Butene-2} - t_{RTBA}) / 1.699(w_{1/2h2-M-Butene-2} + w_{1/2hTBA}) \quad (5)$$

11.4.1 The resolution for this pair at 35 °C must be between 3.25 and 5.25.

11.5 Extraneous column effects, or instrumental effects such as an active injector liner, may cause adsorption of oxygenated compounds, commonly seen and referred to as *tailing*, and may increase their retention. If this effect is caused by instrumental activity, the problem should be corrected. If the column is inherently active, a new column should be obtained. A measure of the tailing can be made and specified by applying a *skewness* calculation, which determines a ratio of the distances from the peak apex perpendicular to the front and back of the peak at 5 % of the peak height. See Annex A1, Fig. A1.3 for an example of this calculation.

$$\text{skewness} = B/A \quad (6)$$

11.5.1 This test method shall be made using the *t*-butanol peak (0.5 %) in the analysis of the column performance

mixture (7.5.5) at 35 °C isothermal. The skewness ratio must be greater than 1.0 and not more than 5.0.

12. Optimization of Instrument Operating Conditions

12.1 The column temperature programming profile is dependent upon the individual column characteristics. Table 2 lists the programming profile determined for a 100 m methyl silicone column with a precolumn as determined in Annex A1. The profile is determined by establishing satisfactory separations for the sets of sample components listed in 12.3. It is not practical to expect complete separation of all components, so the optimum for each column may contain some compromises, also dependent upon any particular other separations deemed important.

12.2 The use of retention indices to numerically express the relative location of components among themselves and to surrounding normal paraffins is a convenient convention. The indices are also useful in providing a system of component identification with complex analyses such as this. There are several schemes for calculating retention indices, the first of which is the Kovats method, developed to express the logarithmic relationship of retention times of a homologous series of compounds when chromatographed isothermally. While this test method is not an isothermal column temperature procedure, it does contain isothermal steps and the longer temperature program step is a slow rate. The use of the Kovats indices provides a closer relationship to previous work in this field than using the linear index format.

12.2.1 The formula for the calculation of Kovats retention indices is:

$$RI_i = 100 \times (n + (\log(t_i) - \log(t_n)) / (\log(t_{n+1}) - \log(t_n))) \quad (7)$$

where:

- RI = retention index,
- n = carbon number of n -paraffin,
- t_i = retention time of component,
- t_n = retention time of preceding n -paraffin, and
- t_{n+1} = retention time of next n -paraffin.

12.3 The following examples show the key or critical separations required for this analysis. Typical retention indices are given, and a description of the effect of instrumental conditions on the separation is provided.

12.3.1 *i-butane/methanol and ethanol/3-methylbutene-1*—The initial starting temperature of 5 °C is dictated by these separations. A lower starting temperature is not necessary and a higher temperature would effect the next set. The retention indices should be about 380 for methanol and 456.5 for ethanol (Fig. 2).

12.3.2 *i-propanol/2-methylbutene-1 and t-butanol/2-methylbutene-2*—*i*-propanol will appear resolved between pentene-1 and 2-methyl-butene-1, *t*-butanol will appear resolved between *c*-pentene-2 and 2-methylbutene-2.

12.3.2.1 Higher temperatures will move the alcohols into the peaks ahead of them. At 35 °C the alcohols will be located ahead of the pentene-1 and *c*-pentene-2, respectively (Fig. 3).

12.3.3 *2,3-dimethylbutane/methyl-t-butylether*—This separation is critical and the 5 °C hold for 10 min determines its success. The retention indices should be about 569.5, 571.5,




FIG. 2 *i*-butane/methanol and ethanol/3-methyl-butene-1




FIG. 3 *i* propanol/2-methyl/butene-1 and *t* butanol/2-methylbutene-2

and 574.0 for 2,3-dimethylbutane, MTBE, and 2-methylpentane, respectively. If the MTBE is too close to the 2,3-DMC₄, use a 9 min initial hold. If too close to the 2-MC₅ use an 11 min hold (Fig. 4).

12.3.4 *1-methylcyclopentene/benzene*—This is a key separation that is used to specify the column selectivity. Changing column temperature produces only slight differences in this resolution (Fig. 5).

12.3.4.1 The 50 °C column temperature is held isothermal until the elution of ethylbenzene. This is variable due to slight differences in the column retention factor.




FIG. 4 2,3-dimethylbutane/methyl-*t* butylether




FIG. 5 1-methylcyclopentene/benzene

12.3.5 *2,3,3-trimethylpentane/toluene*—This is a key separation that is used to specify the column selectivity. Column temperature has very little effect on this resolution, which is controlled by the column selectivity for aromatics (Fig. 6).

12.3.6 *p-xylene/2,3-dimethylheptane* —This is a key separation which limits the maximum length of the precolumn. If the column selectivity is too great the aromatics are retained and this separation is not achieved. If this resolution is excessive and the separation in 12.3.5 is insufficient, the precolumn should be lengthened slightly. Lowering the 50 °C hold temperature to 48 °C will increase this separation (Fig. 7).

12.3.7 *I17 (Unknown)/1,2-methylethylbenzene* —The unknown isoparaffin (I17) appears to be a component of alkylate and must be resolved from the aromatic. If the resolution is incomplete the final column temperature program rate of 1.5 °/min. is adjusted to provide sufficient separation. Increase the rate in 0.1 °/min increments to increase the resolution. This rate is also dictated by the separation requirements in 12.3.8. The proper rate will provide for both separations (Fig. 8).

12.3.8 *1-methylnaphthalene/tridecane* —The recommended final column temperature program rate of 1.5 °/min. should also provide this separation. If the 1-MeNaph/*n*-C₁₃ resolution is incomplete this rate may be adjusted to provide sufficient




FIG. 7 *p-xylene/2,3-dimethylheptane*




FIG. 8 I17 (unknown)/1,2-methylethylbenzene

separation. Lower the rate in 0.1 °/min. increments to increase the resolution (Fig. 9).

13. Calibration

13.1 *Qualitative*—Determine the retention times of components by analyzing known reference mixtures or samples under identical conditions. Calculate retention indices from these data using 12.2. Table A1.1 provides a listing of typical values for this test method.

13.2 *Quantitative, Hydrocarbons*—Use theoretical response factors for correction of the detector response of hydrocarbons determined by this test method, unless response factors have been determined experimentally. The response of an FID to hydrocarbons is determined by the ratio of the molecular weight of the carbon in the analyte to the total molecular weight of the analyte. If experimentally determined response




FIG. 6 2,3,3-trimethylpentane/toluene




FIG. 9 1-methylnaphthalene/tridecane

factors are to be used, they must be determined using known purity individual standards and calculated using Practice D4626. The response factors, as listed in Table 3, are relative to that calculated for heptane. Calculations are based on the following equation:

$$F_i = (((((C_{aw} \times C_n) + (H_{aw} \times H_n)) / C_n) \times 0.83905) / C_{aw}) \quad (8)$$

where:

F_i = relative response factor for a hydrocarbon type group of a particular carbon number.

C_{aw} = atomic weight of carbon 12.011,

C_n = number of carbon molecules in the group,

H_{aw} = atomic weight of hydrogen, 1.008,

H_n = number of hydrogen molecules in the group,
0.83905 is the correction factor with heptane as unity (1.0000), and

0.7487 is used with methane as unity.

13.3 *Quantitative, Oxygenates*—Determine response factors for methanol, ethanol, and other oxygenated compounds experimentally. The principles in Practice D4626 should be applied when determining these response factors. The response of the flame ionization detector for oxygenated compounds is not directly (theoretically) related to mass concentration. A study has indicated that the FID response is linear for the conditions of this test method (see Figs. 10 and 11). Each individual apparatus must be calibrated using gravimetrically prepared standards, covering the sample concentration ranges expected and the scope of this test method. Standards used must comply with the requirements in Section 7. Figs. 10 and 11 present calibration data for six oxygenates as determined in a preliminary cooperative study report for calibration of this test method. Precision data will be prepared when more data becomes available.

14. Sample Analysis Procedure

14.1 Adjust the instrument operating variables to the values specified in Table 1 or as determined in Section 12.

14.2 Set the recorder or integration device, or both, for accurate presentation and collection of the data.

14.3 Inject an appropriate size sample (as determined in Section 10) into the injection port and start the analysis. Obtain a chromatogram and a peak integration report.

15. Calculation

15.1 Identify each peak by matching retention indices (or retention times) with those for known reference standards or sample components. If a computing integrator is used, examine the chromatographic data for proper peak integration. Examine the report to ensure peaks are properly identified.

15.1.1 Proper component identification using retention indices requires the use of *windows* surrounding each RI value in order to account for the analysis to analysis variations. The following windows have been found to provide satisfactory identification for this test method.

Indices	Window
100 – 300	± 15
300 – 400	± 2.6
400 – 500	± 1.5
500 – 885	± 0.6
885 – 900	± 0.5
> 900	± 0.6

15.2 Obtain the area for each peak. Multiply each peak area by its appropriate response factor, taken from Table 2 or determined separately with standards, to obtain corrected peak areas. Use a response factor of 1.000 for unknown peaks.

15.3 If required, determine the concentration of water in the sample using Test Method D1744, or an equivalent method. The total concentration of any other materials not determined by this test method should also be obtained.

15.4 The corrected peak areas are normalized to 100 % or to 100 % minus the concentrations determined in 15.3.

$$\text{component \% (m/m)} = \text{corrected peak area} \quad (9)$$

$$\times (100 - \% \text{ undetected}) / \text{total corrected peak area}$$

16. Report

16.1 Report the concentration of each component as mass %, % (m/m), to the nearest 0.001 % (m/m).

16.2 These individual component data may be grouped by summing the concentration of compounds in each particular group type such as paraffin, isoparaffin, olefin, aromatic, naphthene, oxygenates, and unknowns. Commercially available software may be used to provide this function, as well as calculation of other properties of petroleum liquids. See the caution in 5.3.

TABLE 3 Theoretical FID Relative Response Factors

Carbon No.	Saturated Paraffins	Unsaturated Paraffins	Saturated Naphthenes	Unsaturated Naphthenes	Aromatics
1	1.1207	-	-	-	-
2	1.0503	-	-	-	-
3	1.0268	0.9799	-	-	-
4	1.0151	0.9799	-	-	-
5	1.0080	0.9799	0.9799	0.9517	-
6	1.0034	0.9799	0.9799	0.9564	0.9095
7	1.0000	0.9799	0.9799	0.9598	0.9195
8	0.9975	0.9799	0.9799	0.9623	0.9271
9	0.9955	0.9799	0.9799	0.9642	0.9329
10	0.9940	0.9799	0.9799	0.9658	0.9376
11	0.9927	0.9799	0.9799	0.9671	0.9415
12	0.9916	0.9799	0.9799	0.9681	0.9447
13	0.9907	0.9799	0.9799	0.9690	0.9474
14	0.9899	0.9799	0.9799	0.9698	0.9497
15	0.9893	0.9799	0.9799	0.9705	0.9517

Oxygenates Relative Response Factors

	Lab 1	Lab 2	Lab 3	Lab 4	Ave.	Std. Dev.	%SD	Auto/Oil RRF
Methanol	3.0760	3.0477	2.9779	2.9230	3.0062	0.0691	2.30	3.0965
Ethanol	2.1888	2.0797	2.1755	2.0640	2.1270	0.0642	3.02	2.0953
t-Butanol	1.2975	1.3189	1.3312	1.2989	1.3116	0.0163	1.24	1.3368
MTBE	1.5279	1.5590	1.4860	1.5024	1.5188	0.0318	2.09	1.5016
ETBE	1.3848	1.3720	1.3804	1.3720	1.3773	0.0064	0.46	1.4032
TAME	1.3383	1.2993	1.3598	1.3340	1.3329	0.0250	1.88	1.3775

**DHA Method Oxygenate Linearity Cooperative Study - peak area
Laboratory 4**

Spl							Ave. RF	RRF
MeOH	0.0100	1.0100	5.0500	10.0200	20.0100	29.8300		
	0.4037	34.7643	174.8862	340.9069	717.4781	1046.1427		
	0.3599	33.8017	179.9043	353.4087	717.1507	980.1566		
ave.	0.3818	34.2830	177.3953	347.1578	717.3144	1013.1496		
RF	0.0262	0.0295	0.0285	0.0289	0.0279	0.0294	0.0288	2.9230
EtOH	0.0100	1.0000	5.0000	10.1000	20.1500	30.1800		
	0.2883	50.5190	237.7223	495.9717	967.7888	1526.2755		
	0.4095	46.7438	242.3003	500.4514	1007.0434	1537.3776		
ave.	0.3489	48.6314	240.0113	498.2116	987.4161	1531.8265		
RF	0.0287	0.0206	0.0208	0.0203	0.0204	0.0197	0.0204	2.0640
TBA	0.0099	0.9640	4.9692	9.9583	19.8768	29.7953		
	1.0363	77.5423	408.5969	757.2307	1546.4197	2241.0530		
	1.1869	72.7672	392.8649	775.5192	1550.7498	2346.4085		
ave.	1.1116	75.1548	400.7309	766.3749	1548.5847	2293.7307		
RF	0.0089	0.0128	0.0124	0.0130	0.0128	0.0130	0.0128	1.2989
MTBE	0.0100	0.9992	5.0362	9.9724	20.0248	30.0471		
	0.7645	66.0865	345.4606	713.3773	1332.2069	2041.1591		
	0.5890	65.8994	325.8215	679.7792	1348.4042	2052.4822		
ave.	0.6767	65.9929	335.6411	696.5783	1340.3055	2046.8206		
RF	0.0148	0.0151	0.0150	0.0143	0.0149	0.0147	0.0148	1.5024
ETBE	0.0099	0.9851	4.9255	9.8707	19.6724	29.5727		
	0.4527	69.3251	374.3939	732.8740	1537.9746	2144.9023		
	0.6242	72.7316	374.7065	695.3345	1462.4055	2173.4412		
ave.	0.5384	71.0283	374.5502	714.1042	1500.1901	2159.1718		
RF	0.0183	0.0139	0.0132	0.0138	0.0131	0.0137	0.0135	1.3720
TAME	0.0100	0.9997	4.9788	9.8883	19.1530	29.7144		
	0.3702	75.3456	363.7452	762.9970	1488.8626	2346.1907		
	0.0072	75.1503	380.0280	763.6254	1420.3514	2230.3657		
ave.	0.1887	75.2480	371.8866	763.3112	1454.6070	2288.2782		
RF	0.0530	0.0133	0.0134	0.0130	0.0132	0.0130	0.0132	1.3340
C6	8.5050	8.4750	8.4400	8.4525	8.4525	8.6950		
	890.3467	843.5383	836.5459	803.1739	843.6532	847.7344		
	847.7681	854.2333	840.8679	834.8488	841.6083	802.3011		
ave.	869.0574	848.8858	838.7069	819.0113	842.6307	825.0177		
RF	0.0098	0.0100	0.0101	0.0103	0.0100	0.0105	0.0101	1.0262
C7	8.5050	8.4750	8.4400	8.4525	8.4525	8.6950		
	893.5123	847.7426	868.0640	834.6944	880.9965	869.6032		
	846.4708	858.0901	862.0443	871.7571	882.1653	834.0419		
ave.	869.9916	852.9164	865.0541	853.2258	881.5809	851.8225		
RF	0.0098	0.0099	0.0098	0.0099	0.0096	0.0102	0.0099	1.0000
C8	8.5050	8.4750	8.4400	8.4525	8.4525	8.6950		
	889.7205	846.6591	877.1065	838.8929	890.2631	873.8851		
	839.2188	855.2006	862.7846	884.2601	895.4804	854.5747		
ave.	864.4697	850.9298	869.9455	861.5765	892.8718	864.2299		
RF	0.0098	0.0100	0.0097	0.0098	0.0095	0.0101	0.0098	0.9944
C9	8.5050	8.4750	8.4400	8.4525	8.4525	8.6950		
	883.5337	843.1968	870.7139	832.1808	883.3178	868.7531		
	829.0626	849.0969	854.1742	881.9661	889.9074	860.0512		
ave.	856.2982	846.1469	862.4440	857.0734	886.6126	864.4021		
RF	0.0099	0.0100	0.0098	0.0099	0.0095	0.0101	0.0099	1.0003

FIG. 10 Determination of Oxygenate Response—DHA Speciation Analysis




FIG. 11 Graphical Representation Determination of Oxygenate Response—DHA Speciation Analysis

17. Precision and Bias⁹

17.1 *Repeatability*—The difference in two test results obtained by the same operator with the same apparatus in a given laboratory under constant operating conditions on test samples taken from the same laboratory sample should, in the long run,

⁹ Supporting data is available from ASTM International Headquarters in the form of a research report. Request RR:D02-1518.

in the normal and correct operation of the test method not exceed the values given in Table 4 and Table A1.3 for the gasoline components.

17.2 *Reproducibility*—The difference between two single and independent measurements on test samples taken from the same bulk sample should, in the long run, in the normal and correct operation of the test method, not exceed the values given in Table 4 and Table A1.3 for the gasoline components.

TABLE 4 Repeatability and Reproducibility of DHA Determinations

NOTE 1—The following is a partial list of precision data that has been prepared by statisticians of CS94 in accordance with RR:D2-1007, and represents their best estimate of the cooperative study data. The complete precision data set appears in [Annex A1., Table A1.3](#).

NOTE 2—For each analyte to qualify for a precision statement, it must be present in at least six samples, and detected by at least six laboratories, at least once. The (repeatability standard deviation)/mean value for each analyte/sample combination must be less than or equal to 0.1, as per LOQ requirements which, while not a standard, is what CS94 is recommending.

NOTE 3—

Legend:

- r_{\min} = lower 95 % confidence limit of r_{est} ,
- r_{est} = repeatability estimate in percent of concentration,
- r_{\max} = upper 95 % confidence limit of r_{est} ,
- R_{\min}, R_{est} = for reproducibility,
- R_{\max}
- C_{\min} = lower concentration limit that $r_{\text{est}}, R_{\text{est}}$ is applicable, and
- C_{\max} = upper concentration limit that $r_{\text{est}}, R_{\text{est}}$ is applicable.

Component	Average RI	r_{\min}	r_{est}	r_{\max}	R_{\min}	R_{est}	R_{\max}	C_{\min}	C_{\max}
n-butane	400.00	6.8	9.9	13.9	15.3	32.4	59.1	1.02	3.75
i-pentane	477.45	5.9	7.2	8.7	8.5	14.8	23.8	2.48	13.38
Pentene-1	490.83	5.2	7.5	10.5	9.7	13.8	19	0.06	0.43
n-pentane	500.00	5.2	6.5	8.1	7.1	10.4	14.8	1.06	3.49
Cyclopentane	566.84	3.8	4.9	6.2	7	10.1	14	0.07	0.59
2,3-dimethylbutane	569.24	2.9	3.2	3.5	5.1	8.5	13.1	0.7	1.91
n-hexane	600.00	2	2.4	2.9	3.6	5.1	6.9	0.33	2.52
Methylcyclopentane	625.86	2.2	2.6	3.1	4.5	6.4	8.7	0.37	2.35
1-methylcyclopentene	648.71	1.9	2.7	3.7	7.9	8.7	9.6	0.17	0.82
Benzene	649.92	2.6	3.6	4.8	5.5	9	13.7	0.17	1.58
Cyclohexane	657.81	2.7	3.7	4.9	8.2	14.8	24.3	0.07	0.9
2-methylhexane	667.61	1.6	2.2	2.9	5.1	6.1	7.2	0.39	1.09
2,2,4-trimethylpentane	688.48	2.4	3.2	4.1	7.4	11.4	16.7	0.1	11.26
n-heptane	700.00	2.5	3.4	4.5	7.7	10.8	14.7	0.21	1.06
Methylcyclohexane	717.89	2.8	3.4	4	4.1	5.9	8.2	0.11	1.2
2,3,4-trimethylpentane	746.83	2.3	3.8	6	5.8	7.8	10.3	0.08	4.26
Toluene	751.77	1.9	2.7	3.8	10.8	13.5	16.5	1.99	10.34
2-methylheptane	764.14	3.5	4.9	6.6	4.8	6.1	7.5	0.15	0.63
n-octane	800.00	2.2	3.6	5.5	6.5	15.7	30.9	0.14	0.75
Ethylbenzene	854.65	2.2	3.2	4.4	7.2	10.6	14.9	0.62	2.62
1,3-dimethylbenzene	864.22	2.6	3.3	4.2	9.7	12.5	15.7	1.55	6.66
3-methyloctane	880.24	5.1	8.5	13	8.7	15.5	24.9	0.07	0.29
n-nonane	900.20	3.9	6.4	9.8	8.6	10.3	12.2	0.06	0.34
n-propylbenzene	946.33	2.8	5	8.1	7.6	11.9	17.7	0.21	0.77
1,4-methylethylbenzene	956.22	3.5	5.3	7.7	5.1	7.7	11.1	0.32	1.19
1,3,5-trimethylbenzene	961.92	3.7	5.5	7.7	5.4	8.3	12.1	0.39	1.21
2-methylnonane	971.77	6.5	10.6	16.2	17.5	25.9	36.6	0.03	0.19
1,2,4-trimethylbenzene	983.40	4.2	5.7	7.5	7.8	10.6	13.9	1.19	4.32
n-decane	1000.20	7.5	9.2	11.1	12.1	17.9	25.3	0.03	0.25
1,2,3-trimethylbenzene	1006.88	3.8	5.8	8.5	7.2	8.5	10	0.28	0.96
n-undecane	1100.00	8.6	13.9	21	24.4	40	61.2	0.03	0.18
1,2,3,5-tetramethylbenzene	1108.79	6.4	7.8	9.3	10.2	13.9	18.3	0.21	0.51
Naphthalene	1168.01	6.1	8.5	11.3	12.9	16.9	21.5	0.13	0.4
n-dodecane	1200.00	12.2	16.7	22.1	20.2	32.9	50	0.01	0.11
2-methylnaphthalene	1282.57	7.6	11.1	15.4	17.5	22.3	28	0.05	0.5

17.3 *Bias*—No information can be presented on the bias of the procedure in this test method for measuring hydrocarbon concentrations because no material having an accepted reference value is available.

18. Keywords

18.1 detailed hydrocarbon analysis; DHA; gas chromatography; hydrocarbons; open tubular column; oxygenates; PIONA; PONA

ANNEX
(Mandatory Information)
A1. PROCEDURE FOR ADJUSTING THE SELECTIVITY OF A DHA METHYL SILICONE OPEN TUBULAR COLUMN

A1.1 The successful application of this test method is highly dependent upon the selectivity of the column used. New 100 m × 0.25 mm 0.5 µm methyl silicone open tubular fused silica columns will likely not have sufficient selectivity for aromatics to function properly. Critical to the successful analysis of *reformulated* and oxygenated spark engine motor fuels is column inertness and component selectivity. Inertness of the primary 100 m column affects the retention and adsorption of the oxygenates such as alcohols and ethers, while selectivity for the aromatic compounds is controlled by the liquid phase. Until adequate commercial columns are available, it will be necessary to slightly increase the column selectivity, which is accomplished by the addition of a short precolumn containing a moderately selective liquid phase.

A1.2 Prior to making any precolumn additions to the 100-m methyl silicone capillary column, determine that the main column meets the column specifications outlined in 6.4.1 and determined in Section 9. Section 9 describes the preliminary evaluation of the 100 m methyl silicone capillary column, using a 35 °C isothermal analysis to determine the basic column characteristics of efficiency, retention factor, inertness, and selectivity. Figs. A1.1-A1.3 provide examples of the column quality specification determinations. These determinations may also be made with a precolumn attached, since the precolumn has little if any affect on the results. Fig. A1.4 illustrates that the addition of different lengths of precolumn has negligible influence on the retention characteristics of oxygenated compounds. Poor peak shape and resolution of these oxygenates cannot be corrected by the addition of the precolumn. Tailing peaks may also be the result of an active injector liner or packing material, or both, in the injector liner. An increase in retention of the oxygenates is likely due to column activity. The relative position of the oxygenates to the hydrocarbons is dependent upon column temperature, thus a faulty column oven temperature control could also result in shifted peaks.

A1.3 When necessary, a precolumn is added to the primary 100 m column to adjust the column selectivity for aromatic compounds. Precolumns that have been used successfully are variable lengths of 0.25 mm internal diameter fused silica open tubular column containing a 1.0 µm film thickness of 5 % phenyl methyl silicone. The film thickness is likely not critical, only the total amount of phase. Lengths ranging from 1 m to more than 3 m have been necessary to provide sufficient selectivity, depending on the initial selectivity of the methyl

silicone column used. One metre of 1.0 µm precolumn is equivalent to a 100 m column with 0.5 µm of 0.1 % phenyl methyl silicone liquid phase.

A1.4 Figs. A1.5-A1.8 illustrate the resolution of the methylcyclopentene-1 and benzene pair with a new column and one, two, and three metres of precolumn. The key segment of the chromatogram is expanded to better illustrate the resolution of this component pair.

A1.5 The preliminary evaluation of the 100 m column will provide the user with information regarding the initial length of precolumn with which to start the tuning process. Dependent upon the methylcyclopentene-1 and benzene resolution, an initial precolumn of between 1 and 4 m is selected; which ever provides a resolution greater than 1.2.

A1.6 The final tuning will consist of reducing the precolumn length, probably in increments of 0.25 m, until the proper resolution is achieved between 2,3,3-trimethylpentane and toluene, and 1,4-dimethylbenzene and 2,3-dimethylheptane; using the actual analysis temperature conditions.

A1.7 Fig. A1.9 illustrates graphically the effect of different lengths of precolumn, attached to the same 100 m column. The key component separations are shown. These analyses were made using the conditions given in Table 2. In this case, the use of the 1.25 m precolumn provides the best compromise for the three key separations.

A1.8 Fig. A1.10 illustrates the use of different lengths of precolumn to achieve the specified selectivity for three different 100 m columns. The final precolumn length will provide adequate resolution of all three of the key separations.

A1.9 Figs. A1.11-A1.17 illustrate DHA analyses.

A1.10 Tables A1.4-A1.9 show comparisons between this test method and other methods for several compound types. Multidimensional PIONA is included since it tends to give reasonable peak compound type groupings for total olefins, total paraffins, and total naphthenes. The differences for benzene and toluene among the indicated methods are well within the reproducibilities of the methods. The sample numbers refer to the interlaboratory cooperative study samples. It should be noted that the interlaboratory cooperative study samples included only spark ignition fuels and different results may be obtained with pure blending components.

TABLE A1.1 DHA Component Data

NOTE 1—These data consist of the current physical constants used in the cooperative study. The average RI are those accumulated in a ruggedness test of the *tuning* process. The RFA and CCF gasoline data are the averages determined in the cooperative study. RFA is an industry average regular gasoline and CCF is a California Certification Fuel (reformulated gasoline).

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
Methane	100.00	1.121	16.043	0.2600						
Ethylene	178.10	1.050	28.054	0.3000						
Ethane	200.00	1.051	30.070	0.3399						
Propylene	284.00	1.030	42.081	0.5053				7.173	293.43	0.000
Propane	300.00	1.027	44.097	0.5005	7.270	300.32	0.003	7.266	299.79	0.003
i-Butane	366.15	1.015	58.124	0.5572	8.266	365.46	0.088	8.262	365.29	0.078
Methanol	378.82	3.148	32.040	0.7914				8.506		0.021
Butene-1	390.72	0.980	56.108	0.5951				8.893	390.31	0.019
Isobutylene	391.51	0.980	56.108	0.5951						
1,3-Butadiene	394.93	0.980	54.092	0.6211						
n-Butane	400.00	1.015	58.124	0.5788	9.195	400.00	4.637	9.193	400.00	1.201
Vinyl acetylene	409.00	1.100	54.090	0.6500						
t-Butene-2	412.09	0.980	56.108	0.6042	9.441	411.72	0.002	9.567	412.10	0.013
2,2-Dimethylpropane	415.10	1.008	72.151	0.5910	9.670	415.09	0.036	9.666	415.05	0.017
c-Butene-2	427.74	0.980	56.108	0.6213	9.983	427.70	0.004	10.128	427.73	0.018
1,2-Butadiene	450.00	0.945	54.092	0.6520						
Ethanol	455.33	2.193	46.070	0.7890	11.063	452.52	0.006			
3-Methylbutene-1	460.84	0.980	70.135	0.6272	11.670	460.81	0.010	11.665	460.76	0.020
O ₁	469.00	0.980	70.135	0.6300	12.030	470.03	0.005			
O ₂	474.00	0.980	70.135	0.6300						
i-Pentane	477.45	1.008	72.151	0.6196	12.653	477.15	4.773	12.650	477.13	7.163
Acetone ^A	477.55	1.850	58.080	0.7899				12.649		0.134
1,4-Pentadiene	481.18	0.952	68.119	0.6607				13.464	482.77	0.005
?	483.00				13.122	486.32	0.014	12.675	482.80	0.003
Butyne-2	488.00	0.945	54.092	0.6910						
Pentene-1	490.83	0.980	70.135	0.6405	13.616	490.86	0.152	13.613	490.85	0.091
i-Propanol	493.38	1.400	60.110	0.8000						
2-Methylbutene-1	496.66	0.980	70.135	0.6504	14.074	496.73	0.334	14.071	496.72	0.185
n-Pentane	500.00	1.008	72.151	0.6262	14.341	500.00	3.627	14.339	500.00	1.094
Isoprene	506.02	0.952	68.119	0.6809	14.666	506.00	0.013	14.664	505.98	0.009
?	508.00				14.644	508.07	0.003			
t-Pentene-2	510.56	0.980	70.135	0.6482	14.917	510.41	0.653	14.916	510.41	0.285
3,3-dimethylbutene-1	516.79	1.050	70.135	0.6500	15.277	516.60	0.011	15.141	516.58	0.004
c-Pentene-2	519.53	0.980	70.135	0.6556	15.439	519.25	0.378	15.438	519.28	0.160
t-Butanol	521.64	1.154	74.120	0.7887				15.468	522.58	0.065
?	522.40							16.198	522.66	0.038
2-Methylbutene-2	524.92	0.980	70.135	0.6623	15.765	524.49	1.100	15.763	524.51	0.461
1t,3-Pentadiene	527.97	0.952	68.119	0.6760	15.960	527.59	0.022	15.956	527.56	0.015
3-Methylbutadiene-1,2	535.00	0.952	68.120	0.6500						
Cyclopentadiene	538.05	0.938	67.100	0.6500	16.478	537.58	0.004	16.475	537.57	0.003
2,2-Dimethylbutane	540.54	1.004	86.178	0.6491	16.779	539.78	1.102	16.776	539.75	1.106
1c,3-Pentadiene	541.90	0.952	68.119	0.6910						
?	543.00				16.895	543.44	0.006			
O ₅	547.70	1.020	70.135	0.6500						
O ₆	549.70	1.020	70.135	0.6500						
Cyclopentene	557.21	0.952	68.119	0.7720	18.026	556.65	0.160	18.025	556.67	0.070
n-Propanol	560.00	1.400	60.110	0.8035						
4-Methylpentene-1	562.02	0.980	84.162	0.6673	18.411	561.26	0.050	18.402	561.42	0.021
3-Methylpentene-1	562.81	0.980	84.162	0.6637	18.468	562.21	0.083	18.469	562.26	0.032
Cyclopentane	566.84	0.980	70.135	0.7454	18.811	566.40	0.216	18.813	566.45	0.052
2,3-Dimethylbutane	569.24	1.004	86.178	0.6616	19.003	568.67	1.723	19.001	568.69	1.655
Methyl-t-butylether	570.65	1.417	88.150	0.7405				19.110	570.03	11.282
4-Methyl-c-pentene-2	571.00	0.980	84.162	0.6741	19.154	570.47	0.113			
2,3-Dimethylbutene-1	572.67	0.980	84.162	0.6830	19.306	572.01	0.048	19.520	572.52	0.028
2-Methylpentane	573.70	1.004	86.178	0.6531	19.388	573.19	5.145	19.389	573.23	3.967
4-Methyl-t-pentene-2	575.47	0.980	84.162	0.6736	19.542	574.94	0.167	19.546	575.03	0.083
O ₈	578.00	0.980	84.162	0.6736	20.042		0.002	19.893		0.002
2-Methyl-1,4-pentadiene	579.00	0.980	82.146	0.6940	20.078		0.002			
?	581.00							20.002		0.002
1,5-Hexadiene	581.90	0.980	82.146	0.6923	20.123	581.47	0.002	20.210		0.011
?	583.90				20.250	583.99	0.002			
3-Methylpentane	585.52	1.004	86.178	0.6643	20.477	585.25	2.589	20.476	585.25	2.189
2-Methylpentene-1	590.19	0.980	84.162	0.6848	20.933	590.01	0.241	20.934	590.05	0.103
Hexene-1	591.06	0.980	84.162	0.6780	21.021	590.91	0.127	21.021	590.94	0.059
O ₁₁	592.00	0.980	84.162	0.6780						
i-Butanol	593.50	1.337	74.120	0.8030						
?	596.00				21.474	596.35	0.007			
1c/t,4-Hexadiene	597.14	0.980	84.146	0.7000	21.571	597.22	0.004			
2-Ethylbutene-1	598.95	0.980	84.162	0.6944						

TABLE A1.1 *Continued*

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
n-Hexane	600.00	1.004	86.178	0.6594	21.937	600.00	2.598	21.935	600.00	1.057
Diisopropylether	601.90	1.100	102.180	0.7241						
t-Hexene-3	602.83	0.980	84.162	0.6821	22.169	602.83	0.191	22.170	602.86	0.080
c-Hexene-3	603.56	0.980	84.162	0.6847	22.258	603.60	0.063	22.234	603.65	0.028
t-Hexene-2	605.44	0.980	84.162	0.6827	22.382	605.40	0.347	22.383	605.43	0.157
2-Methylpentene-2	607.86	0.980	84.162	0.6912	22.583	607.77	0.462	22.584	607.80	0.193
4-Methylcyclopentene	609.00				21.685	608.90	0.113	22.589	608.65	0.047
3-Methyl-c-pentene-2	610.54	0.980	84.162	0.6980	22.816	610.51	0.240	22.817	610.54	0.102
3-Methylcyclopentene	611.61	0.980	82.146	0.7622	22.920	611.74	0.055	22.921	611.77	0.025
O ₁₃	613.08	0.980	84.162	0.6920						
c-Hexene-2	614.67	0.980	84.162	0.6920	23.171	614.60	0.194	23.172	614.63	0.088
O ₁₄	617.06	0.980	84.162	0.6920	23.007	617.10	0.004	23.088	617.08	0.002
Ethyl-t-butylether	619.00	1.342	102.180	0.7519						
3,3-Dimethylpentene-1	620.91	0.980	98.189	0.7019	23.722	620.77	0.371	23.723	620.80	0.158
3-Methyl-t-pentene-2	622.11	0.980	84.162	0.7023	23.603	622.17	0.006	23.480	622.19	0.003
2-Butanol	622.40	0.980	74.120	0.8080						
4-4-Dimethyl-t-pentene-2	623.10	0.980	98.189	0.6936						
2,2-Dimethylpentane	624.17	1.000	100.205	0.6738	24.025	624.11	0.084	24.024	624.12	0.128
Methylcyclopentane	625.86	0.980	84.162	0.7486	24.189	625.88	0.963	24.190	625.91	0.355
Cyclic diolefin or triolefin	627.00	0.957	82.140	0.7092						
2,4-Dimethylpentane	630.60	1.000	100.205	0.6727	24.622	630.47	1.036	24.623	630.48	2.437
2,3,3-Tirmethylbutene-1	631.00	0.980	98.189	0.7092						
Cyclic diolefin or triolefin	632.90	0.957	82.140	0.7092	24.846	632.82	0.010	24.844	632.80	0.008
?	634.20				25.835	634.14	0.009			
2,2,3-Trimethylbutane	634.86	1.000	100.205	0.6901	25.049	634.91	0.031	25.050	634.93	0.038
?	636.30				25.186	636.30	0.007	25.091	636.33	0.004
Cyclic diolefin or triolefin	638.30	0.957	82.140	0.7092	25.380	639.26	0.008	25.378	638.27	0.005
O ₁₇	641.97	0.980	84.160	0.7039	25.745	641.92	0.005	25.622	642.14	0.002
3,4-Dimethylpentene-1	642.87	0.980	98.189	0.7022	25.846	642.92	0.014	25.845	642.92	0.008
4,4-Dimethyl-c-pentene-2	646.65	0.980	98.189	0.7039	26.223	646.57	0.024	26.224	646.61	0.011
2,4-Dimethylpentene-1	647.67	0.980	98.189	0.6988	26.332	647.63	0.021	26.334	647.65	0.011
Diolefin	647.70	0.957	82.140	0.6988						
1-Methylcyclopentene	648.71	0.957	82.146	0.7795	26.443	648.69	0.374	26.444	648.72	0.180
Benzene	649.92	0.910	78.114	0.8789	26.580	649.98	1.969	26.579	649.99	1.242
3-Ethylpentene-1	650.00	0.980	98.189	0.7005						
n-Butanol ^A	650.02	1.295	74.120	0.8000						
3-Methylhexene-1	650.95	0.980	98.189	0.6959	26.420	651.56	0.029	26.434	651.55	0.015
2-Methyl-c-hexene-3	652.60	0.980	98.189	0.6980	27.081	652.56	0.018	27.059	652.59	0.009
3,3-dimethylpentane	654.43	1.000	100.205	0.6932	27.057	654.47	0.094	27.055	654.46	0.139
5-Methylhexene-1	655.56	0.980	98.189	0.6965	27.198	655.83	0.031	26.985	656.10	0.016
?	656.93				28.233	656.74	0.014	27.752	656.78	0.007
Cyclohexane	657.81	0.980	84.162	0.7785	27.440	657.97	0.225	27.445	658.05	0.050
2-Methyl-t-hexene-3	661.03	0.980	98.189	0.6941	27.763	660.87	0.057	27.766	660.91	0.027
Diolefin (hexadiene)	661.30	0.980	98.189	0.6941	27.946	661.74	0.007	27.794		0.003
2-Ethyl-3-methylbutene-1	662.60	0.980	98.189	0.7135	27.941	662.47	0.018	27.944	662.51	0.009
4-Methylhexene-1	663.81	0.980	98.189	0.7030	28.087	663.77	0.040	28.089	663.80	0.019
4-Methyl-t/c-hexene-2	666.23	0.980	98.189	0.7040	28.357	666.13	0.107	28.361	666.18	0.051
2-methylhexane	667.61	1.000	100.205	0.6786	28.510	667.45	1.342	28.518	667.54	1.236
2,3-Dimethylpentane	668.84	1.000	100.205	0.6951	28.663	668.79	1.635	28.673	668.88	4.375
5-Methyl-t-hexene-2	669.80	0.980	98.189	0.6971						
1,1-Dimethylcyclopentane	671.25	0.980	98.189	0.7545	28.958	671.32	0.045	28.961	671.35	0.042
t-Amylmethylether	672.48	1.318	102.180	0.7517				28.973		0.002
Cyclohexene	673.69	0.980	82.146	0.8110	29.254	673.82	0.058	29.255	673.84	0.032
3-Methylhexane	675.89	1.000	100.205	0.6871	29.496	675.82	1.449	29.497	675.83	1.450
1,6-Heptadiene	677.40	0.980	98.190	0.7500						
3,4-Dimethyl-c-pentene-2	679.46	0.980	98.189	0.7180	29.935	679.42	0.043	29.938	679.46	0.021
5-Methyl-c-hexene-2	680.00	0.980	98.189	0.7060						
1c,3-Dimethylcyclopentane	681.68	0.980	98.189	0.7448	30.225	681.78	0.261	30.228	681.82	0.114
1t,3-Dimethylcyclopentane	684.37	0.980	98.189	0.7488	30.567	684.51	0.228	30.572	684.54	0.104
3-Ethylpentane	685.98	1.000	100.205	0.6981	30.751	685.96	0.202	30.757	686.01	0.169
1t,2-Dimethylcyclopentane	687.07	0.980	98.189	0.7514	30.911	687.21	0.185	30.921	687.30	0.085
2,2,4-Trimethylpentane	688.48	0.998	114.232	0.6919	31.086	688.57	3.273	31.115	688.81	9.481
Heptene-1	688.60	0.980	98.189	0.6970						
2-Ethylpentene-1	689.58	0.980	98.189	0.6970	31.231	689.58	0.059			
1,5-Heptadiene	691.60	0.980	93.168	0.7500						
O ₂₅	692.89	0.980	98.189	0.6900	31.502	692.93	0.007	31.422	692.95	0.004
3-Methyl-c-hexene-3	694.82	0.980	98.189	0.7181	31.912	694.87	0.070	31.913	694.88	0.034
?	696.80							31.826	696.87	0.004
t-Heptene-3	698.39	0.980	98.189	0.7026	32.392	698.43	0.250	32.393	698.44	0.126
n-Heptane	700.00	1.000	100.205	0.6837	32.605	700.00	1.164	32.604	700.00	0.996
c-Heptene-3	701.00	0.980	98.189	0.7028						
2-Methyl-2-hexene	701.30	0.980	98.189	0.7126						
3-Methyl-c-hexene-2	702.30	0.980	98.189	0.7126	32.916	702.07	0.283	32.917	702.08	0.139

TABLE A1.1 *Continued*

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
3-Methyl-t-hexene-3	702.99	0.980	98.189	0.6941	33.064	703.05	0.103	33.065	703.06	0.049
t-Heptene-2	704.58	0.980	98.189	0.7057	33.306	704.63	0.124	33.307	704.64	0.063
3-Ethylpentene-2	705.96	0.980	98.189	0.7249	33.526	706.06	0.065	33.527	706.07	0.030
c-Heptene-2	708.82	0.980	98.189	0.7116	33.974	708.92	0.221	33.990	709.04	0.129
3-Methyl-t-hexene-2	709.50	0.980	98.189	0.7188						
O ₂₈	710.53	0.980	98.189	0.7188						
2,3-Dimethylpentene-2	712.07	0.980	98.189	0.7322	34.488	712.18	0.123	34.488	712.17	0.059
3-Ethylcyclopentene	713.22	0.980	96.173	0.7830	34.789	713.45	0.010	34.562	713.41	0.004
O ₂₉	715.67	0.980	98.189	0.7190	35.083	715.85	0.018	35.084	715.86	0.009
1c,2-Dimethylcyclopentane	717.13	0.980	98.189	0.7322	35.329	717.35	0.138	35.331	717.36	0.060
Methylcyclohexane	717.89	0.980	98.189	0.7694	35.451	718.09	0.380	35.452	718.09	0.139
O ₃₀	719.00	0.980	98.189	0.7322	35.359	720.75	0.079			
2,2-Dimethylhexane	720.70	0.998	114.232	0.6953	35.884	720.70	0.117	35.854	720.51	0.099
1,1,3-Trimethylcyclopentane ^A	720.72	0.980	112.216	0.7482						
O ₃₂	721.00	0.980	98.189	0.7322						
O ₃₃	722.00	0.980	112.216	0.7322						
O ₃₄	723.00	0.980	112.216	0.7322						
O ₃₅	724.35	0.980	98.189	0.7322	36.531	724.50	0.013	36.398	724.50	0.005
O ₃₆	726.26	0.980	98.189	0.7322	36.872	726.47	0.010	36.652	725.97	0.010
?	727.00							36.637	726.09	0.003
Ethylcyclopentane	728.90	0.980	98.189	0.7664	37.342	729.16	0.146	37.381	729.09	0.072
2,5-Dimethylhexane	730.05	0.998	114.232	0.6935	37.522	730.17	0.422	37.523	730.17	0.693
2,2,3-Trimethylpentane	730.90	0.998	114.232	0.7160	37.775	730.96	0.064	37.682	731.06	0.188
2,4-Dimethylhexane	731.84	0.998	114.232	0.7003	37.846	731.99	0.697	37.848	731.98	1.056
?	733.53				38.081	733.63	0.011	37.823		0.007
O ₃₇	735.18	0.980	98.189	0.7322	38.047	735.41	0.004	38.027	735.34	0.003
1c,2t,4-Trimethylcyclopentane	737.11	0.980	112.216	0.7634	38.824	737.36	0.102	38.825	737.35	0.046
3,3-Dimethylhexane	738.39	0.998	114.232	0.7100	39.052	738.59	0.077	39.053	738.59	0.075
O ₃₈	740.43	0.980	98.189	0.7322	39.432	740.61	0.005	38.887	740.62	0.004
?	742.18				39.825	742.27	0.007	39.387	742.53	0.003
?	743.20				40.364	743.28	0.024	40.227	743.50	0.015
?	743.80				39.701	744.00	0.033	39.868	743.90	0.014
1t,2c,3-Trimethylcyclopentane	744.21	0.980	112.216	0.7704	40.162	744.46	0.077	40.163	744.44	0.033
O ₃₉	745.34	0.980	98.189	0.7322				40.019	744.72	0.005
2,3,4-Trimethylpentane	746.83	0.998	114.232	0.7190	40.667	747.06	0.862	40.678	747.11	2.585
I1	747.91	0.998	114.232	0.7190	40.874	748.12	0.195	40.876	748.11	0.093
O ₄₀	749.37	0.980	98.189	0.7322	41.157	749.56	0.058	41.160	749.54	0.032
2,3,3-Trimethylpentane	750.84	0.998	114.232	0.7262	41.539	751.10	0.525	41.470	751.10	1.716
Toluene	751.77	0.920	92.143	0.8670	41.666	752.08	6.421	41.688	752.18	8.999
O ₄₁	752.20	0.980	112.220	0.7322						
O ₄₂	753.63	0.980	112.220	0.7322	42.037	753.73	0.030	42.362	753.75	0.014
?	754.63				42.054	754.65	0.009	42.183	754.77	0.020
O ₄₃	755.33	0.980	112.220	0.7322	42.351	755.48	0.049	42.334	755.36	0.031
2,3-Dimethylhexane	757.87	0.998	114.232	0.7121	42.890	758.08	0.508	42.898	758.11	0.812
2-Methyl-3-ethylpentane	759.04	0.998	114.232	0.7121	43.139	759.28	0.060	43.149	759.31	0.062
1,1,2-Trimethylcyclopentane	760.33	0.980	112.216	0.7725	43.380	760.44	0.045	43.379	760.42	0.025
O ₄₄	761.73	0.980	112.220	0.7322	43.709	761.99	0.076	43.712	761.97	0.041
O ₄₅	762.20	0.980	112.220	0.7322						
O ₄₆	763.00	0.980	112.220	0.7322						
2-Methylheptane	764.14	0.998	114.232	0.6979	44.199	764.29	0.831	44.198	764.26	0.571
2-ethylhexene-1 ^A	764.20	0.980	112.220	0.7650						
4-Methylheptane	765.62	0.998	114.232	0.7046	44.521	765.78	0.362	44.521	765.75	0.266
3-Methyl-3-ethylpentane	766.62	0.980	114.232	0.7121	44.753	766.83	0.084	44.750	766.80	0.104
3,4-Dimethylhexane	767.18	0.998	114.232	0.7192	44.865	767.35	0.086	44.867	767.33	0.114
1c,2c,4-Trimethylcyclopentane	768.95	0.980	112.216	0.7620	45.430	769.91	0.090	45.427	769.88	0.041
1c,3-Dimethylcyclohexane	769.80	0.980	112.216	0.7625						
3-Methylheptane	771.78	0.998	114.232	0.7058	45.880	771.92	0.911	45.877	771.88	0.651
1c,2t,3-Trimethylcyclopentane	772.98	0.980	112.216	0.7704	56.135	773.05	0.291	46.127	772.98	0.185
3-Ethylhexane	773.76	0.998	114.232	0.7136	46.360	774.03	0.055	46.362	774.01	0.022
1t,4-Dimethylcyclohexane	774.89	0.980	112.216	0.7625	46.689	775.16	0.059	46.621	775.15	0.024
?	775.65				46.439	775.62	0.009			
1,3-Octadiene	777.16	0.980	110.200	0.7650	47.117	777.33	0.011	46.298	777.29	0.006
O ₄₈	778.50	0.980	112.220	0.7322	46.756	779.08	0.004	46.542	779.02	0.003
1,1-Dimethylcyclohexane	780.48	0.980	112.216	0.7809	47.922	780.76	0.009	47.413	780.75	0.005
2,2,5-Trimethylhexane	782.93	0.996	128.259	0.7072	48.473	783.08	0.470	48.473	783.04	0.740
3c-Ethylmethylcyclopentane	784.35	0.980	112.216	0.7670	48.831	784.57	0.130	48.833	784.53	0.060
2,6-Dimethylheptene-1	785.55	0.980	126.240	0.7196	49.152	785.64	0.018	49.046	785.66	0.010
3t-Ethylmethylcyclopentane	786.55	0.980	112.216	0.7670	49.366	786.75	0.081	49.369	786.74	0.036
2t-Ethylmethylcyclopentane	787.86	0.980	112.216	0.7690	49.679	788.03	0.071	49.682	788.00	0.036
Octene-1 ^A	787.87	0.980	112.220	0.7650						
1,1-Methylethylcyclopentane	788.78	0.980	112.216	0.7809	49.894	788.90	0.028	49.896	788.89	0.013
?	789.88				49.436	789.88	0.013	50.166	789.96	0.013
2,2,4-Trimethylhexane	790.75	0.996	128.259	0.7392	50.384	790.88	0.046	50.386	790.86	0.020

TABLE A1.1 *Continued*

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
1t,2-Dimethylcyclohexane	792.77	0.980	112.216	0.7760	50.911	792.96	0.096	50.915	792.94	0.045
t-Octene-4	794.21	0.980	112.216	0.7185	51.252	794.31	0.052	51.259	794.31	0.029
3,5,5-Trimethylhexene-1	795.00	0.980	126.240	0.7196	50.669		0.002			
t-Octene-3	796.00	0.980	112.216	0.7196						
1c,2c,3-Trimethylcyclopentane	797.25	0.980	112.216	0.7792	52.033	797.34	0.151	52.042	797.34	0.080
1t,3-Dimethylcyclohexane	798.80	0.980	112.216	0.7760	52.443	798.90	0.033	52.450	798.89	0.017
n-Octane	800.00	0.998	114.232	0.7025	52.733	800.00	0.811	52.740	800.00	0.502
1c,4-Dimethylcyclohexane	801.05	0.980	112.216	0.7828						
3,3-Dimethylheptene-1	802.50	0.980	126.240	0.7196						
Octene-2	804.40	0.980	112.216	0.7196	53.872	804.52	0.065	53.885	804.54	0.035
O ₅₀	805.50	0.980	112.216	0.7196						
I ₂	806.39	0.996	128.259	0.7300	54.389	806.51	0.112	54.406	806.56	0.052
?	807.00							54.600	807.29	0.020
i-Propylcyclopentane	808.06	0.980	112.216	0.7765	54.822	808.20	0.074	54.840	808.23	0.035
2,4,4-Trimethylhexane	808.50	0.996	128.259	0.7392						
O ₅₂	810.62	0.980	126.240	0.7196	55.508	810.81	0.011	55.545	810.84	0.006
O ₅₃	813.47	0.980	126.240	0.7196	56.268	813.67	0.035	56.295	813.72	0.016
N ₁	815.02	0.980	112.216	0.7800	56.655	815.09	0.012	55.395	814.95	0.007
?	815.60				56.673		0.006			
2,2,3,4-Tetramethylpentane	816.45	0.996	128.259	0.7389	57.221	816.59	0.009	56.554	816.50	0.005
2,3,4-Trimethylhexane	818.10	0.996	128.259	0.7392	57.536	818.33	0.081	57.567	818.38	0.127
N ₂	819.93	0.980	112.216	0.7800	58.170	820.20	0.034	58.093	820.25	0.014
?	820.85				58.049	821.28	0.014	57.612	821.13	0.007
?	821.10				59.376	820.70	0.014	59.939	821.10	0.009
N ₃	822.29	0.980	112.216	0.7800	58.668	822.41	0.039	58.725	822.55	0.022
2,3,3-Trimethylhexene-1	824.74	0.980	126.240	0.6826	59.404	824.98	0.017	59.582	825.08	0.010
?	825.00									
1c,2-Dimethylcyclohexane	826.48	0.980	112.216	0.7962	59.179	826.84	0.023			
1,3,5-Trimethylhexane	827.51	0.996	128.259	0.7219	60.088	827.38	0.122	60.141	827.50	0.084
?	828.95				60.649	828.50	0.006			
?	829.20							59.422		0.002
2,2-Dimethylheptane	829.76	0.996	128.259	0.7105	59.233	829.20	0.012	59.294	829.15	0.005
?	831.80							60.231		0.004
1,1,4-Trimethylcyclohexane	832.56	0.980	126.243	0.7722	61.689	832.81	0.104	61.744	832.94	0.049
N ₄	834.07	0.980	112.216	0.7800	62.028	833.97	0.021	62.098	834.13	0.014
?	834.40				60.862		0.013	60.851		0.008
2,2,3-Trimethylhexane	834.96	0.996	128.259	0.7153	62.399	835.22	0.074	62.472	835.37	0.043
2,4-Dimethylheptane	836.47	0.996	128.259	0.7153	62.878	836.79	0.011	61.246	836.60	0.005
4,4-Dimethylheptane	838.68	0.996	128.259	0.7153	63.454	838.68	0.048	63.530	838.83	0.024
Ethylcyclohexane	840.20	0.980	112.216	0.7839	62.675		0.004	62.624		0.003
n-Propylcyclopentane	841.38	0.980	112.216	0.7763	64.307	842.00	0.008	62.897	841.45	0.004
1c,3c,5-trimethylcyclohexane ^A	841.40	0.980	126.243	0.7697						
2,5-Dimethylheptane	842.63	0.996	128.259	0.7167	64.762	842.88	0.208	64.855	843.09	0.132
3,3-Dimethylheptane	843.96	0.996	128.259	0.7256	65.189	844.25	0.059	65.283	844.44	0.032
3,5-Dimethylheptane	845.02	0.996	128.259	0.7225	65.042	844.55	0.015	64.178		0.002
?	845.60							64.526		0.002
2,6-Dimethylheptane	846.47	0.996	128.259	0.7089	65.987	846.73	0.023	64.434	846.51	0.007
?	847.00							64.667		0.002
1,1,3-Trimethylcyclohexane	848.43	0.980	126.243	0.7870	66.612	848.67	0.024	66.718	848.90	0.011
2,4-Dimethylheptene-1	849.43	0.980	126.240	0.6826	66.462	848.89	0.006	65.309	849.56	0.005
N ₇	850.89	0.980	112.216	0.7800	66.993	850.52	0.008	65.931	851.33	0.005
N ₈	852.36	0.980	112.216	0.7800	68.017	852.57	0.017	67.247	852.64	0.010
N ₁₀	853.04	0.980	126.240	0.7800						
Ethylbenzene	854.65	0.927	106.168	0.8670	68.687	854.92	3.131	68.809	855.18	2.395
N ₁₁	854.70	0.980	126.240	0.7800	67.693		0.015			
1c,2t,4t-Trimethylcyclohexane	856.34	0.980	126.243	0.7800	69.261	856.61	0.060	69.377	857.03	0.037
I ₃	858.51	0.996	128.259	0.7300	70.138	858.77	0.016	68.290	858.74	0.011
?	859.00							68.711		0.002
2-Methyloctene-1	859.80	0.980	126.240	0.6826	70.172	859.99	0.021	70.169	859.87	0.015
?	860.50							69.131		0.002
I ₄	860.89	0.996	128.259	0.7300	70.240	860.78	0.011	69.125	861.19	0.008
?	861.40				69.536		0.005	69.853		0.002
2-Methyloctene-2	862.14	0.980	126.240	0.6826	71.310	862.00	0.014	69.565	862.44	0.009
N ₁₂	863.00	0.980	126.243	0.7800	72.374	863.22	0.017	70.111		0.005
N ₁₃	863.77	0.980	126.243	0.7800						
1,3-Dimethylbenzene	864.22	0.927	106.168	0.8642	72.171	864.48	5.181	72.190	864.82	3.649
1,4-Dimethylbenzene	865.20	0.927	106.168	0.8610	72.389	865.45	2.299	72.556	865.82	1.698
2,3-Dimethylheptane	866.02	0.996	128.259	0.7260	71.154	866.59	0.111	72.052	866.55	0.088
3,4-Dimethylheptane	867.94	0.996	128.259	0.7314	73.307	868.00	0.040	73.490	868.38	0.025
?	868.00							72.160		0.002
3,4-Dimethylheptane	868.78	0.996	128.259	0.7314	73.584	868.75	0.041	73.782	869.18	0.023
N ₁₄	869.70	0.980	126.243	0.7800	74.035	869.96	0.018	72.336	870.12	0.009
I ₅	870.95	0.996	128.259	0.7300	74.478	871.15	0.073	74.658	871.53	0.036

TABLE A1.1 *Continued*

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
4-Ethylheptane	872.73	0.996	128.259	0.7202	75.147	872.96	0.016	75.560	873.30	0.010
4-Methyloctane	873.81	0.996	128.259	0.7202	75.557	874.05	0.237	75.727	874.35	0.104
2-Methyloctane	874.76	0.996	128.259	0.7134	75.918	874.99	0.298	76.089	875.30	0.128
?	875.00							75.116		0.002
N ₁₅	876.00	0.980	126.243	0.7800	76.402	876.26	0.023	74.595	876.35	0.010
1c,2t,3-Trimethylcyclohexane	877.98	0.980	126.243	0.7580	76.601	878.00	0.016	76.726	878.30	0.007
?	878.00				77.313	878.27	0.040	75.982		0.005
3-Ethylheptane	879.11	0.996	128.259	0.7265	77.373	879.55	0.075	77.742	879.58	0.031
3-Methyloctane	880.24	0.996	128.259	0.7205	78.034	880.47	0.336	78.185	880.70	0.144
?	881.04				80.286	881.32	0.016	76.824	881.25	0.004
1c,2t,4c-Trimethylcyclohexane	881.67	0.980	126.243	0.7722	78.628	881.99	0.019	78.869	882.43	0.020
1,1,2-Trimethylcyclohexane	882.78	0.980	126.243	0.8000	78.917	882.72	0.015			
1,2,-Dimethylbenzene	883.47	0.927	106.168	0.8802	79.328	883.76	2.652	79.453	883.89	1.784
?	884.87				77.949	884.74	0.015	77.734	884.66	0.004
I6	885.34	0.996	128.259	0.7300	80.135	885.49	0.022	80.257	885.89	0.023
?	885.88				79.109		0.008			
I7	886.38	0.996	128.259	0.7300	80.474	886.61	0.062	80.608	886.75	0.139
N ₁₈	887.87	0.980	126.243	0.7800	81.060	888.07	0.038	81.621	888.18	0.017
N ₁₉	888.36	0.980	126.243	0.7800	81.279	888.60	0.030	81.398	888.69	0.011
Nonene-1	889.00	0.980	126.240	0.7684						
?	889.40				81.509	889.16	0.040	80.759	889.21	0.016
I8	889.78	0.996	128.259	0.7300	81.888	890.09	0.024	81.476	889.94	0.011
N ₂₀	890.51	0.980	126.243	0.7800	85.070	890.70	0.111	86.286	890.45	0.010
I9	891.29	0.996	128.259	0.7300	82.375	891.28	0.102	82.454	891.24	0.235
i-Butylcyclopentane	892.11	0.980	126.243	0.7809	82.743	892.17	0.017	82.390	892.18	0.008
N ₂₁	892.96	0.980	126.243	0.7800	83.054	892.92	0.030	82.938	892.86	0.015
?	893.20				83.607	893.23	0.030	83.590	893.22	0.009
?	894.00				80.659	893.15	0.033	81.076	893.14	0.015
t-7-Methyloctene-3	895.10	0.980	126.241	0.6826						
N ₂₂	895.99	0.980	126.243	0.7800	84.408	896.11	0.057	84.519	896.14	0.029
?	896.76				84.742	896.88	0.014	84.855	896.93	0.007
N ₂₃ /c-nonene-2	897.24	0.980	126.243	0.7800	84.967	897.41	0.035	85.075	897.44	0.018
t-Nonene-3	897.94	0.980	126.241	0.6826						
?	898.44				84.255	898.49	0.034			
I10	898.70	0.996	128.259	0.7300	85.566	898.78	0.051	85.709	898.90	0.147
?	899.19				87.478	898.94	0.046			
n-Nonane	900.20	0.996	128.259	0.7176	86.082	900.00	0.214	86.186	900.01	0.086
1,1-Methylethylcyclohexane	901.39	0.980	126.243	0.8062	86.378	901.62	0.035	86.476	901.59	0.016
3,7-Dimethyloctene-1	903.40	0.980	140.270	0.7013						
?	904.38				86.929	904.59	0.023	87.029	904.57	0.012
N ₂₅	905.50	0.980	126.243	0.7900	87.138	905.71	0.026	87.239	905.70	0.010
t-2,2,5,5-Tetramethylhexene-3	906.68	0.980	140.270	0.7013	87.352	906.86	0.012	85.841	906.79	0.006
i-Propylbenzene	912.28	0.933	120.195	0.8618	88.419	912.54	0.112	88.510	912.51	0.082
N ₂₆	913.43	0.980	126.243	0.7900						
N ₂₇	914.45	0.980	126.243	0.7900	88.839	914.76	0.045	88.957	914.88	0.029
c-Nonene-3	915.00	0.980	126.240	0.6826	88.198		0.004			
I11	916.40	0.994	142.286	0.7300				89.216	916.24	0.010
i-Propylcyclohexane	917.51	0.980	126.243	0.8022	89.365	917.50	0.020	88.849	917.53	0.009
?	918.60				90.523	918.03	0.006			
I12	921.30	0.994	142.286	0.7300	90.138	921.53	0.042	90.227	921.50	0.082
2,2-Dimethyloctane	922.59	0.994	142.286	0.7245	89.707		0.018	90.591	923.41	0.025
2,4-Dimethyloctane	924.39	0.994	142.286	0.7264	90.743	924.65	0.061	90.829	924.62	0.026
N ₂₈	926.32	0.980	126.243	0.7900	89.078	926.37	0.005	90.227		0.003
N ₂₉	927.99	0.980	126.243	0.7900	91.453	928.31	0.008	88.997	928.03	0.005
2,6-Dimethyloctane	930.83	0.994	142.286	0.7276	91.999	931.09	0.038	92.075	931.02	0.018
2,5-Dimethyloctane	932.66	0.994	142.286	0.7302	92.361	932.90	0.065	92.438	932.86	0.034
?	934.00							92.042		0.002
?	934.50				92.053		0.004			
n-Butylcyclopentane	936.13	0.980	126.243	0.7846	93.070	936.48	0.022			
I13	937.41	0.994	142.286	0.7300	93.309	937.65	0.033	93.261	937.01	0.045
?	937.60							92.567		0.002
N ₃₀	938.04	0.980	140.270	0.8000	92.877	938.25	0.025	93.516	938.28	0.011
I14	940.39	0.994	142.286	0.7300	93.378	940.53	0.009	91.869	940.47	0.005
?	941.00				93.027		0.005			
3,3-Dimethyloctane	942.30	0.994	142.286	0.7390	94.291	942.54	0.064	94.358	942.50	0.028
N ₃₁	943.42	0.980	140.270	0.8000	92.409	943.48	0.006	92.497	943.66	0.004
?	944.55				98.078	945.10	0.018	97.967	945.45	0.006
?	944.95				92.576	944.76	0.019	92.740	944.90	0.009
n-Propylbenzene	946.33	0.933	120.195	0.8620	95.116	946.61	0.627	95.182	946.55	0.401
?	947.54				93.127	947.53	0.020	93.521	947.53	0.008
3,6-Dimethyloctane	948.31	0.994	142.286	0.7363	95.496	948.44	0.023	95.564	948.39	0.011
3-Methyl-5-ethylheptane	949.41	0.994	142.286	0.7264	95.692	949.41	0.022	95.536	949.35	0.010
?	950.00				95.233		0.002	95.262		0.002

TABLE A1.1 *Continued*

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
N ₃₂	951.22	0.980	140.270	0.8000	96.296	951.32	0.018	95.351		0.002
?	951.50				95.460		0.004	95.442		0.002
?	953.00				95.589		0.003	95.580		0.002
1,3-Methylethylbenzene	954.42	0.933	120.195	0.8645	96.789	954.72	2.027	96.840	954.63	1.276
1,4-Methylethylbenzene	956.22	0.933	120.195	0.8612	97.157	956.49	0.878	97.212	956.42	0.571
?	957.40				94.626	957.43	0.011	96.594		0.005
N ₃₃	958.16	0.980	140.270	0.800						
?	958.90				95.762	958.89	0.008	96.900		0.003
?	960.80				96.911		0.006			
1,3,5-Trimethylbenzene	961.92	0.933	120.195	0.8652	98.348	962.16	0.996	98.400	962.09	0.638
2,3-Dimethyloctane	961.99	0.994	142.286	0.7379						
I15	963.67	0.994	142.286	0.7400	98.713	963.86	0.043	98.765	963.81	0.019
N34	964.76	0.980	140.270	0.8000						
?	965.40				98.438		0.003			
I16	966.53	0.994	142.286	0.7400	97.299	966.78	0.011	97.660	966.93	0.007
?	967.10				103.953	967.90	0.018	98.644		0.004
5-Methylnonane	967.89	0.994	142.286	0.7326	99.600	968.02	0.051	99.652	967.98	0.023
I17	969.41	0.994	142.286	0.7400	99.944	969.61	0.116	100.383	969.77	0.216
1,2-Methylethylbenzene	970.33	0.933	120.195	0.8807	100.146	970.57	0.605	100.194	970.52	0.447
2-Methylnonane	971.77	0.994	142.286	0.7264	100.431	971.87	0.115	100.479	971.83	0.048
?	973.00				100.210	973.37	0.017	99.457	973.29	0.011
3-Ethyl-octane	974.47	0.994	142.286	0.7399	101.009	974.55	0.022	100.826	974.53	0.010
?	975.05							100.504		0.003
N35	975.89	0.980	140.270	0.8000	101.669	976.12	0.018	99.640	976.08	0.006
3-Methylnonane	977.26	0.994	142.286	0.7334	101.629	977.38	0.119	101.675	977.35	0.046
?	978.30				101.271	978.34	0.011	100.504	978.16	0.007
N36	979.33	0.980	140.270	0.8000	100.066	979.21	0.008	99.919	979.30	0.005
3-Ethyl-2-methylheptene-2	979.35	0.980	140.270	0.7013						
I18	980.12	0.994	142.286	0.7400	102.306	980.46	0.029	102.362	980.49	0.065
I19	981.56	0.994	142.286	0.7400	101.282	981.67	0.007	99.704	981.50	0.007
1,2,4-Trimethylbenzene	983.40	0.933	120.195	0.8758	103.003	983.63	2.813	103.032	983.55	1.829
t-butylbenzene ^A	983.42	0.933	120.200	0.8665						
I20	985.82	0.994	142.286	0.7400	103.376	985.29	0.014	102.881	985.32	0.011
i-Butylcyclohexane	986.27	0.980	140.270	0.7960	103.606	986.32	0.023	103.402	986.29	0.010
I21	987.40	0.994	142.286	0.7400	103.819	987.26	0.044	103.845	987.19	0.025
?	987.60							104.866	987.79	0.026
I22	988.00	0.994	142.286	0.7400	105.334	988.43	0.018	102.938	987.84	0.019
?	988.60				102.239	988.63	0.009	102.157	988.55	0.005
I23	989.12	0.994	142.286	0.7400	103.648	989.04	0.011	103.593		0.002
N ₃₇	990.53	0.980	140.270	0.8000	104.581	990.68	0.015	104.365	990.61	0.006
?	991.24				104.174	991.37	0.010	103.370	991.29	0.005
Decene-1	992.81	0.990	140.270	0.7408	103.762	992.78	0.009	103.952	992.90	0.004
1t-Methyl-2-n-propylcyclohexane	993.55	0.980	140.270	0.8000						
2,3-Dimethyloctene-2	993.56	0.990	140.270	0.7400						
I24	993.70	0.994	142.286	0.7400	105.255	993.65	0.053	105.375	993.99	0.041
?	994.20							107.662	994.18	0.029
i-Butylbenzene	995.95	0.938	134.222	0.8532	105.781	995.97	0.063	105.813	995.95	0.046
I25	996.84	0.994	142.286	0.7400	105.356	996.81	0.021	105.398	996.72	0.024
Sec-Butylbenzene	997.79	0.938	134.222	0.8620	106.237	997.97	0.054	106.270	997.95	0.040
?	998.70				105.784		0.005	105.732		0.002
?	999.30							105.862		0.004
n-Decane	1000.20	0.994	142.286	0.7300	106.708	999.99	0.080	106.737	999.97	0.038
I26	1001.71	0.993	156.313	0.7400	106.952	1001.70	0.011	106.990	1001.72	0.016
N ₃₈	1003.39	0.980	140.260	0.8000	107.189	1003.32	0.028	107.218	1003.28	0.017
1,2,3-Trimethylbenzene	1006.88	0.933	120.195	0.8944	107.705	1006.91	0.539	107.732	1006.87	0.323
1,3-Methyl-i-propylbenzene	1009.84	0.938	134.222	0.8610	108.117	1009.73	0.058	108.144	1009.70	0.048
N ₃₉	1011.33	0.980	154.290	0.8000						
1,4-Methyl-i-propylbenzene	1013.24	0.938	134.222	0.8573	108.602	1013.08	0.034	108.638	1013.12	0.025
I27	1014.33	0.993	156.313	0.7400	106.759	1014.42	0.005	105.253	1014.26	0.002
I28	1015.86	0.993	156.313	0.7400	112.126	1016.63	0.012	113.222	1016.31	0.010
I29	1017.87	0.993	156.313	0.7400	107.874	1017.86	0.012	106.471	1017.20	0.009
2-3-Dihydroindene	1019.44	0.918	118.179	0.9640	109.504	1019.21	0.341	109.532	1019.21	0.163
?	1022.40				109.027		0.006	109.056		0.002
Sec-butylcyclohexane	1023.07	0.980	140.270	0.8140	107.271	1022.40	0.008	108.837	1022.14	0.007
I30	1024.82	0.993	156.313	0.7400	110.213	1023.97	0.028	110.225	1023.86	0.013
?	1026.50				108.990	1026.17	0.022	107.795	1026.59	0.011
1,2-Methyl-i-propylbenzene	1027.73	0.938	134.222	0.8766	110.677	1027.07	0.031	110.675	1026.86	0.017
?	1028.40				110.218		0.002			
3-Ethyl-nonane	1029.40	0.993	156.313	0.7440	110.240		0.003	110.276		0.002
?	1031.13				111.214	1030.68	0.011	111.238	1030.69	0.015
N ₄₀	1032.29	0.980	154.290	0.8000						
I31	1033.20	0.993	156.313	0.7400	111.495	1032.55	0.048	111.515	1032.54	0.020
I32	1036.92	0.993	154.290	0.8000	112.055	1036.34	0.021	111.809	1036.22	0.008

TABLE A1.1 *Continued*

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
?	1038.53				112.295	1037.88	0.012	112.295	1037.71	0.008
1,3-Diethylbenzene	1039.97	0.938	134.222	0.8639	112.538	1039.49	0.205	112.563	1039.49	0.112
1,3-Methyl-n-propylbenzene	1042.60	0.938	134.222	0.8609	112.944	1042.17	0.400	112.967	1042.18	0.232
I33	1044.35	0.993	156.313	0.7400	112.500	1044.04	0.020	112.163	1043.97	0.008
1,4-Diethylbenzene	1045.25	0.938	134.222	0.8620						
1,4-Methyl-n-propylbenzene	1046.40	0.938	134.222	0.8584	113.528	1046.00	0.237	113.550	1046.00	0.135
n-Butylbenzene	1047.48	0.938	134.222	0.8610	113.688	1047.04	0.121	113.710	1047.05	0.063
1,3-Dimethyl-5-ethylbenzene	1049.78	0.938	134.222	0.8800	114.044	1049.36	0.392	114.064	1049.35	0.218
1,2-Diethylbenzene	1051.72	0.938	134.222	0.8799	114.349	1051.35	0.043	114.372	1051.36	0.024
I34	1051.80	0.993	156.313	0.7400				113.736		0.003
t-Decahydronaphthalene	1053.12	0.980	154.290	0.8000	110.769	1052.71	0.004	110.626	1052.67	0.002
N41	1054.60	0.980	154.290	0.8000	114.769	1054.07	0.017	111.829	1054.53	0.008
?	1055.80							114.310		0.007
?	1056.50				115.938	1056.62	0.013	114.197	1056.55	0.007
1,2-Methyl-n-propylbenzene	1057.87	0.938	134.222	0.8736	115.304	1057.54	0.150	115.324	1057.56	0.078
I35	1058.87	0.993	156.313	0.7400	114.740		0.004	114.800		0.003
?	1059.00				114.838		0.002			
?	1059.50				114.918		0.003			
I36	1060.15	0.993	156.313	0.7400	115.058		0.008	115.622	1061.18	0.013
I37	1062.62	0.993	156.313	0.7400	116.030	1062.17	0.045	116.044	1062.16	0.020
?	1063.96				113.810	1063.86	0.006	115.398		0.005
I38	1065.53	0.993	156.313	0.7400	116.492	1065.12	0.038	116.508	1065.12	0.015
1,4-Dimethyl-2-ethylbenzene	1068.05	0.938	134.222	0.8772	116.905	1067.76	0.264	116.920	1067.77	0.140
A3	1068.90	0.938	134.222	0.8594						
1,3-Dimethyl-4-ethylbenzene	1069.53	0.938	134.222	0.8594	117.158	1069.36	0.307	117.173	1069.38	0.158
I39	1071.12	0.993	156.313	0.7400	114.335	1071.04	0.015	114.937	1071.08	0.005
?	1072.49				113.681	1072.50	0.003	116.744		0.006
?	1073.00				116.736		0.005			
I40	1074.39	0.993	156.313	0.7400	117.933	1074.24	0.178	118.598	1073.91	0.068
1,2-Dimethyl-4-ethylbenzene	1075.25	0.938	134.222	0.8745	118.068	1075.08	0.426	118.079	1075.09	0.250
?	1076.00				117.400		0.002			
?	1077.00							117.638		0.003
?	1078.00				115.592	1078.25	0.007	114.048	1078.31	0.005
I41	1079.65	0.993	156.313	0.7400	118.759	1079.41	0.012	117.600	1079.52	0.006
1,3-Dimethyl-2-ethylbenzene	1080.68	0.938	134.222	0.8904	118.945	1080.60	0.031	118.958	1080.62	0.017
I42	1081.60	0.993	156.313	0.7400	114.969	1081.44	0.003	114.988	1081.58	0.003
?	1083.35							118.280		0.002
?	1083.60							118.402		0.003
I43	1084.18	0.993	156.313	0.7400	119.495	1084.00	0.016	118.753	1084.00	0.008
?	1085.30				118.946	1085.34	0.009	116.463	1085.44	0.004
?	1086.54				119.122	1086.45	0.009	118.927		0.006
?	1087.50							117.379	1088.22	0.004
?	1088.80				120.278	1088.86	0.011	117.484	1088.96	0.004
Undecene-1	1090.45	0.980	154.300	0.7503	120.533	1090.42	0.022	120.544	1090.44	0.012
1,4-Methyl-t-butylbenzene	1092.00	0.942	148.240	0.8500	120.778	1091.92	0.038	120.788	1091.96	0.018
1,2-Dimethyl-3-ethylbenzene	1093.12	0.938	134.222	0.8921	120.985	1093.18	0.113	120.993	1093.22	0.058
?	1094.89				117.966	1094.87	0.008	117.921	1094.88	0.004
?	1095.78				118.105	1095.78	0.013	118.273	1095.78	0.007
1,2-Ethyl-i-propylbenzene	1097.22	0.942	148.240	0.8900	121.661	1097.33	0.011	119.139	1097.36	0.005
?	1098.54				118.119	1098.85	0.009	119.095	1098.88	0.003
?	1099.00				120.899		0.005	125.366	1099.10	0.004
n-Undecane	1100.00	0.993	156.313	0.7440	122.105	1100.03	0.053	122.106	1100.03	0.020
1,4-Ethyl-i-propylbenzene	1102.50	0.942	148.240	0.8900	122.417	1102.56	0.012	121.163	1102.56	0.006
1,2,4,5-Tetramethylbenzene	1104.83	0.938	134.222	0.8875	122.718	1104.99	0.234	122.720	1105.00	0.116
1,2-Methyl-n-butylbenzene	1107.30	0.942	148.240	0.8900						
1,2,3,5-Tetramethylbenzene	1108.79	0.938	134.222	0.8903	123.207	1108.93	0.319	123.208	1108.94	0.158
?	1110.82				119.539	1110.82	0.006	122.376		0.005
?	1112.39				122.367		0.002			
?	1113.53				123.033	1112.50	0.047	123.661	1112.56	0.007
?	1115.92				126.210	1113.03	0.045	122.689		0.003
?	1117.49				124.100	1116.08	0.023	123.509	1115.99	0.008
?	1119.00				120.791	1117.53	0.007	124.824	1116.58	0.011
?	1119.50							123.251		0.001
?	1120.13				124.606	1120.12	0.017	124.435	1120.07	0.006
?	1121.30				124.756	1121.30	0.015	123.419	1121.29	0.005
1,2-Methyl-t-butylbenzene	1122.80	0.942	148.240	0.8900						
?	1124.62				123.227	1124.59	0.006	124.027		0.006
?	1126.18				127.869	1126.31	0.007			
5-Methylindan	1127.35	0.938	132.200	0.8900	125.539	1127.50	0.300	125.538	1127.50	0.122
?	1127.30				124.651		0.002			
I43	1131.42	0.992	170.340	0.7530	126.032	1131.37	0.020	125.747	1131.30	0.008
4-Methylindan	1133.70	0.938	132.200	0.8900	126.337	1133.74	0.072	126.333	1133.72	0.031
?	1134.90				134.919	1134.95	0.032	135.048	1134.95	0.013

TABLE A1.1 *Continued*

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
1,2-Ethyl-n-propylbenzene	1136.52	0.942	148.240	0.8900	126.698	1136.56	0.105	126.692	1136.53	0.043
2-Methylindan	1138.11	0.938	132.200	0.9034	126.908	1138.22	0.289	126.903	1138.20	0.121
1,3-Methyl-n-butylbenzene	1140.67	0.942	148.240	0.8900	127.238	1140.79	0.029	127.233	1140.74	0.012
1,3-Di-i-propylbenzene	1142.70	0.945	162.272	0.8900	127.496	1142.79	0.103	127.490	1142.77	0.050
s-Pentylbenzene	1144.27	0.942	148.240	0.8900	127.697	1144.35	0.084	127.692	1144.33	0.033
?	1146.90				129.097	1146.97	0.008	126.758		0.004
n-Pentylbenzene	1149.04	0.942	148.240	0.8900	128.303	1149.00	0.123	128.297	1148.99	0.053
?	1149.83				127.775	1149.87	0.039	129.737	1149.75	0.015
1t-M-2-(4-MP)cyclopentane	1151.80	0.980	168.320	0.8000						
1,2-Di-propylbenzene	1153.16	0.945	162.272	0.8900	128.847	1153.19	0.045	128.839	1153.17	0.019
?	1154.09				128.970	1154.13	0.046	128.961	1154.11	0.019
?	1155.98				127.861		0.005	127.862		0.004
?	1157.64				129.427	1157.63	0.051	128.886	1157.45	0.021
?	1158.00							130.655	1158.06	0.023
1,4-Di-i-propylbenzene	1159.52	0.945	162.272	0.8900	129.675	1159.53	0.079	129.666	1159.51	0.033
Tetrahydronaphthalene	1163.30	0.924	132.206	0.9695	130.166	1163.27	0.034	130.156	1163.24	0.013
?	1165.13				128.975		0.004	128.924		0.002
?	1166.34				129.516	1166.43	0.076	130.568	1166.34	0.030
Naphthalene	1168.01	0.896	128.174	1.0253	130.803	1168.08	0.438	130.792	1168.05	0.190
?	1168.50				129.780		0.004	129.802		0.006
1-t-Butyl-3,5-dimethylbenzene	1169.25	0.945	162.272	0.8900	140.612	1169.00	0.013			
1,4-Ethyl-t-butylbenzene	1173.72	0.945	162.272	0.8900	131.554	1173.75	0.083	131.544	1173.73	0.030
I45	1177.88	0.942	170.300	0.7530	132.112	1177.91	0.124	132.101	1177.89	0.047
I46	1179.46	0.942	170.300	0.7530	132.325	1179.51	0.062	132.313	1179.48	0.022
?	1181.20				131.043		0.003	131.107		0.003
I47	1183.44	0.942	170.300	0.7530	132.864	1183.52	0.083	132.850	1183.47	0.032
I48	1187.14	0.942	170.300	0.7530	133.358	1187.20	0.071	133.347	1187.17	0.027
1,3-Di-n-propylbenzene	1188.64	0.945	162.272	0.8900	133.560	1188.67	0.077	133.547	1188.66	0.032
A5	1190.24	0.945	162.272	0.8900	133.778	1190.29	0.052	133.765	1190.26	0.023
?	1191.00				132.431		0.009	132.415		0.005
Dodecene-1	1192.19	0.980	168.330	0.7584	129.949	1192.19	0.012	132.673		0.005
?	1193.83				130.211	1193.93	0.012	130.350	1193.95	0.004
?	1194.60				132.870	1194.59	0.011			
?	1196.00				129.167	1196.71	0.006	133.044		0.003
A6	1198.52	0.945	162.272	0.8900	134.904	1198.56	0.040	134.888	1198.53	0.016
n-Dodecane	1200.00	0.992	170.340	0.7530	135.106	1200.07	0.046	135.089	1200.03	0.016
?	1202.51				135.377	1202.51	0.013	135.454	1202.41	0.007
?	1204.12				132.955	1204.05	0.025	132.926	1204.09	0.009
?	1205.70				132.109	1205.60	0.002	134.038		0.004
?	1208.41				136.041	1208.47	0.029	136.025	1208.45	0.010
1,3,5-Triethylbenzene	1211.79	0.945	162.272	0.8897	136.427	1211.94	0.015	135.785	1211.76	0.005
?	1212.90				132.913	1213.33	0.015	1213.196	1213.37	0.007
?	1213.71				135.006	1213.74	0.017	139.107	1214.01	0.006
?	1215.50				147.556	1215.90	0.028	141.394	1215.65	0.014
?	1216.27				137.053	1217.68	0.059	137.053	1217.63	0.020
?	1217.50				135.296		0.036			
?	1220.12				133.673	1220.92	0.007	135.640		0.005
?	1220.90				135.780	1220.94	0.009	131.628	1220.92	0.002
?	1222.36				137.009	1222.52	0.057	136.986	1222.46	0.021
?	1223.70				144.395	1223.59	0.051	144.409	1223.50	0.020
?	1225.08				137.897	1225.05	0.061	137.878	1224.99	0.023
?	1228.60				138.310	1228.69	0.031	138.291	1228.66	0.011
?	1230.00				136.676		0.020	136.675		0.008
1,2,4-Triethylbenzene	1230.83	0.945	162.272	0.8897	138.551	1230.60	0.019	137.335	1230.73	0.008
?	1232.23				139.172	1232.17	0.027	136.248	1232.12	0.009
?	1235.00				149.881	1235.00	0.008	137.282		0.013
?	1236.42				140.655	1236.40	0.030	141.732	1236.33	0.013
?	1237.42				136.485	1237.39	0.038	136.151	1237.46	0.014
?	1238.00				150.631	1238.00	0.009	137.580		0.002
1,4-Methyl-n-pentylbenzene	1241.71	0.945	162.272	0.8897	139.782	1241.65	0.111	139.761	1241.62	0.041
?	1242.50				151.463	1243.85	0.047			
?	1244.15				140.120	1244.60	0.040	140.045	1244.10	0.018
?	1245.00				142.998	1246.66	0.013	136.979	1245.15	0.005
?	1246.48				140.387	1246.86	0.016	140.890	1246.61	0.006
?	1248.73				140.651	1249.22	0.015	139.999	1248.99	0.006
?	1251.16				140.918	1251.54	0.033	140.895	1251.51	0.012
n-Hexylbenzene	1252.85	0.945	162.272	0.8897	141.130	1253.37	0.048	141.107	1253.34	0.019
?	1254.25				137.178	1254.09	0.006	139.194		0.003
?	1255.61				138.479	1255.85	0.108	136.643	1255.74	0.042
?	1257.39				144.513	1257.88	0.040	146.044	1257.31	0.030
?	1259.54				138.912	1259.82	0.053	138.882	1259.85	0.020
?	1262.15				142.153	1262.20	0.020	142.129	1262.17	0.007
?	1262.55				139.337	1263.68	0.017	137.487	1263.60	0.006

TABLE A1.1 *Continued*

Component	Average RI	RRF	MW	Relative Density	RFA Gasoline			CCF Gasoline		
					Min.	Index	Mass %	Min.	Index	Mass %
?	1264.03				144.425	1265.33	0.014	149.847	1265.65	0.008
?	1265.92				142.689	1266.78	0.043	142.663	1266.76	0.016
?	1266.71				155.002	1268.55	0.025	154.983	1268.25	0.010
?	1269.02				139.934	1269.13	0.031	139.907	1269.18	0.011
I49	1270.79	0.991	184.370	0.7560	143.152	1270.72	0.016	138.284	1270.93	0.005
?	1271.58				140.243	1271.91	0.011	138.379	1271.81	0.004
?	1273.13				143.553	1273.26	0.020	143.558	1273.31	0.009
1,2,3,4,5-Pentamethylbenzene	1274.04	0.942	148.240	1.0000	143.534	1273.98	0.094	143.510	1273.98	0.035
?	1276.70							141.702		0.004
?	1277.23				143.318	1277.27	0.007			
?	1279.96				144.231	1279.89	0.023	144.205	1279.88	0.008
2-Methylnaphthalene	1282.57	0.903	143.170	1.0200	144.522	1282.34	0.431	144.494	1282.33	0.171
?	1286.59				144.932	1285.80	0.034	142.460	1285.86	0.014
?	1287.50				157.493	1285.55	0.036	152.473	1286.23	0.013
?	1288.77				145.273	1288.67	0.032	144.965	1288.52	0.013
Tridecene-1	1290.10	0.980	182.350	0.7658				143.309		0.002
?	1292.41				150.934	1292.15	0.015	143.429		0.002
?	1293.81				145.906	1293.96	0.020	144.668	1293.93	0.007
?	1295.08				142.847	1295.31	0.019	142.817	1295.36	0.008
1-Methylnaphthalene	1297.72	0.903	143.170	1.0200	146.330	1297.50	0.175	146.303	1297.49	0.074
n-Tridecane	1300.00	0.991	184.370	0.7564	146.635	1300.11	0.040	146.604	1300.09	0.013
C ₁₄ ⁺	1300.50						0.361			0.127

^A Components known to co-elute. Unknown components whose group type are known are named with a letter (that is, O for olefin) and a consecutive number. If consecutive numbers are missing, they have been identified by name.



TABLE A1.2 Analyses of Naphtha, Reformate, Alkylate, and Other Refinery Streams

NOTE 1—These data consist of analyses of various refinery samples and the purpose for this table is to illustrate that the DHA test method is equally applicable to other refinery samples as well as to the analysis of spark-ignition engine gasolines. This table presents the retention times and retention indices of the components covering a range of different concentrations as well as the actual analyses.

Component	Cooperative Study Average RI	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D					
		Sample: PONA-Va *NJ* Reference Mixture	Index	Mass %	Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate	Min.	Index	Mass %	Sample: Platformate Auto/Oil C-Matrix Reformate	Min.	Index	Mass %	Sample: No. 1 Ref Naphtha ASTM No. 1 Reference Sample	Min.	Index	Mass %
Methane	100.00															
Ethylene	178.10															
Ethane	200.00	6.70	201.5	0.033												
Propylene	284.00															
Propane	300.00	7.20	300.0	0.166												
i-Butane	366.15	8.13	366.0	0.099										8.15	366.0	0.350
Methanol	378.82	8.42	378.6	0.362												
Butene-1	390.72	8.74	390.6	0.087												
Isobutylene	391.51	8.76	391.5	0.090												
1,3-Butadiene	394.93	8.87	395.1	0.005												
n-Butane	400.00	9.02	400.0	0.315	9.02	400.0	0.155	9.04	400.0	0.596	9.04	400.0	1.844			
Vinyl acetylene	409.00															
t-Butene-2	412.09	9.38	412.0	0.353												
2,2-Dimethylpropane	415.10	9.46	414.6	0.005												
c-Butene-2	427.74	9.92	427.5	0.438												
1,2-Butadiene	450.00															
Ethanol	455.33	11.13	454.8	6.914												
3-Methylbutene-1	460.84	11.43	460.5	0.418												
O ₁	469.00															
O ₂	474.00															
i-Pentane	477.45	12.43	477.2	9.757	12.43	477.2	11.716	12.45	477.1	3.248	12.45	477.1	0.158	12.46	477.1	5.163
*Acetone	477.55															
1,4-Pentadiene	481.18	12.70	481.1	0.004												
Butyne-2	488.00															
Pentene-1	490.83	13.40	490.8	0.951												
i-Propanol	493.38	13.58	493.1	0.310												
2-Methylbutene-1	496.66	13.87	496.7	1.811												
n-Pentane	500.00	14.14	500.0	1.850	14.14	500.0	0.599	14.17	500.0	2.693	14.17	500.0	0.428	14.18	500.0	2.875
Isoprene	506.02	14.47	506.0	0.072												
t-Pentene-2	510.56	14.74	510.7	1.641												
3,3-Dimethylbutene-1	516.79	15.09	516.8	0.016												
c-Pentene-2	519.53	15.27	519.7	0.888												
t-Butanol	521.64	15.37	521.4	0.250												
2-Methylbutene-2	524.92	15.61	525.2	1.980	15.61	525.3	0.005	15.64	525.3	0.060	15.64	525.3	0.060	15.65	525.2	0.165
1t,3-Pentadiene	527.97	15.81	528.3	0.042												
3-Methylbutadiene-1,2	535.00															
Cyclopentadiene	538.05	16.48	538.3	0.015												
2,2-Dimethylbutane	540.54	16.64	540.7	0.087	16.64	540.6	0.005	16.67	540.6	0.185	16.67	540.6	0.067	16.68	540.6	0.330
1c,3-Pentadiene	541.90															
O ₅	547.70															
O ₆	549.70															
Cyclopentene	557.21	17.91	557.5	0.157												
n-Propanol	560.00															
4-Methylpentene-1	562.02	18.31	562.5	0.062												
3-Methylpentene-1	562.81	18.37	563.2	0.097												
Cyclopentane	566.84	18.70	567.1	1.140	18.70	567.1	0.007	18.72	567.1	0.094	18.72	567.0	0.190	18.73	567.1	0.232
2,3-Dimethylbutane	569.24	18.91	569.6	0.611	18.90	569.5	2.601	18.93	569.6	0.346	18.94	569.6	0.228	18.95	569.6	1.522
Methyl-t-butylether	570.65	19.01	570.8	0.731												
4-Methyl-c-pentene-2	571.00															
2,3-Dimethylbutene-1	572.67	19.21	573.1	0.061												
2-Methylpentane	573.70	19.30	574.1	2.213	19.29	573.9	0.754	19.32	574.1	1.878	19.33	574.0	1.857	19.34	574.0	2.536



TABLE A1.2 Continued

Component	Cooperative Study Average RI	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D		
		Sample: PONA-Va *NJ* Reference Mixture	Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate	Sample: Platformate Auto/Oil C-Matrix Reformate	Sample: No. 1 Ref Naphtha ASTM No. 1 Reference	Sample: ASTM D02 Reference	Sample: ASTM Indolene Standard	Min.	Index	Mass %	Min.	Index	Mass %
4-Methyl-t-pentene-2	575.47	19.46	575.9	0.175	19.49	575.9	0.012	19.50	575.9	0.027			
O ₈	578.00												
2-Methyl-1,4-pentadiene	579.00												
1,5-Hexadiene	581.90												
3-Methylpentane	585.52	20.38	585.7	1.450	20.40	585.7	1.441	20.42	585.7	1.831			
2-Methylpentene-1	590.19	29.83	590.4	0.288	20.86	590.4	0.015	20.88	590.4	0.041			
Hexene-1	591.06	20.92	591.2	0.157	20.95	591.3	0.005	20.97	591.2	0.020			
O ₁₁	592.00												
i-Butanol	593.50												
1c/t,4-Hexadiene	597.14												
2-Ethylbutene-1	598.95												
n-Hexane	600.00	21.83	600.0	1.754	21.85	600.0	1.735	21.87	600.0	2.573			
Diisopropylether	601.90												
t-Hexene-3	602.83	22.06	602.9	0.323	22.09	603.0	0.010	22.11	603.0	0.035			
c-Hexene-3	603.56	22.11	603.6	0.076	22.15	603.8	0.002	22.32	605.5	0.049			
r-Hexene-2	605.44	22.27	605.5	0.436	22.30	605.6	0.014	22.52	608.0	0.073			
2-Methylpentene-2	607.86	22.47	608.0	0.547	22.50	608.0	0.028						
4-Methylcyclopentene													
3-Methyl-c-pentene-2	610.54	22.68	610.6	0.308	22.71	610.7	0.018	22.73	610.6	0.041			
3-Methylcyclopentene	611.61	22.76	611.5	0.040	23.07	614.8	0.008	22.81	611.5	0.007			
O ₁₃	613.08	22.90	613.2	0.003									
c-Hexene-2	614.67	23.04	614.8	0.250									
O ₁₄	617.06	23.23	617.1	0.005									
Ethyl-t-butylether	619.00												
3,3-Dimethylpentene-1	620.91	23.58	621.1	0.602	23.60	621.1	0.026	23.63	621.1	0.053			
3-Methyl-t-pentene-2	622.11	23.70	622.4	0.009									
2-Butanol	622.40												
4,4-Dimethyl-pentene-2	623.10												
2,2-Dimethylpentane	624.17	23.86	624.3	0.053	23.88	624.2	0.118	23.91	624.2	0.174			
Methylcyclopentane	625.86	23.99	625.7	1.057	24.02	625.7	0.310	24.04	625.7	0.982			
Cyclic Diolefin or triolefin	627.00												
2,4-Dimethylpentane	630.60	24.45	630.8	0.768	24.47	630.7	0.243	24.50	630.7	1.039			
2,3,3-Trimethylbutene-1	631.00												
Cyclic Diolefin or triolefin	632.90	24.67	633.1	0.013									
2,2,3-Trimethylbutane	634.86	24.83	634.7	0.046	24.86	634.9	0.010	24.88	634.8	0.091			
Cyclic diolefin or triolefin	636.38	25.00	636.6	0.010									
4,4-Dimethyl-c-pentene-2	638.30	25.16	638.4	0.011	26.00	646.6	0.005	26.04	646.8	0.006			
O ₁₇	641.97	25.52	642.0	0.006									
3,4-Dimethylpentene-1	642.87	25.60	642.9	0.017									
2,4-Dimethylpentene-1	646.65	25.99	646.8	0.029									
Diolefin	647.67	26.06	647.6	0.021									
1-Methylcyclopentene	648.71	26.18	648.7	0.328	26.32	649.8	1.786	26.33	649.7	0.445	26.24	648.7	
Benzene	649.92	26.29	649.8	0.541							26.35	649.8	
3-Ethylpentene-1	650.00												
*n-Butanol	650.02												
3-Methylhexene-1	650.95	26.42	651.0	0.038									
2-Methyl-c-hexene-3	652.60	26.60	652.8	0.020	26.61	652.7	0.003						



TABLE A1.2 Continued

Component	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D								
	Cooperative Study Average RI	Sample: PONA-Va *NJ* Reference Mixture	Index	Min.	Mass %	Index	Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate	Min.	Mass %	Index	Sample: No. 1 Ref Naphtha	Min.	Mass %	Index	Sample: ASTM Indolene Standard	Min.	Mass %	Index
1,1-Dimethylcyclohexane	780.48	46.55	780.2	0.099				46.62	0.291	780.0	46.67	780.3	0.047					
2,2,5-Trimethylhexane	782.93	47.16	782.9	0.274				47.22	0.010	783.0	47.25	783.0	1.175					
3-Ethylmethylcyclopentane	784.35	47.44	784.1	0.199				47.50	0.037	784.2	47.52	784.2	0.054					
2,6-Dimethylheptene-1	785.55	47.64	785.0	0.034				48.01	0.034	786.4	48.03	786.2	0.049					
3-Ethylmethylcyclopentane	786.55	47.95	786.4	0.154				48.32	0.037	787.8	48.34	787.5	0.094					
2-Ethylmethylcyclopentane	787.86	48.26	787.7	0.172				48.79	0.012	789.8	48.82	789.5	0.026					
*Octene-1	787.87																	
1,1-Methylethylcyclopentane	788.78	48.54	788.9	0.039				48.70	0.050	789.3	48.70	789.3	0.141					
	789.88	48.73	789.7	0.023														
2,2,4-Trimethylhexane	790.75	49.00	790.8	0.053														
1t,2-Dimethylcyclohexane	792.77	49.41	792.6	0.314				49.47	0.021	792.7	49.49	792.4	0.014					
t-Octene-4	794.21	49.83	794.3	0.075														
3,5,5-Trimethylhexene-1	795.00																	
t-Octene-3	796.00																	
1c,2c,3-Trimethylcyclopentane	797.25	50.57	797.3	0.200				50.61	0.018	797.3	50.68	797.3	0.014					
1t,3-Dimethylcyclohexane	798.80	50.91	798.7	0.051														
n-Octane	800.00	51.23	800.0	1.978				51.27	0.837	800.0	51.36	800.0	1.159					
1c,4-Dimethylcyclohexane	801.05																	
3,3-Dimethylheptene-1	802.50																	
Octene-2	804.40	52.33	804.5	0.082														
O ₅₀	805.50																	
I ₂	806.39	52.79	806.3	0.109				52.92	0.119	806.6			0.041					
i-Propylcyclopentane	808.06	53.21	808.0	0.123									0.028					
2,4,4-Trimethylhexane	808.50																	
O ₅₂	810.62	53.85	810.5	0.013														
O ₅₃	812.50																	
N ₁	813.47	54.60	813.4	0.044														
2,2,3,4-Tetramethylpentane	815.02	54.93	814.6	0.031														
	816.45	55.35	816.2	0.014														
2,3,4-Trimethylhexane	818.10	55.77	817.8	0.057				55.75	0.585	817.5			0.013					
N ₂	819.93	56.21	819.4	0.063														
	820.85	56.45	820.3	0.016														
N ₃	822.29	56.89	821.9	0.069														
2,3,3-Trimethylhexene-1	824.74	57.55	824.3	0.024														
1c,2-Dimethylcyclohexane	826.48	57.95	825.7	0.084				58.04	0.174	825.6	58.04	825.9	0.030					
2,3,5-Trimethylhexane	827.51	58.23	826.8	0.172														
2,2-Dimethylheptane	829.76	59.66	831.8	0.530				59.71	0.030	831.8	59.79	831.7	0.264					
1,1,4-Trimethylcyclohexane	834.07	60.06	833.1	0.031														
N ₄	834.96	60.42	834.4	0.263														
2,2,3-Trimethylhexane	836.47	60.80	835.6	0.030				60.12	0.022	833.2	60.17	833.1	0.020					
								60.49	0.059	834.5	60.55	834.3	0.183					
2,4-Dimethylheptane								60.91	0.061	835.6	60.91	835.6						



TABLE A1.2 Continued

Component	Cooperative Study Average RI	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D		
		Sample: PONA-Va *NJ* Reference Mixture	Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate	Sample: Platformate Auto/Oil C-Matrix Reformate	Sample: No. 1 Ref Naphtha	Sample: ASTM D02 Reference Standard	Sample: ASTM Indolene Standard	Sample: No. 1 Ref Naphtha	Sample: ASTM D02 Reference Standard	Sample: ASTM Indolene Standard	Sample: No. 1 Ref Naphtha	Sample: ASTM D02 Reference Standard	Sample: ASTM Indolene Standard
		Min.	Index	Mass %	Min.	Index	Mass %	Min.	Index	Mass %	Min.	Index	Mass %
4,4-Dimethylheptane	838.68	61.43	837.8	0.323	61.41	837.5	0.020	61.55	837.7	1.205	61.58	837.9	0.175
Ethylcyclohexane	840.20												
n-Propylcyclopentane	841.38	62.20	840.4	0.076	62.20	840.2	0.011	62.33	840.3	0.156	62.35	840.5	0.027
*1c,3c,5-Trimethyl-cyclohexane	841.40												
2,5-Dimethylheptane	842.63	62.68	841.9	0.326	62.66	841.7	0.192	62.74	842.0	0.179	62.84	842.1	0.212
3,3-Dimethylheptane	843.96	63.09	843.3	0.086	63.04	842.9	0.020	63.16	843.4	0.062	63.27	843.5	0.069
3,5-Dimethylheptane	845.02	63.30	843.9	0.038									
2,6-Dimethylheptane	846.47	63.79	845.5	0.083									
1,1,3-Trimethyl-cyclohexane	848.43	64.40	847.5	0.070									
2,4-Dimethylheptene-1	849.43												
N ₇	850.89												
N ₆	852.36	65.77	851.8	0.046				65.98	852.0	0.103	66.03	852.2	0.020
N ₁₀	853.04												
Ethylbenzene	854.65	66.41	853.7	0.728	66.43	853.7	0.032	66.64	854.3	7.364	66.61	854.0	2.653
N ₁₁	854.70												
1c,2i,4i-Trimethyl-cyclohexane	856.34	66.91	855.3	0.222	66.90	855.1	0.049	67.04	855.3	0.580	67.08	855.5	0.106
I ₃	858.51	67.63	857.5	0.032									
2-Methyloctene-1	859.80	68.01	858.6	0.030									
I ₄	860.89	68.51	860.1	0.033									
2-Methyloctene-2	862.14	69.05	861.7	0.021									
N ₁₂	863.00	69.25	862.3	0.016									
N ₁₃	863.77												
1,3-Dimethylbenzene	864.22	69.69	836.6	1.828				69.98	864.3	10.485	69.94	863.9	5.660
1,4-Dimethylbenzene	865.20	70.06	864.6	0.693				70.31	865.2	4.479	70.28	864.9	2.475
2,3-Dimethylheptane	866.02	70.36	865.5	0.261	70.32	865.3	0.127						
3,4-Dimethylheptane	867.94	70.99	867.0	0.102	70.93	867.0	0.019	71.09	867.5	0.029	71.15	867.4	0.049
3,4-Dimethylheptane	868.78	71.30	868.2	0.042	71.23	867.9	0.021	71.38	868.3	0.029	71.48	868.4	0.032
N ₁₄	869.70	71.62	869.1	0.033									
I ₅	870.95	72.11	870.5	0.106				72.23	870.7	0.053	72.28	870.6	0.051
4-Ethylheptane	872.73	72.83	872.5	0.025									
4-Methyloctane	873.81	73.19	873.5	0.393				73.27	873.6	0.217	73.34	873.6	0.187
2-Methyloctane	874.76	73.55	874.4	0.560				73.63	874.6	0.215	73.69	874.5	0.222
N ₁₅	876.00	73.95	875.5	0.045	73.53	874.3	0.006	74.05	875.4	0.072	74.10	875.7	0.018
1c,2i,3-Trimethyl-cyclohexane	877.98	74.66	877.5	0.073				74.76	877.4	0.208	74.86	877.7	0.033
3-Ethylheptane	879.11	75.19	878.9	0.135									
3-Methyloctane	880.24	75.62	880.0	0.577				75.24	878.9	0.063	75.32	878.9	0.066
1c,2i,4c-Trimethyl-cyclohexane	881.67	76.05	881.2	0.061	76.23	881.6	0.042	75.68	880.1	0.288	75.75	880.1	0.280
1,1,2-Trimethyl-cyclohexane	882.78	76.49	882.3	0.065				76.17	881.1	0.090	76.40	881.8	0.124
1,2-Dimethylbenzene	883.47	76.83	883.2	1.011	76.49	882.3	0.007	76.60	882.3	0.158	76.64	882.5	0.014
I ₆	884.87	77.43	884.8	0.022	76.86	883.2	0.062	76.98	883.5	5.364	76.99	883.4	3.212
I ₇	886.38	77.99	886.2	0.054	77.60	885.2	0.044	77.56	884.8	0.021	77.48	884.6	0.003
N ₁₈	887.87				77.97	886.1	0.296	77.91	885.7	0.020	77.76	885.4	0.082
N ₁₉	888.36	78.70	888.1	0.323				78.08	886.1	0.022	78.13	886.3	0.141
Nonene-1	889.00	78.91	888.6	0.036				78.62	887.8	0.011	78.83	888.1	0.172
I ₈	889.40	79.08	889.0	0.051				79.02	888.5	0.059	79.09	888.9	0.109
N ₂₀	889.51	79.32	889.6	0.112				79.15	888.9	0.109	79.45	889.7	0.047
I ₉	891.29	80.20	891.8	0.033	79.80	890.8	0.442	79.90	890.5	0.030	79.95	890.9	0.141
								80.26	891.6	0.057	80.30	891.8	0.011



TABLE A1.2 Continued

Component	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D			
	Coopera- tive Study Average RI	Sample: PONA-Va *NJ* Reference Mixture	Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate	Sample: Platformate Auto/Oil C-Matrix Reformate	Sample: No. 1 Ref Naphtha ASTM No. 1 Reference	Sample: ASTM Indolene Standard ASTM D02 Reference Standard	Index	Mass %	Min.	Index	Mass %	Min.	Index
i-Butylcyclopentane	892.11	80.54	892.7	0.091	80.70	892.7	0.116	80.69	892.8	0.024	80.69	892.8	0.024
N ₂₁	892.96												
	894.00												
t-7-Methyloctene-3	895.10												
N ₂₂	895.99	81.89	896.0	0.113	81.81	895.5	0.068	81.25	894.2	0.003	81.25	894.2	0.003
	896.76	82.22	896.8	0.019									
N ₂₃ /c-nonene-2	897.24	82.46	897.4	0.070									
t-Nonene-3	897.94												
	898.44												
I10	898.70	82.93	898.5	0.081	83.01	898.7	0.242	83.04	898.5	0.029	83.15	898.8	0.110
	899.19	83.17	899.1	0.026	83.33	899.4	0.007	83.26	899.0	0.140	83.15	898.8	0.110
n-Nonane	900.20	83.55	900.0	1.495									
1,1-Methylethylcyclo- hexane	901.39				83.57	900.0	0.224	83.68	900.0	4.771	83.64	900.0	0.691
3,7-Dimethyloctene-1	903.40												
N ₂₄	904.38	84.17	903.4	0.033	84.27	903.2	0.078	84.58	905.1	0.013	84.58	905.1	0.013
N ₂₅	905.50	84.43	904.9	0.108	85.54	904.7	0.056						
t-2,2,5,5- Tetramethylhexene-3	906.68												
	911.02	85.55	910.9	0.008	85.60	910.5	0.019	85.90	912.3	0.295	85.90	912.3	0.207
i-Propylbenzene	912.28	85.80	912.3	0.061	85.83	912.3	0.295	85.87	911.9	0.132	85.90	912.3	0.207
N ₂₆	913.43												
N ₂₇	914.45	86.15	914.1	0.145	86.27	914.7	0.020	86.22	913.8	0.493	86.25	914.1	0.068
c-Nonene-3	915.00												
i-Propylcyclohexane	917.51	86.75	917.3	0.086	86.50	915.9	0.015	86.82	917.0	0.234	86.85	917.4	0.036
	918.60	87.06	919.0	0.011	87.79	917.4	0.009	87.07	918.3	0.025	87.07	918.3	0.025
I12	921.30	87.53	921.4	0.098	87.57	921.4	0.013	87.60	921.1	0.228	87.64	921.5	0.093
2,2-Dimethyloctane	922.59												
2,4-Dimethyloctane	924.39	88.13	924.6	0.135	88.16	924.5	0.029	88.20	924.2	0.274	88.21	924.5	0.042
N ₂₈	926.32	88.49	926.4	0.018	88.56	926.1	0.051	88.56	926.1	0.051	88.56	926.1	0.051
N ₂₉	927.89	88.75	927.8	0.037	88.82	927.5	0.101	88.82	927.5	0.101	88.85	927.8	0.014
2,6-Dimethyloctane	930.83	89.33	930.7	0.221	89.39	930.8	0.014	89.40	930.5	0.800	89.43	930.8	0.107
2,5-Dimethyloctane	932.66	89.75	932.9	0.144	89.72	932.7	0.021	89.81	932.6	0.245	89.84	932.9	0.050
	934.50				90.51	936.7	0.053						
n-Butylcyclopentane	936.13	90.38	936.1	0.078	90.44	935.8	0.212	90.48	935.8	0.212	90.48	936.2	0.028
I13	937.41	90.68	937.7	0.078	90.71	937.6	0.015	90.76	937.4	0.179	90.66	937.1	0.031
N ₃₀	938.04	90.83	938.4	0.059	90.84	938.2	0.016	90.93	938.3	0.125	90.78	937.7	0.028
I14	940.39	91.21	940.3	0.042	91.26	939.9	0.121	91.26	939.9	0.121	91.26	939.9	0.121
3,3-Dimethyloctane	942.30	91.66	942.5	0.295	91.68	942.4	0.039	91.73	942.3	1.121	91.75	942.6	0.158
N ₃₁	943.42				92.15	944.4	0.241	92.15	944.4	0.241	92.15	944.7	0.015
	944.55	92.07	944.6	0.077									
	944.95												
n-Propylbenzene	946.33	92.43	946.4	0.242	92.46	946.3	1.417	92.49	946.1	0.203	92.27	945.2	0.011
	947.54	92.70	947.7	0.056	92.73	947.6	0.009	92.76	947.4	0.190	92.80	947.8	0.024
3,6-Dimethyloctane	948.31	92.85	948.5	0.051	92.89	948.4	0.007	92.92	948.2	0.069	92.80	947.8	0.024
3-Methyl-5- ethylheptane	949.41	93.08	949.6	0.125	93.15	949.4	0.669	93.15	949.4	0.669	93.18	949.6	0.063
N ₃₂	951.22	93.39	951.1	0.058	93.48	951.0	0.158	93.48	951.0	0.158	93.48	951.0	0.158
1,3- Methylethylbenzene	954.42	94.07	954.4	0.925	94.16	954.6	4.421	94.13	954.1	0.725	94.19	954.6	2.428
1,4- Methylethylbenzene	956.22	94.44	956.2	0.359	94.51	956.2	1.972	94.50	955.9	0.300	94.55	956.3	1.063



TABLE A1.2 Continued

Component	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D			
	Coopera- tive Study Average RI	Sample: PONA-Va *NJ* Reference Mixture	Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate	Sample: Platformate Auto/Oil C-Matrix Reformate	Sample: No. 1 Ref Naphtha ASTM No. 1 Reference	Sample: ASTM Indolene Standard ASTM D02 Reference Standard	Index	Mass %	Min.	Index	Mass %	Index	Mass %
N ₃₃	958.16	94.79	957.9	0.066	95.69	961.9	2.101	94.86	957.7	0.383	94.89	957.9	0.037
1,3,5-Trimethylbenzene	961.92	95.63	961.9	0.544	96.08	963.7	0.025	95.70	961.7	0.985	95.73	962.0	1.370
2,3-Dimethyloctane	961.99												
I15	963.67	96.05	963.9	0.081	96.96	967.9	0.023	96.11	963.7	0.144	96.14	963.9	0.025
N ₃₄	964.76	96.20	964.6	0.024				96.27	964.4	0.158			
I16	966.53												
5-Methylonane	967.89	96.93	968.1	0.159	97.43	970.1	1.356	97.00	967.9	0.262	97.03	968.1	0.033
I17	969.41	97.25	969.6	0.219	97.78	971.7	0.051	97.82	971.7	0.700	97.84	971.9	0.082
1,2-Methyl- Methyl-ethylbenzene	970.33	97.39	970.2	0.306	98.35	974.4	0.013	98.07	972.9	0.100	98.43	974.7	0.027
2-Methylonane	971.77	97.75	971.9	0.294	98.96	977.2	0.060	98.65	975.6	0.120	99.03	977.4	0.077
3-Ethyl- Ethyl-ethylbenzene	973.13	98.04	973.3	0.049	99.92	981.5	0.010	99.99	981.8	0.049	99.67	980.4	0.111
I18	974.47	98.33	974.6	0.080	100.32	983.3	5.960	100.28	983.1	1.477	100.34	983.4	3.921
3-Ethyl- Ethyl-ethylbenzene	975.89	98.57	975.7	0.059				100.45	983.9	0.266			
N ₃₅	977.26	98.93	977.4	0.260				100.93	986.0	0.445	100.96	986.2	0.026
I19	978.30	99.14	978.3	0.030	101.12	987.8	0.029	101.18	987.2	0.113	101.18	987.2	0.050
3-Ethyl-2- methylheptene-2	979.33				101.19	988.5	0.038	101.54	988.8	0.181			
I20	981.56	99.51	980.0	0.048	101.80	990.4	0.035	101.86	990.2	0.091			
1,2,4-Trimethylbenzene	983.40	100.22	983.2	1.286	102.02	991.4	0.019	102.13	991.4	0.022			
*t-Butylbenzene	983.42				102.53	993.6	0.078	102.51	993.1	0.122			
I21	984.20	100.72	985.5	0.035	102.60	993.7	0.051	102.75	994.2	0.173	102.72	994.1	0.098
I22	985.82	100.87	986.2	0.081	103.17	996.2	0.023	103.05	995.5	0.065	103.07	995.6	0.049
i-Butylcyclohexane	986.27	101.12	987.3	0.082	103.31	996.9	0.017	103.46	997.3	0.129	103.30	996.6	0.030
I23	987.40	101.38	988.5	0.059	103.80	999.0	0.008	104.00	999.6	0.048	103.52	997.6	0.086
I24	988.60	101.80	990.4	0.035	104.18	1001.1	0.028	104.08	1000.0	3.484	104.06	1000.0	0.211
I25	989.12	102.02	991.4	0.019	104.90	1005.9	1.075	104.34	1001.9	0.051	104.29	1001.7	0.020
N ₃₇	990.53	102.53	993.6	0.078	105.16	1008.2	0.007	104.56	1003.5	0.056	104.96	1006.5	0.751
Decene-1	991.24				105.34	1009.1	0.132	104.94	1006.2	0.561	104.96	1006.5	0.751
11-Methyl-2-n- propylcyclohexane	992.81				105.83	1012.6	0.039	105.25	1008.4	0.087	105.40	1009.6	0.078
2,3-Dimethyloctene-2	993.55				105.94	1013.7	0.006	105.38	1009.4	0.170	105.89	1013.1	0.026
I26	993.56				106.33	1016.5	0.013	105.86	1012.8	0.137	105.89	1013.1	0.026
I27	994.53	102.70	994.3	0.052	106.02	1014.0	0.038	106.02	1014.0	0.038			
i-Butylbenzene	995.95	103.00	995.7	0.027	106.33	1016.5	0.013	106.43	1017.0	0.052			
I28	996.20												
I29	996.84	103.10	996.1	0.030									
Sec-butylbenzene	997.79	103.44	997.6	0.051									
n-Decane	998.70	103.99	1000.0	0.739									
I30	1000.20												
I31	1001.71												
N ₃₈	1003.39	104.46	1003.4	0.049									
1,2,3-Trimethylbenzene	1006.88	104.87	1006.4	0.311									
I32	1008.70	105.16	1008.5	0.018									
1,3-Methyl-i- propylbenzene	1009.84	105.33	1009.7	0.057									
N ₃₉	1011.33												
1,4-Methyl-i- propylbenzene	1013.24	105.81	1013.1	0.050									
I33	1014.33	105.97	1014.2	0.013									
I34	1015.86	106.19	1015.8	0.013									

TABLE A1.2 Continued

Component	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D					
	Cooperative Study Average RI	Sample: PONA-Va		Sample: HFLA C-Matrix		Sample: Platformate		Sample: No. 1 Ref Naphtha		Sample: ASTM Indolene Standard		Sample: ASTM D02 Reference Standard			
		NJ Reference Mixture Index	Min.	Mass %	Auto/Oil C-Matrix Alkylate Min.	Index	Mass %	Auto/Oil C-Matrix Reformate Min.	Index	Mass %	ASTM No. 1 Reference Mixture Min.	Index	Mass %	ASTM D02 Reference Standard Index	Mass %
I29	1017.87	106.45	0.028	106.73	1019.4	0.011	106.64	1018.4	0.463	106.52	1017.5	0.102	106.71	1018.9	0.208
2-3-Dihydroindene	1019.44	106.62	0.257	107.15	1022.3	0.008				106.68	1018.7	0.146			
	1022.40									106.96	1020.6	0.101			
Sec-butylcyclohexane	1023.07	107.23	0.056	107.35	1023.7	0.009				107.29	1023.0	0.230	107.28	1022.9	0.016
	1024.50	107.45	0.051							107.50	1024.4	0.095	107.48	1024.4	0.018
I30	1024.82	107.68	0.034				107.68	1025.7	0.029	107.63	1025.4	0.055			
	1025.70									107.77	1026.3	0.044			
1,2-Methyl-i-propylbenzene	1026.50														
3-Ethylnonane	1027.73	107.90	0.109				107.91	1027.4	0.013	107.96	1027.7	0.747	107.98	1027.8	0.035
N ₄₀	1029.40	108.48	0.063	108.38	1030.9	0.044				108.55	1031.7	0.376	108.49	1031.4	0.217
I31	1031.13														
	1032.29									108.71	1032.9	0.105			
	1032.60	108.67	0.089							108.85	1033.8	0.115	108.82	1033.7	0.012
	1035.50														
I32	1036.92	109.17	0.053							109.24	1036.5	0.242	109.27	1036.8	0.017
	1038.53	109.45	0.020	109.37	1037.7	0.014	109.64	1039.4	0.477	109.50	1038.3	0.060	109.48	1038.2	0.101
1,3-Diethylbenzene	1039.97	109.63	0.173							109.73	1039.9	0.215	109.70	1039.7	0.143
	1040.50			109.92	1041.5	0.030	110.04	1042.1	0.942						
1,3-Methyl-n-propylbenzene	1042.60	110.01	0.257							110.06	1042.2	0.260	110.06	1042.2	0.737
I33	1044.35	110.31	0.064							110.36	1044.2	0.329			
1,4-Diethylbenzene	1045.25														
1,4-Methyl-n-propylbenzene	1046.40	110.57	0.152	110.60	1046.1	0.008	110.59	1045.9	0.596	110.62	1046.0	0.110	110.65	1046.2	0.264
n-Butylbenzene	1047.48	110.72	0.097							110.77	1047.0	0.147	110.79	1047.2	0.080
1,3-Dimethyl-5-ethylbenzene	1049.78	111.07	0.249				110.74	1046.9	0.224	111.12	1049.4	0.184	111.15	1049.6	0.402
							111.10	1049.4	0.934						
1,2-Diethylbenzene	1051.72	111.32	0.038				111.36	1051.1	0.070	111.27	1050.4	0.042	111.42	1051.4	0.027
I34	1051.80	111.49	0.040							111.55	1052.3	0.246	111.68	1053.1	0.023
t-Decahydronaphthalene	1053.12														
N ₄₁	1054.60	111.83	0.032	112.83	1061.1	0.038	112.88	1061.1	0.038	111.88	1054.6	0.093	112.93	1061.5	0.400
	1055.80	111.95	0.013	113.10	1062.9	0.007	113.05	1062.5	0.009	112.05	1055.7	0.062	113.20	1063.3	0.059
1,2-Methyl-n-propylbenzene	1057.87	112.26	0.123	113.47	1065.4	0.006	113.49	1065.4	0.009	112.31	1057.4	0.176	113.57	1065.8	0.037
I35	1058.87	112.46	0.018				113.82	1067.6	0.502	112.50	1058.7	0.047	113.87	1067.8	0.229
1,4-Dimethyl-2-ethylbenzene	1060.15	112.59	0.014							112.66	1059.8	0.036	114.11	1069.3	0.230
A3	1062.62	113.03	0.091	113.10	1062.9	0.007	114.05	1069.2	0.530	113.08	1062.6	0.199	114.27	1070.4	0.202
1,3-Dimethyl-4-ethylbenzene	1063.96	113.47	0.033	114.17	1070.0	0.025	114.74	1073.7	0.050	113.52	1065.5	0.231	114.80	1073.8	0.022
I39	1065.53	113.80	0.185				114.92	1074.9	0.886	113.67	1066.5	0.052	114.98	1075.0	0.419
	1068.05	113.80	0.185							113.85	1067.7	0.108			
1,3-Dimethyl-4-ethylbenzene	1069.53	114.01	0.316												
I39	1071.12	114.32	0.033	114.17	1070.0	0.025	114.74	1073.7	0.050	114.05	1069.0	0.380	114.80	1073.8	0.022
	1072.49						114.92	1074.9	0.886	114.36	1071.0	0.061	114.98	1075.0	0.419
I40	1074.39	114.72	0.147												
1,2-Dimethyl-4-ethylbenzene	1075.25	114.89	0.360												



TABLE A1.2 Continued

Component	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D		
	Coopera- tive Study Average RI	Sample: PONA-Va *NJ* Reference Mixture Index	Mass %	Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate Min.	Sample: Platformate Auto/Oil C-Matrix Reformate Index	Mass %	Sample: No. 1 Ref Naphtha ASTM No. 1 Reference Min.	Sample: No. 1 Ref Naphtha ASTM No. 1 Reference Index	Mass %	Sample: ASTM Indolene Standard ASTM D02 Reference Standard Min.	Sample: ASTM Indolene Standard ASTM D02 Reference Standard Index	Mass %
	1076.00	115.06	1075.8	0.005								
I41	1077.91	115.28	1077.5	0.017			115.30	1077.2	0.080			
1,3-Dimethyl-2- ethylbenzene	1079.65	115.72	1080.3	0.053	115.74	1080.2	0.045	116.65	1079.4	0.064	115.82	1080.5
I42	1080.68	115.94	1081.7	0.012			116.07	1082.2	0.035	116.00	1081.7	0.168
I43	1081.60	116.32	1084.1	0.017	115.90	1081.3	0.015	116.35	1084.0	0.044		
	1085.30	116.44	1084.9	0.016			116.49	1084.9	0.055			
	1086.54						116.71	1086.3	0.072	116.70	1086.2	0.013
	1088.20	1088.20					117.08	1088.7	0.059	117.00	1088.1	0.024
	1088.88	117.04	1088.8	0.027								
Undecene-1	1090.45	117.30	1090.4	0.035	117.50	1091.5	0.051	117.55	1091.7	0.063	117.40	1090.7
1,4-Methyl-t- butylbenzene	1092.00	117.51	1091.8	0.032								
1,2-Dimethyl-3- ethylbenzene	1093.12	117.66	1092.7	0.115	117.68	1092.7	0.226	117.71	1092.7	0.118	117.74	1092.8
	1094.89	118.05	1095.2	0.016								
	1095.78	118.15	1095.8	0.020	118.34	1097.0	0.015	118.17	1095.6	0.052		
1,2-Ethyl-i- propylbenzene	1097.22	118.34	1097.0	0.014							118.42	1097.1
	1098.54	118.53	1098.3	0.018	118.61	1098.7	0.013	118.61	1098.4	0.034		
n-Undecane	1100.00	118.81	1100.0	0.381	118.82	1100.0	0.009	118.86	1100.0	0.753	118.86	1099.9
	1101.00										119.07	1101.6
1,4-Ethyl-i- propylbenzene	1102.50	119.13	1102.7	0.018								
1,2,4,5- Tetramethylbenzene	1104.83	119.32	1104.4	0.147	119.33	1104.4	0.473	119.36	1104.3	0.073	119.39	1104.4
1,2-Methyl-n- butylbenzene	1107.30											
1,2,3,5- Tetramethylbenzene	1108.79	119.77	1108.3	0.200	119.79	1108.3	0.621	119.82	1108.3	0.073	119.85	1108.3
	1110.82										120.14	1110.8
	1112.39	120.16	1111.5	0.028								
	1115.92	120.65	1115.8	0.023	120.75	1116.4	0.007					
	1120.13	121.18	1120.2	0.023								
	1121.30	121.30	1121.2	0.025								
1,2-Methyl-t- butylbenzene	1122.80							121.36	1121.4	0.015		
	1124.62											
5-Methylindan	1127.35	121.95	1126.7	0.283	121.97	1126.7	0.372	122.09	1127.5	0.019	121.83	1125.1
	1129.83										122.02	1126.7
I43	1131.42	122.52	1131.5	0.026								
4-Methylindan	1133.70	122.75	1133.4	0.072	122.76	1133.4	0.124	122.57	1131.5	0.012	122.52	1130.9
	1134.90	122.90	1134.6	0.031				122.79	1133.4	0.011	122.82	1133.4
	1136.52	123.09	1136.2	0.078	123.11	1136.2	0.161	123.13	1136.2	0.022	123.01	1135.0
1,2-Ethyl-n- propylbenzene	1138.11	123.23	1137.4	0.266								
2-Methylindan	1140.67	123.60	1140.4	0.036	123.56	1140.0	0.009	123.25	1137.4	0.353	123.31	1137.5
1,3-Methyl-n- butylbenzene	1142.70	123.81	1142.1	0.071	123.82	1142.2	0.172	123.85	1142.2	0.019	123.88	1142.2
1,3-Di-i-propylbenzene	1144.27	124.03	1144.0	0.094								
s-Pentylbenzene	1148.00	124.52	1148.0	0.047	124.54	1148.0	0.053				124.10	1144.0
	1149.04	124.62	1148.8	0.049							124.59	1148.0
n-Pentylbenzene	1149.83	124.71	1149.5	0.047	124.73	1149.6	0.034					



TABLE A1.2 Continued

Component	File: DHA94098.D			File: DHA94101.D			File: DHA94105.D			File: DHA94108.D		
	Coopera- tive Study Average RI	Sample: PONA-Va *NJ* Reference Mixture	Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate	Sample: Platformate Auto/Oil C-Matrix Reformate	Sample: No. 1 Ref Naphtha ASTM No. 1 Reference	Sample: ASTM Indolene Standard ASTM D02 Reference	Index	Mass %	Min.	Index	Mass %	Min.
1,1-M-2-(4- MP)cyclopentane	1151.80											
1,2-Di-i-propylbenzene	1153.16	125.11	1152.8	0.042	125.13	1152.8	0.071	125.18	1152.9	0.015	125.18	1152.9
	1154.09	125.23	1153.8	0.056	125.24	1153.8	0.066	125.29	1153.7	0.012	125.29	1153.7
	1157.64	125.65	1157.2	0.045	125.67	1157.3	0.068	125.72	1157.3	0.021	125.72	1157.3
1,4-Di-i-propylbenzene	1159.52	125.91	1159.4	0.110	125.90	1159.2	0.085	125.95	1159.1	0.017	125.95	1159.1
	1161.30							126.29	1161.8	0.013	126.29	1161.8
Tetrahydronaphthalene	1163.30	126.41	1163.3	0.051	126.37	1162.9	0.007	126.61	1164.5	0.021	126.61	1164.5
	1165.13				126.74	1165.9	0.117	126.80	1166.0	0.021	126.80	1166.0
Naphthalene	1166.34	126.73	1166.0	0.060	126.91	1167.3	0.891	126.96	1167.3	0.148	126.96	1167.3
1-t-Butyl-3,5- dimethylbenzene	1168.01	126.89	1167.3	0.285								
	1169.25	127.15	1169.3	0.015								
1,4-Ethyl-t-butylbenzene	1173.72	127.68	1173.6	0.099	127.66	1173.3	0.010	127.81	1174.1	0.057	127.81	1174.1
	1177.88	128.15	1177.3	0.114	128.18	1177.4	0.036	128.15	1176.8	0.043	128.15	1176.8
1,46	1179.46	128.34	1178.8	0.063	128.29	1178.3	0.010					
	1181.20				128.35	1178.8	0.010					
1,47	1183.44	128.85	1182.9	0.081	128.87	1182.9	0.031					
	1187.14	129.31	1186.5	0.068	129.33	1186.6	0.011					
1,3-Di-n-propylbenzene	1188.64	129.53	1188.2	0.075	129.55	1188.3	0.077	129.60	1188.3	0.026	129.60	1188.3
A ₅	1190.24	129.73	1189.8	0.058	129.75	1189.8	0.050	129.80	1189.9	0.022	129.80	1189.9
Dodecene-1	1192.19											
A ₆	1198.52	130.79	1198.1	0.037	130.80	1198.1	0.045	130.86	1198.1	0.017	130.86	1198.1
n-Dodecane	1200.00	131.03	1200.0	0.332				131.10	1200.0	0.014	131.10	1200.0
	1202.51	131.30	1202.6	0.025								
1,3,5-Triethylbenzene	1208.41	131.86	1208.0	0.027	131.84	1207.7	0.009					
	1211.79											
	1216.27	132.68	1215.9	0.030	132.68	1215.8	0.009					
	1217.50	132.83	1217.3	0.061	132.86	1217.4	0.024	132.91	1217.6	0.009	132.91	1217.6
	1222.36	133.32	1222.0	0.060	133.33	1221.9	0.023					
	1223.70	133.52	1223.9	0.055	133.53	1223.9	0.022					
	1225.08	133.66	1225.2	0.025	133.65	1225.0	0.014	133.72	1225.3	0.006	133.72	1225.3
	1228.64	133.97	1228.2	0.040	134.03	1228.6	0.010	134.15	1229.8	0.004	134.15	1229.8
1,2,4-Triethylbenzene	1230.00											
	1230.83											
	1232.23	134.40	1232.2	0.052	134.87	1236.5	0.002					
	1236.42	134.84	1236.4	0.058								
	1237.42	134.98	1237.7	0.012	135.35	1241.0	0.083					
1,4-Methyl-n- pentylbenzene	1241.71	135.34	1241.1	0.110				135.39	1241.1	0.015	135.39	1241.1
	1244.15	135.63	1243.8	0.068								
	1246.48	135.86	1246.0	0.013	135.89	1246.1	0.007					
	1248.73	136.12	1248.4	0.013	136.12	1248.3	0.007					
	1251.16	136.38	1250.8	0.029								
n-Hexylbenzene	1252.85	136.57	1252.5	0.052	136.84	1254.9	0.055					
	1255.61	136.84	1255.1	0.101								
	1257.39	137.11	1257.6	0.029	137.25	1258.7	0.021					
	1259.54	137.25	1258.9	0.050								
	1262.15	137.61	1262.2	0.041								
	1266.71	138.08	1266.5	0.062								
	1269.02	138.31	1268.6	0.030								
1,49	1270.79	138.43	1269.8	0.015	138.85	1273.5	0.067					
1,2,3,4,5- Pentamethylbenzene	1274.04	138.82	1273.3	0.107								
	1277.23							139.25	1277.0	0.010	139.25	1277.0

TABLE A1.2 Continued

Component	Cooperative Study Average RI		File: DHA94099.D Sample: PONA-Va *NJ* Reference Mixture		File: DHA94101.D Sample: HFLA C-Matrix Auto/Oil C-Matrix Alkylate		File: DHA94105.D Sample: No. 1 Ref Naphtha ASTM No. 1 Reference Sample		File: DHA94108.D Sample: ASTM Indolene Standard ASTM D02 Reference Standard	
	Min.	Index	Min.	Mass %	Min.	Index	Min.	Mass %	Min.	Index
2-Methylnaphthalene	1279.96	1279.4	139.49	0.022	139.77	1281.8	0.786	139.82	1282.1	0.059
	1282.57	1281.7	139.75	0.353						
	1286.59	1285.1	140.11	0.010						
	1287.50	1286.0	140.22	0.034						
Tridecene-1	1290.10	1288.2	140.46	0.029	140.44	1296.8	0.377	140.84	1291.5	0.011
	1292.41	1294.6	141.16	0.042						
1-Methylnaphthalene	1295.08	1296.9	141.42	0.152	141.44	1296.8	0.377	141.49	1297.3	0.035
n-Tridecane	1300.00	1300.0	141.77	0.234						
C14+	1300.50							141.85	1300.6	0.007

TABLE A1.3 Repeatability and Reproducibility of DHA Determinations

NOTE 1—The following data has been prepared by statisticians of CS94 in accordance with RR:D02-1007, and represents their best estimate of the cooperative study data. Not all of the data qualified for this evaluation since:

(a) For each analyte to qualify for a precision statement, it must be present in at least six samples, and detected by at least six laboratories, at least once.

(b) The (repeatability standard deviation)/mean value for each analyte/sample combination must be less than or equal to 0.1, in accordance with LOQ requirements which, while not a standard, is what CS94 is recommending.

NOTE 2—

Legend:

- r_{min} = lower 95 % confidence limit of r_{est}
- r_{est} = repeatability estimate in percent of concentration,
- r_{max} = upper 95 % confidence limit of r_{est}
- $R_{min}, R_{est}, R_{max}$ = for reproducibility,
- C_{min} = lower concentration limit that r_{est}, R_{est} is applicable, and
- C_{max} = upper concentration limit that r_{est}, R_{est} is applicable.

Component	Average RI	r_{min}	r_{est}	r_{max}	R_{min}	R_{est}	R_{max}	C_{min}	C_{max}
i-Butane	366.15	5.6	10.1	16.4	22.8	46.1	81.5	0.06	0.38
Butene-1	390.72	6.4	11.1	17.6	31.2	64.5	115.8	0.01	0.14
n-Butane	400.00	6.8	9.9	13.9	15.3	32.4	59.1	1.02	3.75
2,2-Dimethylpropane	415.10	3.3	8.8	18.6	32.1	50.1	73.7	0.01	0.02
i-Pentane	477.45	5.9	7.2	8.7	8.5	14.8	23.8	2.48	13.38
Pentene-1	490.83	5.2	7.5	10.5	9.7	13.8	19.0	0.06	0.43
2-Methylbutene-1	496.66	4.9	6.9	9.4	8.3	12.9	19.0	0.14	0.86
n-Pentane	500.00	5.2	6.5	8.1	7.1	10.4	14.8	1.06	3.49
t-Pentene-2	510.56	4.5	6.5	9.0	7.2	10.3	14.4	0.28	1.16
c-Pentene-2	519.53	4.7	6.3	8.1	7.6	13.2	20.9	0.16	0.63
2-Methylbutene-2	524.92	4.3	6.0	8.1	7.8	11.4	15.9	0.50	1.85
1,3-Pentadiene	527.97	6.7	14.0	25.3	17.0	25.3	35.9	0.01	0.06
2,2-Dimethylbutane	540.54	3.1	4.7	6.8	6.4	9.9	14.6	0.08	2.18
Cyclopentene	557.21	4.0	5.8	8.1	7.6	10.5	13.9	0.07	0.27
4-Methylpentene-1	562.02	2.7	4.2	6.1	8.9	11.3	14.0	0.02	0.09
3-Methylpentene-1	562.81	3.5	5.0	6.9	6.1	8.8	12.0	0.03	0.12
Cyclopentane	566.84	3.8	4.9	6.2	7.0	10.1	14.0	0.07	0.59
2,3-Dimethylbutane	569.24	2.9	3.2	3.5	5.1	8.5	13.1	0.70	1.91
2-Methylpentane	573.70	2.5	2.9	3.4	5.1	6.6	8.4	1.06	5.80
4-Methyl-t-pentene-2	575.47	6.5	8.7	11.3	18.1	28.0	41.1	0.08	0.28
3-Methylpentane	585.52	2.6	2.9	3.2	4.0	5.6	7.5	0.60	2.50
2-Methylpentene-1	590.19	2.3	2.7	3.2	3.8	5.4	7.4	0.11	0.45



TABLE A1.3 Continued

Component	Average RI	r _{min}	r _{est}	r _{max}	R _{min}	R _{est}	R _{max}	C _{min}	C _{max}
Hexene-1	591.06	3.1	4.2	5.5	5.9	8.2	10.9	0.06	0.26
n-Hexane	600.00	2.0	2.4	2.9	3.6	5.1	6.9	0.33	2.52
t-Hexene-3	602.83	1.9	3.2	4.9	7.4	11.8	17.7	0.08	0.35
t-Hexene-2	605.44	2.5	2.8	3.2	4.5	6.4	8.9	0.16	0.71
2-Methylpentene-2	607.86	2.4	3.0	3.8	6.1	13.1	24.1	0.22	0.97
3-Methyl-c-pentene-2	610.54	2.1	2.5	2.9	5.3	6.9	8.7	0.11	0.48
3-Methylcyclopentene	611.61	3.0	4.6	6.8	10.1	12.8	15.9	0.02	0.10
c-Hexene-2	614.67	2.6	3.3	4.1	4.8	5.4	6.1	0.09	0.40
3,3-Dimethylpentene-1	620.91	1.9	2.5	3.1	4.3	5.7	7.5	0.17	0.75
2,2-Dimethylpentane	624.17	3.3	4.3	5.5	4.4	8.2	13.7	0.01	0.09
Methylcyclopentane	625.86	2.2	2.6	3.1	4.5	6.4	8.7	0.37	2.35
2,4-Dimethylpentane	630.60	1.6	2.5	3.8	4.2	5.3	6.6	0.20	1.94
2,2,3-Trimethylbutane	634.86	3.2	4.1	13.1	12.5	20.4	31.1	0.02	0.08
3,4-Dimethylpentene-1	642.87	5.5	7.1	30.6	9.7	22.3	43.0	0.01	0.03
4,4-Dimethyl-c-pentene-2	646.65	3.8	5.6	8.0	6.1	11.0	17.9	0.01	0.11
2,4-Dimethylpentene-1	647.67	3.8	6.7	10.8	11.9	13.5	15.3	0.01	0.04
1-Methylcyclopentene	648.71	1.9	2.7	3.7	7.9	8.7	9.6	0.17	0.82
Benzene	649.92	2.6	3.6	4.8	5.5	9.0	13.7	0.17	1.58
5-Methylhexene-1	655.56	5.3	7.5	10.1	18.0	30.4	47.6	0.01	0.22
Cyclohexane	657.81	2.7	3.7	4.9	8.2	14.8	24.3	0.07	0.90
2-Methyl-t-hexene-3	661.03	3.2	6.3	11.0	16.8	23.4	31.6	0.03	0.14
2-Ethyl-3-Methylbutene-1	662.60	4.1	8.9	16.5	89.5	117.0	149.7	0.01	0.04
4-Methylhexene-1	663.81	2.5	4.5	7.2	11.7	14.7	18.3	0.02	0.09
4-Methyl-t-c-hexene-2	666.23	2.1	3.2	4.7	5.4	7.3	9.5	0.05	0.28
2-Methylhexane	667.61	1.6	2.2	2.9	5.1	6.1	7.2	0.39	1.09
2,3-Dimethylpentane	668.84	1.6	2.3	3.2	5.4	6.4	7.5	0.33	3.16
3-Methylhexane	675.89	1.9	2.8	4.0	4.7	5.6	6.7	0.37	1.08
3-4-Dimethyl-c-pentene-2	679.46	3.6	5.9	9.1	11.6	23.4	41.2	0.02	0.14
1c,3-dimethylcyclopentane	681.68	1.7	2.7	4.0	5.4	7.5	10.1	0.11	0.56
1t,3-Dimethylcyclopentane	684.37	1.7	2.8	4.4	5.4	7.7	10.5	0.08	0.48
3-Ethylpentane	685.98	3.1	3.7	4.3	5.6	8.3	11.9	0.08	0.26
2,2,4-Trimethylpentane	688.48	2.4	3.2	4.1	7.4	11.4	16.7	0.10	11.26
3-Methyl-c-hexene-3	694.82	3.2	6.5	11.6	7.3	12.9	20.8	0.03	0.17
t-Heptene-3	698.39	2.6	3.2	3.8	5.0	8.1	12.3	0.11	0.67
n-Heptane	700.00	2.5	3.4	4.5	7.7	10.8	14.7	0.21	1.06
3-Methyl-c-hexene-2	702.30	1.5	2.5	4.0	4.6	7.2	10.6	0.13	0.75
3-Methyl-t-hexene-3	702.99	2.2	4.1	6.7	8.1	10.1	12.3	0.05	0.26
t-Heptene-2	704.58	2.6	4.2	6.4	5.9	7.1	8.4	0.06	0.34
3-Ethylpentene-2	705.96	4.7	8.3	13.4	14.5	17.9	21.8	0.03	0.16
c-Heptene-2	708.82	1.9	3.2	5.0	7.0	7.8	8.7	0.12	0.63
2,3-Dimethylpentene-2	712.07	2.4	3.7	5.5	7.8	9.1	10.5	0.06	0.57
O29	715.67	6.8	11.5	18.0	15.7	22.6	31.3	0.01	0.08
1c,2-Dimethylcyclopentane	717.13	2.4	3.7	5.3	9.5	11.6	13.9	0.05	0.20
Methylcyclohexane	717.89	2.8	3.4	4.0	4.1	5.9	8.2	0.11	1.20
2,2-Dimethylhexane	720.70	4.2	7.2	11.3	10.2	15.4	22.3	0.02	0.10
2,5-Dimethylhexane	730.05	2.5	3.0	3.5	4.9	6.2	7.7	0.16	1.12
2,4-Dimethylhexane	731.84	3.5	4.3	5.2	6.5	9.0	12.1	0.29	1.39
1c,2t,4-Trimethylcyclopentane	737.11	3.0	4.2	5.6	6.9	7.9	9.0	0.03	0.17
2,3,4-Trimethylpentane	746.83	2.3	3.8	5.8	5.8	7.8	10.3	0.08	4.26
I1	747.91	2.2	6.2	13.4	10.1	21.1	38.1	0.09	0.59
Toluene	751.77	1.9	2.7	3.8	10.8	13.5	16.5	1.99	10.34
2,3-Dimethylhexane	757.87	2.2	3.7	5.7	5.0	6.9	9.2	0.22	1.23
1,1,2-Trimethylcyclopentane	760.33	7.6	13.5	21.8	13.6	25.7	43.4	0.02	0.26
O44	761.73	4.2	9.4	17.7	12.7	20.5	30.8	0.03	0.24
2-Methylheptane	764.14	3.5	4.9	6.6	4.8	6.1	7.5	0.15	0.63
4-Methylheptane	765.62	4.1	6.2	9.1	7.1	9.1	11.5	0.05	0.29
3-Methyl-3-ethylpentane	766.62	5.4	7.1	9.3	11.0	15.8	21.8	0.05	0.10
3-Methylheptane	771.78	2.1	3.4	5.3	4.4	5.1	5.9	0.13	0.71

TABLE A1.3 Continued

Component	Average RI	r_{min}	r_{est}	r_{max}	R_{min}	R_{est}	R_{max}	C_{min}	C_{max}
1c,2t,3-Trimethylcyclopentane	772.98	3.4	4.4	5.7	5.9	7.4	9.1	0.06	0.29
3-Ethylhexane	773.76	6.9	10.8	15.9	13.1	23.8	39.1	0.01	0.07
1t,4-Dimethylcyclohexane	774.89	9.8	18.0	29.8	22.2	49.5	93.1	0.01	0.11
2,2,5-Trimethylhexane	782.93	3.1	4.3	5.7	5.7	8.0	10.8	0.14	2.21
3c-Ethylmethylcyclopentane	784.35	5.7	12.1	22.1	9.7	22.2	42.5	0.05	0.21
3t-Ethylmethylcyclopentane	786.55	7.4	9.7	12.4	8.4	22.1	46.0	0.02	0.11
2t-Ethylmethylcyclopentane	787.86	8.9	11.4	14.2	12.6	26.7	48.7	0.02	0.12
1t,2-Dimethylcyclohexane	792.77	3.7	6.7	11.0	7.0	10.1	14.0	0.04	0.29
t-Octene-4	794.21	12.7	14.5	16.6	16.7	21.9	28.1	0.02	0.18
1c,2c,3-Trimethylcyclopentane	797.25	3.5	6.0	9.4	6.3	9.2	12.9	0.07	0.51
1t,3-Dimethylcyclohexane	798.80	6.8	9.3	12.4	11.9	16.3	21.7	0.02	0.10
n-Octane	800.00	2.2	3.6	5.5	6.5	15.7	30.9	0.14	0.75
Octene-2	804.40	4.9	8.2	12.7	11.1	20.1	32.9	0.03	0.23
I2	806.39	4.7	10.9	20.9	8.7	19.7	37.4	0.05	0.36
i-Propylcyclopentane	808.06	4.1	10.7	22.0	8.4	19.8	38.4	0.03	0.18
2,3,4-Trimethylhexane	818.10	2.5	5.0	8.9	5.1	7.2	9.8	0.05	0.37
N2	819.93	7.7	11.5	16.4	6.8	17.4	35.7	0.01	0.06
N3	822.29	8.0	13.3	20.7	11.3	23.2	41.5	0.01	0.09
2,3,5-Trimethylhexane	827.51	3.9	5.4	7.1	7.5	31.4	82.4	0.07	0.12
1,1,4-Trimethylcyclohexane	832.56	8.6	17.1	29.9	21.7	36.0	55.6	0.03	0.26
2,2,3-Trimethylhexane	834.96	8.4	13.1	19.2	14.8	20.3	27.0	0.05	0.09
2,5-Dimethylheptane	842.63	6.8	8.8	11.1	7.8	12.3	18.2	0.11	0.18
Ethylbenzene	854.65	2.2	3.2	4.4	7.2	10.6	14.9	0.62	2.62
1,3-Dimethylbenzene	864.22	2.6	3.3	4.2	9.7	12.5	15.7	1.55	6.66
1,4-Dimethylbenzene	865.20	3.6	4.2	5.0	10.4	14.1	18.5	0.62	2.97
I5	870.95	6.6	11.5	18.3	13.8	28.4	50.9	0.02	0.13
4-Methyloctane	873.81	4.4	7.6	12.0	5.9	11.2	18.9	0.05	0.20
2-Methyloctane	874.76	4.6	8.2	13.3	6.0	10.4	16.6	0.07	0.35
3-Ethylheptane	879.11	8.5	13.4	20.1	27.7	38.7	52.5	0.02	0.09
3-Methyloctane	880.24	5.1	8.5	13.0	8.7	15.5	24.9	0.07	0.29
1,2-Dimethylbenzene	883.47	2.1	3.2	4.7	8.8	11.5	14.8	0.83	3.85
I6	885.34	10.2	17.0	26.4	23.3	44.5	75.8	0.02	0.06
I7	886.38	6.5	9.0	12.0	7.8	21.1	44.7	0.05	0.32
N22	895.99	7.9	16.5	29.6	16.7	29.4	47.2	0.03	0.22
N23/c-nonene-2	897.24	5.7	15.9	34.2	26.1	48.7	81.6	0.02	0.15
I10	898.70	4.5	12.3	26.0	9.3	31.1	73.3	0.04	0.44
n-Nonane	900.20	3.9	6.4	9.8	8.6	10.3	12.2	0.06	0.34
i-Propylbenzene	912.28	3.2	5.0	7.3	8.3	15.1	24.9	0.04	0.33
N27	914.45	3.6	12.3	29.2	9.0	21.1	40.8	0.02	0.14
I12	921.30	4.4	11.0	22.2	8.7	21.2	42.2	0.03	0.34
2,4-Dimethyloctane	924.39	6.1	12.1	21.3	16.8	26.0	38.1	0.03	0.11
2,6-Dimethyloctane	930.83	7.0	14.6	26.4	15.4	27.7	45.3	0.02	0.10
2,5-Dimethyloctane	932.66	5.6	11.2	19.6	15.3	22.0	30.4	0.04	0.13
3,3-Dimethyloctane	942.30	4.3	10.4	20.6	7.5	17.5	34.0	0.03	0.11
n-Propylbenzene	946.33	2.8	5.0	8.1	7.6	11.9	17.7	0.21	0.77
3,6-Dimethyloctane	948.31	7.6	14.9	25.7	18.1	38.9	71.4	0.01	0.04
1,3-Methylethylbenzene	954.42	3.7	5.2	7.2	7.6	10.2	13.3	0.81	2.61
1,4-Methylethylbenzene	956.22	3.5	5.3	7.7	5.1	7.7	11.1	0.32	1.19
1,3,5-Trimethylbenzene	961.92	3.7	5.5	7.7	5.4	8.3	12.1	0.39	1.21
2-Methylnonane	971.77	6.5	10.6	16.2	17.5	25.9	36.6	0.03	0.19
3-Methylnonane	977.26	7.5	13.5	22.1	23.5	41.0	65.5	0.03	0.16
I18	980.12	7.3	15.7	28.8	14.9	30.0	52.9	0.04	0.82
1,2,4-Trimethylbenzene	983.40	4.2	5.7	7.5	7.8	10.6	13.9	1.19	4.32
I21	987.40	6.9	18.6	39.3	16.9	49.6	109.2	0.03	0.27
I24	994.53	6.9	12.2	19.6	18.9	31.1	47.7	0.04	0.36
n-Decane	1000.20	7.5	9.2	11.1	12.1	17.9	25.3	0.03	0.25
N38	1003.39	9.7	19.8	35.2	25.8	47.6	79.3	0.01	0.12
1,2,3-Trimethylbenzene	1006.88	3.8	5.8	8.5	7.2	8.5	10.0	0.28	0.96



TABLE A1.3 Continued

Component	Average RI	r _{min}	r _{est}	r _{max}	R _{min}	R _{est}	R _{max}	C _{min}	C _{max}
2,3-dihydroindene	1019.44	5.2	7.8	11.3	8.8	10.8	13.0	0.18	0.37
I30	1024.82	7.5	13.9	23.3	20.4	61.8	138.4	0.01	0.15
1,2-Methyl-i-propylbenzene	1027.73	17.3	19.5	22.0	66.4	99.4	141.9	0.02	0.08
?	1038.53	6.8	16.0	31.1	16.8	37.4	70.2	0.02	0.16
1,3-Diethylbenzene	1039.97	5.0	7.3	10.1	9.9	13.7	18.5	0.08	0.22
1,3-Methyl-n-propylbenzene	1042.60	4.0	6.4	9.5	6.7	13.5	23.6	0.30	0.68
1,4-Methyl-n-propylbenzene	1046.40	4.6	8.7	14.7	7.8	12.7	19.4	0.16	0.32
1,3-Dimethyl-5-ethylbenzene	1049.78	3.4	6.0	9.8	7.6	10.6	14.3	0.22	0.51
1,2-Methyl-n-propylbenzene	1057.87	6.3	9.3	13.2	20.9	30.3	42.1	0.11	0.20
		4.5	11.4	23.2	14.4	36.1	73.1	0.06	0.56
I38	1065.53	7.9	16.6	30.0	28.4	37.4	48.2	0.02	0.09
1,4-Dimethyl-2-ethylbenzene	1068.05	4.1	6.7	10.3	15.3	27.8	45.7	0.15	0.38
1,2-Dimethyl-4-ethylbenzene	1075.25	5.1	7.0	9.3	13.8	18.7	24.6	0.29	0.71
Undecene-1	1090.45	8.2	17.8	32.9	20.5	35.4	56.1	0.01	0.09
1,2-Dimethyl-3-ethylbenzene	1093.12	6.8	10.3	14.8	15.2	18.4	22.1	0.09	0.19
1,2-Ethyl-i-propylbenzene	1097.22	14.3	31.3	58.2	51.7	74.6	103.4	0.02	0.06
n-Undecane	1100.00	8.6	13.9	21.0	24.4	40.0	61.2	0.03	0.18
1,2,4,5-Tetramethylbenzene	1104.83	6.1	8.3	11.1	12.5	16.0	20.1	0.15	0.36
1,2,3,5-Tetramethylbenzene	1108.79	6.4	7.8	9.3	10.2	13.9	18.3	0.21	0.51
5-Methylindan	1127.35	6.6	9.2	12.4	7.5	9.3	11.4	0.06	0.34
I43	1131.42	9.1	15.0	23.0	18.8	30.8	47.1	0.02	0.35
4-Methylindan	1133.70	8.0	13.8	21.9	15.3	20.9	27.6	0.02	0.10
1,2-Ethyl-n-propylbenzene	1136.52	7.4	12.0	18.3	25.3	39.6	58.4	0.02	0.11
2-Methylindan	1138.11	5.6	7.7	10.2	6.4	8.9	12.0	0.08	0.34
1,3-Di-i-propylbenzene	1142.70	7.9	9.5	11.2	16.2	17.9	19.7	0.06	0.18
1,2-Di-i-propylbenzene	1153.16	13.9	22.4	34.0	21.3	36.0	56.2	0.01	0.06
?	1157.64	12.8	20.7	31.4	24.5	44.1	71.9	0.02	0.06
1,4-Di-i-propylbenzene	1159.52	11.1	18.6	28.8	18.6	35.6	60.8	0.01	0.09
Naphthalene	1168.01	6.1	8.5	11.3	12.9	16.9	21.5	0.13	0.40
1,4-Ethyl-t-butylbenzene	1173.72	9.8	12.0	14.5	18.0	28.1	41.3	0.04	0.30
I48	1187.14	9.1	14.3	21.1	16.8	30.7	50.8	0.01	0.09
1,3-Di-n-propylbenzene	1188.64	9.9	14.4	19.9	18.9	24.2	30.4	0.02	0.08
A5	1190.24	11.5	19.2	29.8	26.7	30.3	34.2	0.02	0.06
A6	1198.52	13.7	23.3	36.7	28.1	39.5	53.5	0.01	0.05
n-Dodecane	1200.00	12.2	16.7	22.1	20.2	32.9	50.0	0.01	0.11
1,4-Methyl-n-pentylbenzene	1241.71	8.2	14.1	22.3	16.5	31.2	52.7	0.01	0.14
1,2,3,4,5-Pentamethylbenzene	1274.04	11.4	13.9	16.7	23.1	29.7	37.5	0.01	0.11
2-Methylnaphthalene	1282.57	7.6	11.1	15.4	17.5	22.3	28.0	0.05	0.50
1-Methylnaphthalene	1297.72	7.3	11.0	15.8	14.0	21.0	30.1	0.02	0.22




FIG. A1.1 Column Retention Factor Calculation (9.1)

Efficiency (n): $5.545(10.124 / 0.032)^2 = 555,016$
Resolution (R): $2(10.817 - 10.547) / 1.699(0.034 + 0.034) = 4.673$




FIG. A1.2 Column Efficiency and Resolution Calculations (9.2 and 9.3)







FIG. A1.3 Column Inertness - Peak Skewness Calculation (11.5)



0.5 μm_{df} methylsilicone column + 1.25 m 1 μm_{df} precolumn



0.5 μm_{df} methylsilicone column + 1.5 m 1 μm_{df} precolumn



0.5 μm_{df} methylsilicone column + 2.0 m 1 μm_{df} precolumn

FIG. A1.4 Oxygenates Separations - Effect of Different Precolumn Lengths

$$R = 2(21.658 - 21.590) / 1.699(0.08 + 0.08) = 0.500$$




FIG. A1.5 PONA-V Standard—Analysis through Benzene New DHA Column—Analyzed at 35 °C Isothermal (Conditions in Accordance with Table 2)

$$R = 2(21.025 - 20.925) / 1.699(0.076 + 0.081) = 0.750$$




FIG. A1.6 PONA-V Standard—Analysis through Benzene New DHA Column plus 1 m × 0.25 mm 1 μm Precolumn Analyzed at 35 °C Isothermal

$$R = 2(21.258 - 21.125) / 1.699(0.079 + 0.081) = 0.978$$




FIG. A1.7 PONA-V Standard—Analysis through Benzene New DHA Column plus 2 m × 0.25 mm 1 μm Precolumn Analyzed at 35 °C Isothermal

$$R = 2(22.642 - 22.458) / 1.699(0.089 + 0.093) = 1.190$$




FIG. A1.8 DHA Calibration Standard—Analysis through Benzene New DHA Column plus 3 m × 0.25 mm 1 μm Precolumn Analyzed at 35 °C Isothermal







FIG. A1.9 Key Separations—Effect of Different Precolumn Lengths Same Primary Column. Conditions in Accordance with Table 2 Top to Bottom—1.00 m, 1.25 m, 1.50 m, and 2.00 m Precolumn



(Old) 0.5 μm_{df} methylsilicone column + 1.25 m 1 μm_{df} precolumn




0.5 μm_{df} methylsilicone column + 2.4 m 1 μm_{df} precolumn




100 m \times 0.25 mm HP 0.5 μm_{df} methylsilicone column + 3.0 m 1 μm_{df} precolumn


FIG. A1.10 Key Separations—Tuning of Different Columns Conditions in Accordance with Table 2



PONA-Va Mixture




RFA Gasoline




CCF Gasoline


FIG. A1.11 DHA Analyses—Methane through Hexane



PONA-Va Mixture




RFA Gasoline




CCF Gasoline


FIG. A1.12 DHA Analyses—Hexane through Heptane



PONA-Va Mixture




RFA Gasoline




CCF Gasoline


FIG. A1.14 DHA Analyses—Octane through Nonane



PONA-Va Mixture




RFA Gasoline




CCF Gasoline


FIG. A1.15 DHA Analyses—Nonane through Decane



PONA-Va Mixture




RFA Gasoline




CCF Gasoline


FIG. A1.16 DHA Analyses—Decane through Dodecane



PONA-Va Mixture



RFA Gasoline



CCF Gasoline

FIG. A1.17 DHA Analyses—Dodecane through Tetradecane

TABLE A1.4 Benzene

Sample	Benzene (wt %)	
	D5580	D6730
2	1.52	1.58
6	1.05	1.12
8	1.10	1.15
10	1.13	1.19
13	0.14	0.17
14	0.62	0.69
Average	0.93	0.98

TABLE A1.5 Toluene

Sample	Toluene (wt %)	
	D5580	D6730
2	4.3	4.5
6	2.1	2.0
8	10.1	10.3
10	5.0	5.2
13	3.3	3.3
14	4.4	4.7
Average	4.9	5.0

TABLE A1.6 Total Aromatics

Sample	Total Aromatics (wt %)		
	D5580	PIONA ^A	D6730
2	30.3	28.2	30.2
6	18.9	18.7	18.3
8	49.1	49.0	47.6
10	23.9	24.5	23.1
13	19.7	19.8	19.3
14	23.8	24.6	24.2
Average	27.6	27.5	27.1

^A Multidimensional PIONA

TABLE A1.7 Total Olefins

Sample	Total Olefins (wt %)	
	PIONA ^A	D6730
2	7.1	4.5
6	9.8	8.7
8	6.6	6.1
10	15.1	12.9
13	11.1	10.6
14	24.6	19.5
Average	12.4	10.9

^A Multidimensional PIONA

TABLE A1.8 Total Oxygenates

Sample	Total Oxygenates (wt %)	
	PIONA ^A	D6730
2 ^B	15.3	15.1
6 ^B	7.0	7.8
8 ^B	4.2	4.3
10 ^C	>8	10.5
13 ^B	20.5	20.2
14 ^B	2.8	2.9
Average	N/A	10.1

^A Multidimensional PIONA

^B Major oxygenate = MTBE

^C Major oxygenate = Ethanol

TABLE A1.9 Total Paraffins and Total Naphthenes

Sample	Total Paraffins (wt %)		Total Naphthenes (wt %)	
	PIONA ^A	D6730	PIONA ^A	D6730
8	35.6	38.0	2.2	2.7
10	41.1	45.5	5.6	6.5
13	42.6	46.0	1.3	2.1
14	34.1	41.3	5.9	9.3
Average	38.4	42.7	3.8	5.2

^A Multidimensional PIONA

APPENDIX

(Nonmandatory Information)

X1. BIBLIOGRAPHY

X1.1 The following publications on DHA analyses may be useful as background and are recommended to the user of these test methods.

X1.1.1 Johansen, N.G., and Ettore, L.S., "Retention Index Values of Hydrocarbons on Open Tubular Columns Coated with Methyl Silicone Liquid Phases," *Chromatographia*, Vol 5, No. 10, October 1982.

X1.1.2 Johansen, N.G., Ettore, L.S., and Miller, R.L., "Quantitative Analysis of Hydrocarbons by Structural Group Type in Gasolines and Distillates. Part 1," *Journal of Chromatography*, 256, 1983, pp. 393-417.

X1.1.3 Kopp, V.R., Bones, C.J., Doerr, D.G., Ho, S.P., and Schubert, A.J., "Heavy Hydrocarbon/Volatility Study: Fuel Blending and Analysis for the Auto/Oil Air Quality Improvement Research Program," SAE Paper No. 930143, March 1993.


X1.1.4 Schubert, A.J., and Johansen, N.J., "Cooperative Study to Evaluate a Standard Test Method for the Speciation of Gasolines by Capillary Gas Chromatography," SAE Paper No. 930144, March 1993.

X1.1.5 Di Sanzo, F. P., and Giarrocco, V. G., "Analysis of Pressurized Gasoline-Range Liquid Hydrocarbon Samples by Capillary Column and PIONA Analyzer Gas Chromatography," *Journal of Chromatographic Science*, Vol 26, June 1988, pp. 258-266.

X1.1.6 Durand, J. P., Beboluene, J. J., and Ducrozet, A., "Detailed Characterization of Petroleum Products with Capillary GC Analyzers," *Analisis*, 23, 1995, pp. 481-483.

X1.1.7 Canadian General Standards Board: CAN/CGSB –3.0, No. 14.3-94, "Test Method for Individual Hydrocarbon Component Analysis (IHA) in Spark Ignition Engine Fuels by Gas Chromatography."

X1.1.8 French Standard NF N07-086, December 1995, "Determination of Hydrocarbon Type Contents in Motor Gasolines from Detailed Analysis Capillary Gas Chromatography."

 **D6730 – 01 (2016)**

ASTM International takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.

This standard is copyrighted by ASTM International, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (www.astm.org). Permission rights to photocopy the standard may also be secured from the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923, Tel: (978) 646-2600; <http://www.copyright.com/>