

Standard Guide for Ecological Considerations for the Use of Oil Spill Dispersants in Freshwater and Other Inland Environments, Ponds and Sloughs¹

This standard is issued under the fixed designation F1209; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ε) indicates an editorial change since the last revision or reapproval.

1. Scope

- 1.1 This guide covers the use of oil spill dispersants to assist in the control of oil spills. The guide is written with the goal of minimizing the environmental impacts of oil spills; this goal is the basis on which the recommendations are made. Aesthetic and socioeconomic factors are not considered, although these and other factors are often important in spill response.
- 1.2 Spill responders have available several means to control or clean up spilled oil. In this guide, the use of dispersants is given equal consideration with other spill countermeasures. It is not considered as a "last resort" after all other methods have failed.
- 1.3 This is a general guide only. Oil, as used in this guide, includes crude oils and refined petroleum products. Differences between individual dispersants or between different oil products are not considered.
- 1.4 The guide is organized by habitat type, for example, small ponds and lakes, rivers and streams, and land. It considers the use of dispersants primarily to protect habitats from impact (or to minimize impacts).
- 1.5 This guide applies only to freshwater and other inland environments. It does not consider the direct application of dispersants to subsurface waters.
- 1.6 In making dispersant use decisions, appropriate government authorities should be consulted as required by law.
- 1.7 The values stated in SI units are to be regarded as standard. No other units of measurement are included in this standard.
- 1.8 This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appro-

priate safety and health practices and determine the applicability of regulatory limitations prior to use.

2. Referenced Documents

2.1 ASTM Standards:²

F2532 Guide for Determining Net Environmental Benefit of Dispersant Use

3. Significance and Use

- 3.1 This guide is meant to aid local and regional response teams who may use it during spill response planning and spill events.
- 3.2 This guide should be adapted to site specific circumstance.

4. Environment Covered—Ponds and Sloughs

- 4.1 Ponds and sloughs are small isolated water bodies that are not part of major water systems. They have a dynamic ecology, and a wide mixture of animal and plant species. In northern regions, these water bodies may freeze to the bottom and do not sustain a wide variety of aquatic species.
- 4.2 While most of these bodies are naturally occurring and exist throughout the year, some may be man-made. In arid climates, the existence of these bodies may be seasonal.
 - 4.3 The characteristics of these water bodies are:
 - 4.3.1 Open water area of less than 10 hectares,
 - 4.3.2 Shallow water with a maximum depth of 1 to 1.5 m,
 - 4.3.3 Soft bottom with a high organic content,
 - 4.3.4 May have alkaline or acidic water,
 - 4.3.5 No well defined inlet or outlet, and
- 4.3.6 Poorly defined shoreline consisting of floating mats of vegetation.

5. Background

5.1 The effects of oil and dispersed oil on these aquatic environments have been the subject of numerous studies. The

¹ This guide is under the jurisdiction of ASTM Committee F20 on Hazardous Substances and Oil Spill Responseand is the direct responsibility of Subcommittee F20.13 on Treatment.

Current edition approved March 1, 2014. Published March 2014. Originally approved in 1989. Last previous edition approved in 2008 as F1209 – 08. DOI: 10.1520/F1209-14.

² For referenced ASTM standards, visit the ASTM website, www.astm.org, or contact ASTM Customer Service at service@astm.org. For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.

studies have involved both intentional experimental spills and studies undertaken during actual spill situations (1-6).

- 5.2 There have been a number of studies on the impact of oil and oil/dispersant mixtures on microbiological systems and plankton (7-16).
- 5.3 The principal biotic components of such water bodies are a variety of fauna and flora. The aquatic flora include algae (planktonic and attached) and floating or submerged vascular plants. Terrestrial flora include grasses, moss, lichens, herbs, forbs, and woody plants.
- 5.4 The fauna include invertebrates (molluscs, crustaceans, worms), fish, a variety of waterfowl and seabirds (ducks, loons, gulls, terns, and herons), mammals, such as beaver and muskrat, and in many areas, significant human activity. There have been some studies on the effects of oil and oil/dispersant mixtures on fauna (17, 18). The distribution and composition of species is a function of climate, local geography and soil type, and human use of the area (19).
- 5.5 Studies of the effects of oil on vegetation include laboratory work on the effects on willows, moss, lichen, black spruce, and sedge. These studies showed that the pouring of oil directly on the vegetation was lethal to willow and moss, and caused some nonpermanent damage to the other species (20-29).
- 5.6 Floating leaves are wetted and penetrated by oil from slicks while younger floating leaves whose cuticles are intact are more resistant to penetration by oil. Penetration occurs most readily in areas where there is damage to the leaf caused by insects or mechanical abrasion.
- 5.7 Oil is retained in areas of dense vegetation and is released slowly. Dispersants applied soon after a spill appear to speed the movement of the slick through the vegetation if there is some water movement. Only the oil that contacts the leaves causes some damage. Leaves and stems that are subsurface are not damaged by the presence of oil. Oil slicks that are restricted to contact of the stem areas near the water surface are not very injurious.
- 5.8 Field studies in northern environments have shown little long term effects of spilled oil. There have been few studies on the effects of dispersed oil in these environments.
- 5.9 Although oil may cause extensive damage to vegetation growing in low-lying wetlands and shorelines of bog lakes, plants that grow from rhizomes or healthy portions of aerial

- tissue (sedge and willow) have a high recovery potential. Long-term damage to root systems could, however, slow recovery of the vegetation in impacted areas. This may be of less importance in areas with an elevated water table (bog lakes).
- 5.10 Laboratory studies of the foliar damage to moss, lichen, willow, blackspruce, and sedge by dispersants indicate some damage, but this is generally not lethal. Plants treated with dispersants recover in most cases. However, foliar damage may increase the plant's susceptibility to pathogens.
- 5.11 Field studies on the effects of the direct spraying of plants with dispersants indicated that the short-term effect of dispersants to individual plants was less than that incurred through natural causes. No long term effects were found and seasonal growth was not affected.
- 5.12 Studies of the effects of dispersants and dispersed oil on the microbiological community show that the characteristics of the dispersant are critical. The dispersant should have no residues or components toxic to microbes or microbial processes.

6. General Considerations for Making Dispersant Use Decisions

- 6.1 The dispersant use decision is, in this case as most others, one of trade-offs. The use of dispersants can reduce the adverse effects of spilled oil on certain biological species at the expense of other components of the ecosystem.
- 6.2 Guide F2532 should be followed before making a decision to use dispersants in a river or creek.
- 6.3 In most cases, the mortality of individual creatures is of less concern than the destruction of habitat. The repopulation of areas after the spill will occur naturally when an area becomes a suitable habitat for a given species.

7. Recommendations

7.1 Dispersant use in ponds and sloughs, and their bordering vegetation, should be considered if a spill poses a significant threat to indigenous wildlife or its habitat. In evaluating the potential for dispersant use, consideration should be given to the alternatives of leaving the oil untreated or the use of mechanical recovery equipment. In many cases, a spill response operation can cause serious damage to a pond or slough habitat, or a disruption of nesting and breeding activities.

8. Keywords

8.1 dispersants; environmental sensitivity; freshwater; inland; oil spill; oil spill dispersants; ponds; sloughs

³ The boldface numbers in parentheses refer to the list of references at the end of this guide.



REFERENCES

- (1) Miller, M. C., Vestal, J. R., Mozley, S., Butler, M., and Hobbie, J. E., "Effects of Prudhoe Crude Oil Spills on Coastal Tundra Ponds," *Energy/Environment II. U.S. EPA-600/9-77-012*, 1977, pp. 521–529.
- (2) Brown, H. M., Goudey, J. S., Fogh, J. M., Cheng, S. K., Dale, M., Hoddinott, J., Quaife, L. R., and Westlake, D. W. S., "Dispersion of Spilled Oil in Freshwater Systems: Field Trial of a Chemical Dispersant," Oil and Chemical Pollution, Vol 6, 1990, pp.37–54.
- (3) Nagy, E., Scott, B. F., and Hart, J., "The Fate of Oil and Oil-Dispersant Mixtures in Freshwater Ponds," *Science of the Total Environment*, Vol. 35, No. 2, 1984, pp. 115–133.
- (4) Scott, B. F., and Glooschenko, V., "Impact of Oil and Oil-Dispersant Mixtures on Flora and Water Chemistry in Freshwater Ponds," Science of the Total Environment, Vol 35, No. 2, 1984, pp. 169–190.
- (5) Scott, B. F., Nagy, E., Sherry, J. P., Dutka, B. J., Glooschenko, V., Snow, N. B., and Wade, P. J., "Ecological Effects of Oil-Dispersant Mixtures in Fresh Water," *Proceedings of 1979 Oil Spill Conference*, Los Angeles, pp. 565–571.
- (6) Debruyn, A.M.H., Wernick, B.G., Stefura, C., Mcdonald, B.G., Rudolph, B.-L., Patterson, L., Chapman, P.M. In situ experimental assessment of lake whitefish development following a freshwater oil spill (2007) Environmental Science and Technology, 41 (20), pp. 6983-6989.
- (7) Bergstein, P. E., and Vestal, J. R., "Crude Oil Biodegradation in Arctic Tundra Ponds," *Arctic*, Vol 31, No. 3, 1978, pp. 158–169.
- (8) Atlas, R. M., and Busdosh, M., "Microbial Degradation of Petroleum in the Arctic," *Proceedings of the Third International Biodegradation Symposium, Applied Science*, eds. J. M. Sharpley and A. M. Kaplan, 1976, pp. 79–85.
- (9) Breuil, C., and Shindler, D. B., "Effects of Crude Oil on Bacterial and Fungal Populations in Fresh Water Artificial Ponds," (Abstract), Annual Meeting of American Society for Microbiology 77, 1977, p. 269.
- (10) Dutka, B. J., Sherry, J., Scott, B. F., and Kwan, K. K., "Effects of Oil-Dispersant Mixtures on Fresh Water Microbial Populations," *Canadian Research*, Vol 13, No. 5, 1980, pp. 58–62.
- (11) Federle, T. W., Vestal, J. R., Hater, G. R., and Miller, M. C., "Effects of Prudhoe Bay Crude Oil on Primary Production and Zooplankton in Arctic Tundra Thaw Ponds," *Marine Environmental Research*, Vol 2, No. 1, 1979, pp. 3–18.
- (12) Miller, M. C., Alexander, V., and Barsdate, R. J., "The Effects of Oil Spills on Phytoplankton in an Arctic Lake and Ponds," *Arctic*, Vol 31, No. 3, 1978, pp. 192–218.
- (13) Miller, M. C., Hater, G. R., and Vestal, J. R., "Effect of Prudhoe Crude Oil on Carbon Assimilation by Planktonic Algae in an Arctic Pond," *Environmental Chemistry and Cycling Processes: Proceedings of a Symposium*, April 28 to May 1,1976, Augusta, GA.
- (14) Scott, B. F., and Shindler, D. B., "Impact of Crude Oil on Planktonic Freshwater Ecosystems," Proceedings of the Conference on Assessment of Ecological Impacts of Oil Spills, American Institute of Biological Sciences, Arlington, VA, 1978, pp. 712–734.
- (15) Shindler, D. B., Scott, B. F., and Carlisle, D. B., "Effect of Crude Oil on Populations of Bacteria and Algae in Artificial Ponds Subject to Winter Weather and Ice Formation," *Verhlug International Verein Limnologie*, Vol 19, No. 5, 1975, pp. 2138–2144.
- (16) Stanley, D. W., "Productivity of Epipelic Algae in Tundra Ponds and a Lake near Barrow, Alaska," *Ecology*, Vol 57, No. 5, 1976, pp. 1015–1024.

- (17) Bhattacharyya, S., Klerks, P. L., Nyman, J. A., "Toxicity to Freshwater Organisms from Oils and Oil Spill Chemical Treatments in Laboratory Microcosms," *Environmental Pollution*, Vol 122, No. 2, 1 April 2003, pp. 205–215.
- (18) Klerks, P. L., Nyman, J. A., Bhattacharyya, S., "Relationship Between Hydrocarbon Measurements and Toxicity to a Chironomid, Fish Larva and Daphnid for Oils and Oil Spill Chemical Treatments in Laboratory Freshwater Marsh Microcosms," *Environmental Pollution*, Vol 129, No. 3, June 2004, pp. 345–353.
- (19) Mozley, S. C., and Butler, M. G., "Effects of Crude Oil on Aquatic Insects of Tundra Ponds," Arctic, Vol 31, No. 3, 1978, pp. 229–240.
- (20) Blunden, D., Dale, M., Goudey, S., and Hoddinot, J., "The Effects of Oil Spill Chemicals on Shoreline Plants of Northern Freshwater Ecosystems," *Proceedings of the International Conference on Oil* and Freshwater: Chemistry, Biology Technology, ed., J. H. Vandermeulen, Pergamon, pp. 403–409.
- (21) Barsdate, R. J., "Ecologic Changes in an Arctic Tundra Pond Following Exposure to Crude Oil," (Abstract only), Proceedings of the Symposium on the Impact of Oil Resource Development on Northern Plant Communities, 17 August 1972, Institute of Arctic Biology, University of Alaska, 1973, p. 52.
- (22) Barsdate, R. J., Alexander, V., and Benoit, R. E., "Natural Oil Seeps at Cape Simpson, Alaska: Aquatic Effects," Proceedings of the Symposium on Impact of Oil Resource Development on Northern Plant Communities, 17 August 1972, Institute of Arctic Biology, University of Alaska, 1973, pp. 91–95.
- (23) Goudey, J. S., Dale, M., and Hoddinot, J., "The Effects of Oil Spill Chemicals on Transpiration, CO₂ Exchange and Cuticular Structure in Salix Interior," *Canadian Journal of Botany*, Vol 63, No. 12, 1985, pp. 2340–2344.
- (24) Goudey, J. S., Dale, M., and Hoddinot, J., "Effects of Oil Spill Chemicals on CO₂ Assimilation by the Fruticose Lichen Cladina Mitis," Freshwater Oilspill Research Project, (FOSRP) Report, Esso Resources Canada Limited, Calgary, Alberta, Canada, 1986.
- (25) Goudey, J. S., Dale, M., and Hoddinot, J., "Effects of Crude Oil and Chemically Treated Oil on Shoreline Plant Communities of Northern Bog Lakes," FOSRP Report, Esso Resources Canada Limited, Calgary, Alberta, Canada, 1986.
- (26) Hutchinson, T. C., and Freedman, W., "Effects of Experimental Crude Oil Spills on Subarctic Boreal Forest Vegetation near Norman Wells, N.W.T., Canada," *Canadian Journal of Botany*, Vol 56, No. 19, 1978, pp. 2424–2433.
- (27) Helebust, J. A., Hanna, B., Sheath, R. G., Gergis, M., and Hutchinson, T. C., "Experimental Crude Oil Spills on a Small Subarctic Lake in the MacKenzie Valley, N.W.T.: Effects on Phytoplankton, Periphyton, and Attached Aquatic Vegetation," Proceedings of 1975 Oil Spill Conference, San Francisco, 1975, pp. 509–515.
- (28) Walker, D. A., Webber, P. J., Everett, K. R., and Brown, J., "Effects of Crude and Diesel Oil Spills on Plant Communities at Prudhoe Bay, Alaska and the Derivation of Oil Spill Sensitivity Maps," *Arctic*, Vol 31, No. 3, 1978, pp. 242–259.
- (29) Wein, R. W., and Bliss, L. C., "Experimental Crude Oil Spills on Arctic Plant Communities," *Journal of Applied Ecology*, Vol 10, No. 3, 1973, pp. 671–682.



ASTM International takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.

This standard is copyrighted by ASTM International, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (www.astm.org). Permission rights to photocopy the standard may also be secured from the ASTM website (www.astm.org/COPYRIGHT/).