

Cosmetics — Microbiology — Enumeration of yeast and mould

ICS 07.100.99; 71.100.70

National foreword

This British Standard is the UK implementation of EN ISO 16212:2011. It is identical with ISO 16212:2008. It supersedes BS ISO 16212:2008, which is withdrawn.

The UK participation in its preparation was entrusted to Technical Committee CW/217, Cosmetics.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with a British Standard cannot confer immunity from legal obligations.

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 31 October 2008

© BSI 2011

ISBN 978 0 580 72865 5

Amendments/corrigenda issued since publication

Date	Comments
31 July 2011	This corrigendum renumbers BS ISO 16212:2008 as BS EN ISO 16212:2011

EUROPEAN STANDARD

EN ISO 16212

NORME EUROPÉENNE

EUROPÄISCHE NORM

June 2011

ICS 71.100.70

English Version

Cosmetics - Microbiology - Enumeration of yeast and mould (ISO 16212:2008)

Cosmétiques - Microbiologie - Dénombrement des levures
et des moisissures (ISO 16212:2008)

Kosmetik - Mikrobiologie - Zählung von Hefen und
Schimmelpilzen (ISO 16212:2008)

This European Standard was approved by CEN on 12 May 2011.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: Avenue Marnix 17, B-1000 Brussels

Foreword

The text of ISO 16212:2008 has been prepared by Technical Committee ISO/TC 217 “Cosmetics” of the International Organization for Standardization (ISO) and has been taken over as EN ISO 16212:2011 by Technical Committee CEN/TC 392 “Cosmetics” the secretariat of which is held by NEN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by December 2011, and conflicting national standards shall be withdrawn at the latest by December 2011.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and the United Kingdom.

Endorsement notice

The text of ISO 16212:2008 has been approved by CEN as a EN ISO 16212:2011 without any modification.

Contents

Page

Foreword.....	iv
1 Scope	1
2 Normative references	1
3 Terms and definitions.....	1
4 Principles	2
4.1 General.....	2
4.2 Plate count.....	2
4.3 Membrane filtration.....	2
5 Diluents, neutralizers and culture media.....	3
5.1 General.....	3
5.2 Neutralizing diluents and diluents	3
5.3 Diluent for yeast suspension (tryptone sodium chloride solution).....	4
5.4 Culture media	4
6 Apparatus and glassware	5
7 Strain of microorganisms	5
8 Handling of cosmetic products and laboratory samples	6
9 Procedure	6
9.1 General recommendation	6
9.2 Preparation of the initial suspension.....	6
9.3 Counting methods	6
10 Counting of colonies (plate counts and membrane filtration methods)	7
11 Expression of results	8
11.1 Method of calculation for plate count.....	8
11.2 Interpretation	8
12 Neutralization of the antifungal properties of the product.....	10
12.1 General.....	10
12.2 Preparation of inoculum	10
12.3 Validation of counting methods	11
13 Test report	12
Annex A (informative) Other neutralizing diluents	13
Annex B (informative) Other diluents	15
Annex C (informative) Other culture media	16
Annex D (informative) Neutralizers of antifungal activity of preservatives and rinsing liquids	18
Bibliography	19

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 16212 was prepared by Technical Committee ISO/TC 217, *Comestics*.

Cosmetics — Microbiology — Enumeration of yeast and mould

1 Scope

This International Standard gives general guidelines for enumeration of yeast and mould present in cosmetics by counting the colonies on selective agar medium after aerobic incubation.

In order to ensure product quality and safety for consumers, it is advisable to perform an appropriate microbiological risk analysis so as to determine the types of cosmetic products to which this International Standard is applicable. Products considered to present a low microbiological risk include those with low water activity, hydro-alcoholic products, products with extreme pH values, etc.

Because of the large variety of cosmetic products within this field of application, this method might not be suited to some products in every detail (e.g. certain water-immiscible products). Other methods (e.g. automated) can be used for the test presented here provided that their equivalence has been demonstrated or the method has been otherwise validated.

Yeast enumerated can be identified using suitable identification tests, for example tests described in the standards listed in the Bibliography. Mould enumerated can be identified by other appropriate methods, if necessary.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 21148, *Cosmetics — Microbiology — General instructions for microbiological examination*

EN 12353, *Chemical disinfectants and antiseptics — Preservation of test organisms used for the determination of bactericidal, mycobactericidal, sporicidal and fungicidal activity*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

yeast

single-cell fungus, which multiplies mainly vegetatively by budding, able to grow under the test conditions specified in this International Standard

3.2

mould

mycelium forming microfungus, including spores and conidia, able to grow under the test conditions specified in this International Standard

3.3

product

portion of an identified cosmetic product received in the laboratory for testing

3.4

sample

portion of the product (at least 1 g or 1 ml) which is used in the test to prepare the initial suspension

3.5

initial suspension

suspension (or solution) of a sample in a defined volume of an appropriate liquid (diluent, neutralizer, broth or combination thereof)

3.6

sample dilution(s)

dilution(s) of the initial suspension

4 Principles

4.1 General

This method involves enumeration of colonies on a selective agar medium. The possible inhibition of fungal growth by the sample shall be neutralized to allow the detection of viable microorganism (see Reference [2]). In all cases and whatever the methodology, the neutralization of the antifungal properties of the product shall be checked and validated (see References [3], [5], [6]).

4.2 Plate count

Plate count consists of the following steps.

- a) Preparation of poured plates or spread plates, using a specified culture medium, and inoculation of the plates using a defined quantity of the initial suspension or dilution of the product.
- b) Aerobic incubation of the plates at $25\text{ °C} \pm 2,5\text{ °C}$ for 3 d to 5 d.
- c) Counting of the number of colony-forming units (CFU) and calculation of the amount of yeast and mould per millilitre or per gram of product.

NOTE An alternative condition for incubation is $22,5\text{ °C} \pm 2,5\text{ °C}$ for 5 d to 7 d using the culture medium without antibiotic.

4.3 Membrane filtration

Membrane filtration consists of the following steps.

- a) Transfer of a suitable amount of the sample, prepared by a valid method, in the filtration apparatus, wetted with a small volume of an appropriate sterile diluent. Immediate filtration and washing according to the validated procedure. Transfer of the membrane filter on to the surface of the specified agar medium as specified in ISO 21148.
- b) Aerobic incubation of the membranes at $25\text{ °C} \pm 2,5\text{ °C}$ for 3 d to 5 d.
- c) Counting of the number of colony-forming units (CFU) and calculation of the amount of yeast and mould per millilitre or per gram of product.

NOTE An alternative condition for incubation is $22,5\text{ °C} \pm 2,5\text{ °C}$ for 5 d to 7 d using the culture medium without antibiotic.

5 Diluents, neutralizers and culture media

5.1 General

General specifications are given in ISO 21148. When water is mentioned in a formula, use distilled water or purified water as specified in ISO 21148.

The following diluents, neutralizers and culture media are suitable for enumeration of yeasts and moulds. Other diluents, neutralizers and culture media may be used if they have been demonstrated to be suitable for use.

5.2 Neutralizing diluents and diluents

5.2.1 General

The diluent is used to disperse the sample. It may contain neutralizers if the sample to be tested has antifungal properties. The efficacy of the neutralization shall be demonstrated before the determination of the count (see Clause 12). Information relative to suitable neutralizers is given in Annex D.

5.2.2 Neutralizing diluent

5.2.2.1 Fluid casein digest-soy lecithin-polysorbate 20 medium (SCDLP 20 broth)

5.2.2.1.1 Composition

— pancreatic digest of casein	20,0 g
— soy lecithin	5,0 g
— polysorbate 20	40 ml
— water	960 ml

5.2.2.1.2 Preparation

Dissolve the polysorbate 20 in 960 ml of water by mixing while heating in a water bath at $49\text{ °C} \pm 2\text{ °C}$. Add pancreatic digest of casein and soy lecithin. Heat for about 30 min to effect solution. Mix and dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min. After sterilization, the pH shall be equivalent to $7,3 \pm 0,2$ when measured at room temperature.

5.2.2.2 Other neutralizing diluents

Other neutralizing diluents may be used as appropriate (see Annex A and Annex D).

5.2.3 Diluent

5.2.3.1 Fluid A

5.2.3.1.1 Composition

— peptic digest of animal tissue	1,0 g
— water	1 000 ml

5.2.3.1.2 Preparation

Dissolve 1 g of peptone in water to make 1 l. Heat with frequent agitation. Dispense into suitable containers. Sterilize in the autoclave at 121 °C for 15 min. After sterilization the pH shall be equivalent to $7,1 \pm 0,2$ when measured at room temperature.

5.2.3.2 Other diluents

Other diluents may be used as appropriate (see Annex B).

5.3 Diluent for yeast suspension (tryptone sodium chloride solution)

5.3.1 Composition

— tryptone, pancreatic digest of casein	1,00 g
— sodium chloride	8,50 g
— water	1 000 ml

5.3.2 Preparation

Dissolve the components in the water by mixing while heating. Dispense into suitable containers. Sterilize in the autoclave at 121 °C for 15 min. After sterilization the pH shall be equivalent to $7,0 \pm 0,2$ when measured at room temperature.

5.4 Culture media

5.4.1 General

Culture media may be prepared as follows, or from dehydrated culture media according to the manufacturer's instructions. Ready-to-use media may be used when their composition and/or growth yields are comparable to those of the formulae given herein.

5.4.2 Sabouraud dextrose chloramphenicol agar medium (SDCA)

5.4.2.1 Composition

— dextrose	40,0 g
— peptic digest of animal tissue	5,0 g
— pancreatic digest of casein	5,0 g
— chloramphenicol	0,050 g
— agar	15,0 g
— water	1 000 ml

5.4.2.2 Preparation

Dissolve the components (including the chloramphenicol) or the dehydrated complete medium in the water by mixing while heating. Dispense the medium into suitable containers. Sterilize in an autoclave at 121 °C for 15 min. After sterilization the pH shall be equivalent to $5,6 \pm 0,2$ when measured at room temperature.

NOTE For known and non-contaminated products (with bacteria), the media are used without chloramphenicol.

5.4.3 Other media

Other media may be used as appropriate (see Annex C).

5.4.4 Agar medium for cultivation of reference strain: Sabouraud dextrose agar medium (SDA)

5.4.4.1 Composition

— dextrose	40,0 g
— peptic digest of animal tissue	5,0 g
— pancreatic digest of casein	5,0 g
— agar	15,0 g
— water	1 000 ml

5.4.4.2 Preparation

Dissolve the components or the dehydrated complete medium in the water by mixing while heating. Dispense the medium into suitable containers. Sterilize in an autoclave at 121 °C for 15 min. After sterilization the pH shall be equivalent to $5,6 \pm 0,2$ when measured at room temperature.

6 Apparatus and glassware

The laboratory equipment, apparatus and glassware are described in ISO 21148.

7 Strain of microorganisms

For testing the efficacy of neutralizers, one yeast reference strain is used.

Candida albicans ATCC¹⁾ 10231 or equivalent strain (IP²⁾ 48.72 or NCPF³⁾ 3179 or NBRC⁴⁾ 1594 or KCTC⁵⁾ 17205 or TISTR⁶⁾ 5779) or other equivalent national collection strain.

The selected yeast strain being considered more susceptible to antifungicidal activity than moulds may be accepted as representative of fungi (yeast and mould) for the validation of the methodology. However, in case of specific needs, the test for the efficacy of neutralizers may be performed with an additional mould reference strain, using a suitable protocol for the preparation of a calibrated inoculum (e.g. see EN 13624:2003^[11], 5.4.1.4).

The culture should be reconstituted according to the procedures provided by the supplier of reference strain.

The strain may be kept in the laboratory in accordance with EN 12353.

-
- 1) ATCC = American Type Culture Collection.
 - 2) IP = Institut Pasteur.
 - 3) NCPF = National Collection of Pathogenic Fungi.
 - 4) NBRC = National Biological Resource Center.
 - 5) KCTC = Korean Collection for Type Culture.
 - 6) TISTR = Thailand Institute of Scientific and Technological Research.

8 Handling of cosmetic products and laboratory samples

If necessary, store products to be tested at room temperature.

Do not incubate, refrigerate or freeze products (3.3) and samples (3.4) before or after analysis.

Sampling of cosmetic products to be analysed should be carried out as described in ISO 21148. Analyse samples as described in ISO 21148 and according to the procedure described in Clause 9.

9 Procedure

9.1 General recommendation

Use sterile material, equipment and aseptic techniques to prepare the sample, initial suspension and dilutions. In the case of the preparation of the initial suspension, the time that elapses between the end of the preparation and the moment the inoculum comes into contact with the culture medium shall not exceed 45 min, unless specifically mentioned in the established protocols or documents.

9.2 Preparation of the initial suspension

9.2.1 General

The initial suspension is prepared from a sample of at least 1 g or 1 ml of the well-mixed product under test.

Note *S*, the exact mass or volume of the sample.

The initial suspension is usually a 1:10 dilution. Larger volumes of diluent may be required if high levels of contamination are expected and/or if antifungicidal properties are still present in the 1:10 dilution.

9.2.2 Water-miscible products

Transfer the sample, *S*, of product to an appropriate volume (e.g. 9 ml) of neutralizing diluent (see 5.2.2) or diluent (see 5.2.3).

Note *d*, the dilution factor.

9.2.3 Water-immiscible products

Transfer the sample, *S*, of product to a suitable container containing a suitable quantity of solubilizing agent (e.g. polysorbate 80 solution). Disperse the sample within the solubilizing agent and add an appropriate volume (e.g. 9 ml) of neutralizing diluent (see 5.2.2) or diluent (see 5.2.3).

Note *d*, the dilution factor.

9.3 Counting methods

9.3.1 Dilutions for counting methods

Usually, the initial suspension is the first counted dilution. If needed, additional serial dilutions (e.g. 1:10 dilution) may be performed from the initial suspension using the same diluent (according to the expected level of contamination of the product).

Generally, counting is performed using at least two Petri dishes, but it is possible to use only one Petri dish in case of routine testing, or if counts are performed on successive dilutions of the same sample or according to previous results.

9.3.2 Plate-count methods

9.3.2.1 Pour-plate method

In Petri dishes of 85 mm to 100 mm in diameter, add 1 ml of the initial suspension and/or sample dilution prepared as validated (see Clause 12) and pour 15 ml to 20 ml of the melted agar medium (see 5.4.2) kept in a water bath at no more than 48 °C. If larger Petri dishes are used, the amount of agar medium is increased accordingly.

Mix the initial suspension and/or sample dilution with the medium, carefully rotating or tilting the plates sufficiently to disperse them. Allow the mixture in the Petri dishes to solidify on a horizontal surface at room temperature.

9.3.2.2 Surface spread method

In Petri dishes of 85 mm to 100 mm in diameter, put 15 ml to 20 ml of the melted agar medium (see 5.4.2) kept in a water bath at no more than 48 °C. If larger Petri dishes are used, the volume of the agar is increased accordingly.

Allow plates to cool and solidify, for example in a microbiological cabinet or in an incubator. Spread over the surface of the medium a measured volume of not less than 0,1 ml of the initial suspension and/or sample dilution prepared as validated (see Clause 12).

9.3.2.3 Membrane filtration method

Use a membrane having a nominal pore size $\leq 0,45 \mu\text{m}$.

Transfer a suitable amount of the initial suspension or of the sample dilution prepared as validated (preferably representing at least 1 g or 1 ml of the product) on to the membrane.

Filter immediately and wash the membrane (follow the procedure developed during the validation, see Clause 12).

Transfer the membrane on to the surface of the agar medium (see 5.4.2).

9.3.2.4 Incubation

Unless otherwise stated, invert the inoculated dishes and place them in the incubator set at $25 \text{ °C} \pm 2,5 \text{ °C}$ for 3 d to 5 d or use the alternative condition (see Notes in 4.2 and 4.3). After incubation, the dishes shall, if possible, be examined immediately. Alternatively, they may be stored, unless otherwise specified, for up to a maximum of 24 h in the refrigerator at $5 \text{ °C} \pm 3 \text{ °C}$.

NOTE 1 In certain cases, where there is a potential for confusing particles from the product with counted colonies, it can be useful to prepare duplicate dishes containing the same sample dilutions and agar medium which are stored in the refrigerator for comparison with incubated dishes.

NOTE 2 An intermediate check can be performed where both yeast and mould are suspected.

10 Counting of colonies (plate counts and membrane filtration methods)

After incubation, count the colonies:

- in Petri dishes containing 15 colonies to 150 colonies; if less than 15 colonies are counted, see 11.2.3;
- on membranes containing 15 colonies to 150 colonies; if less than 15 colonies are counted, see 11.2.3.

11 Expression of results

11.1 Method of calculation for plate count

Calculate the number, N , of microorganisms present in the sample, S , using:

- m , the arithmetic mean of the counts obtained from the duplicates (1),
- c , the number of colonies counted on a single plate (2) or
- wm , the weighted mean of the counts obtained from two successive dilutions (3),

according to the following formulae:

$$N = ml(V \times d) \quad (1)$$

$$N = cl(V \times d) \quad (2)$$

$$N = wml(V \times d) \quad (3)$$

where:

- m is the arithmetic mean of the counts obtained from the duplicates;
- V is the volume of inoculum applied to each dish, in millilitres;
- d is the dilution factor corresponding to the dilution made for the preparation of the initial suspension (see 9.2) or for the first counted dilution;
- c is the number of colonies counted on a single plate.

The weighted mean of the colonies counted from two successive dilutions, \bar{x}_c , is given by:

$$\bar{x}_c = \frac{\Sigma c}{(n_1 + 0,1n_2)}$$

where

- Σc is the sum of colonies counted on all the dishes retained from two successive dilutions;
- n_1 is the number of dishes counted for the initial suspension (or for the first counted dilution);
- n_2 is the number of dishes counted for the 1:10 dilution of the initial suspension (or for the second counted dilution).

Round off the result calculated to two significant figures. For this, if the last figure is below 5, the preceding figure is not modified; if the last figure is 5 or more, the preceding figure is increased by one unit. Proceed stepwise until two significant figures are obtained. Note the number, N , obtained.

11.2 Interpretation

11.2.1 The inherent variability of plate count should be taken into account. Two results should only be considered different if the difference exceeds 50 % or, when expressed logarithmically, the difference exceeds 0,3 log.

For a count to be precise, only plates or membranes with more than 15 colonies and less than 150 colonies shall be taken into account. Check that the counts are obtained from dilutions validated according to the chosen method (see Clause 12).

11.2.2 Where the number of CFU is more than 15 and less than 150 on plates or membranes, express the result as follows:

- if S is at least 1 g or 1 ml, and V is at least 1 ml, the number of yeast and mould per millilitre or per gram of the sample = N/S ;
- if S is less than 1 g or 1 ml, and/or V is less than 1 ml, the number of yeast and mould in the sample (note the tested quantity of sample taking into account S and V) = N ;

where S is the mass or the volume of the sample (see 9.2).

Express the result as a number between 1,0 and 9,9 multiplied by the appropriate power of 10 (see examples).

11.2.3 Where the number of CFU is less than 15 on plates or membranes, express the result as follows:

- if S is at least 1 g or 1 ml, and V is at least 1 ml, the estimated number of yeast and mould per millilitre or per gram of the sample = N/S ;
- if S is less than 1 g or 1 ml, and/or V is less than 1 ml, the estimated number of yeast and mould in the sample = N ;

where S is the mass or the volume of the sample (see 9.2).

Express the result as a number between 1,0 and 9,9 multiplied by the appropriate power of 10 (see examples).

11.2.4 Where no colony is observed, the result is reported as follows:

- there is less than $1/d \times V \times S$ of yeast and mold per gram or millilitre of the product (S is at least 1 g or 1 ml);
- there is less than $1/d \times V$ of yeast and mould in the sample S (note the tested quantity of sample taking into account S and V) (S is less than 1 g or 1 ml);

where

d is the dilution factor of the initial suspension (see 9.2);

V is 1 (for counting with the pour-plate method and for membrane filtration) or 0,1 (for the spread plate method) (see example).

EXAMPLE 1 Two dishes for one dilution

$S = 1$ g; $V = 1$; counts obtained for the dilution 10^{-1} : 38 and 42

(1) $N = m/(V \times d) = 40/(1 \times 10^{-1}) = 40/0,1 = 400$ or 4×10^2 yeast and mould per millilitre or per gram of the sample.

EXAMPLE 2 One dish for one dilution

$S = 1$ g; $V = 1$; count obtained for the dilution 10^{-1} : 60

(2) $N = c/(V \times d) = 60/(1 \times 10^{-1}) = 60/0,1 = 600$ or 6×10^2 yeast and mould per millilitre or per gram of the sample.

EXAMPLE 3 Two dishes for two dilutions

$S = 1$ g; $V = 1$; counts obtained for the dilution 10^{-2} : 235 and 282; for the dilution 10^{-3} : 31 and 39

$$(3) N = wml/(V \times d) = 235 + 282 + 31 + 39/1(2 + 0,1 \times 2) \times 10^{-2} = 587/0,022 = 26\ 682.$$

Rounding the result as specified above gives 27 000 or $2,7 \times 10^4$ yeast and mould per millilitre or per gram of the sample.

EXAMPLE 4 Two membrane filters for one dilution

$S = 1$ g; $V = 1$; counts obtained for the dilution 10^{-1} : 18 and 22

$$(1) N = ml/(V \times d) = 20/(1 \times 10^{-1}) = 20/0,1 = 200 \text{ or } 2 \times 10^2 \text{ yeast and mould per millilitre or per gram of the sample.}$$

EXAMPLE 5 One membrane filter for one dilution

$S = 1$ g; $V = 1$; count obtained for the dilution 10^{-1} : 65

$$(2) N = c/(V \times d) = 65/(1 \times 10^{-1}) = 65/0,1 = 650 \text{ or } 6,5 \times 10^2 \text{ aerobic yeast and mould per millilitre or per gram of the sample.}$$

EXAMPLE 6 Two membrane filters for two dilutions

$S = 1$ g; $V = 1$; counts obtained for the dilution 10^{-1} : 121 and 105; for the dilution 10^{-2} : 15 and 25

$$(3) N = wml/(V \times d) = 121 + 105 + 15 + 25/1(2 + 0,1 \times 2) \times 10^{-1} = 266/0,22 = 1\ 209.$$

Rounding the result as specified above gives 1 200 or $1,2 \times 10^3$ yeast and mould per millilitre or per gram of the sample.

EXAMPLE 7 Two dishes for one dilution

$S = 1$ g; $V = 1$; estimated counts obtained for the dilution 10^{-1} : 28 and 22

$$(1) N = ml/(V \times d) = 25/(1 \times 10^{-1}) = 25/0,1 = 250 \text{ or } 2,5 \times 10^2 \text{ yeast and mould per millilitre or per gram of the sample.}$$

The estimated number is 250 or $2,5 \times 10^2$ yeast and mould per millilitre or per gram of the sample.

EXAMPLE 8

$S = 1$ g; $V = 1$; estimated counts obtained for the dilution 10^{-1} : 0 and 0

$$(1) N = \leq 1/d \times V \times S, \leq (1/0,1) \times 1 \times 1 \leq 10 \text{ yeast and mould per millilitre or per gram of the sample.}$$

The estimated number is less than 10 yeast and mould per millilitre or per gram of the sample.

12 Neutralization of the antifungicidal properties of the product

12.1 General

The different tests described below demonstrate that the microorganisms can grow under the conditions of analysis.

12.2 Preparation of inoculum

Prior to the test, inoculate the surface of the non-selective agar medium, Sabouraud dextrose agar (SDA), with *Candida albicans*. Incubate the plate at $32,5 \text{ }^\circ\text{C} \pm 2,5 \text{ }^\circ\text{C}$ for 18 h to 24 h.

To harvest the culture, use a sterile loop, streak the surface of the culture and re-suspend in the diluent (see 5.2) to obtain a calibrated suspension of about 1×10^6 CFU/ml (e.g. determined using spectrophotometer); see ISO 21148.

Use this calibrated suspension and its dilutions within 2 h.

12.3 Validation of counting methods

12.3.1 Principle

Mix the neutralized sample (initial suspension or sample dilution according to the antifungal activity or the low solubility of the product) with a dilution of the reference strain. Plate on a Petri dish or filter on a membrane. After incubation, check the nature of the colonies and compare the count with a control (without the sample).

If the count is less than 50 % (0,3 log) of the control, modify the procedure (diluent, neutralization agents or a combination of both, see Annex D). Failure of the inoculum to grow invalidates the test unless possible contamination of the product with this microorganism is unlikely.

12.3.2 Validation of the pour-plate method

Mix 9 ml of the initial suspension and/or the sample dilution(s) in neutralizing diluent (or other, see 5.2) with 1 ml of a suspension of microorganisms containing 1 000 CFU/ml to 3 000 CFU/ml. Transfer 1 ml in a Petri dish (preferably in duplicate) and pour 15 ml to 20 ml of the melted agar medium (see 5.4.2) kept in a water bath at no more than 48 °C. In parallel, prepare and plate a control using the same diluent and the same suspension of microorganisms, but without the sample.

After incubation for 3 d to 5 d at $25 \text{ °C} \pm 2,5 \text{ °C}$ or using the alternative condition (see Notes in 4.2 and 4.3), count the colonies on the plates and compare the counts obtained for the test and for the control. The diluent and the counting method are validated at the 1:10 dilution (when 1 ml of the initial suspension is used) if the validation count is at least 50 % (0,3 log) of the control count.

12.3.3 Validation of the surface spread method

Mix 9 ml of the initial suspension in neutralizing diluent (or other, see 5.2) with 1 ml of a suspension of microorganisms containing 10 000 CFU/ml to 30 000 CFU/ml (or less if 0,5 ml or 1 ml are spread). Spread at least 0,1 ml on a solidified agar plate (see 5.4.2), preferably in duplicate. In parallel, prepare and plate a control using the same diluent and the same suspension of microorganisms, but without the sample.

After incubation for 3 d to 5 d at $25 \text{ °C} \pm 2,5 \text{ °C}$ or using the alternative condition (see Notes in 4.2 and 4.3), count the colonies on the plates and compare the counts obtained for the test and for the control. The diluent and the counting method are validated at the 1:10 dilution (when 1 ml of the initial suspension is used) if the validation count is at least 50 % (0,3 log) of the control count.

12.3.4 Validation of the membrane filtration method

Mix to the volume of initial suspension or of the sample dilution used in the test (see 9.3.2.3) a suitable amount of a calibrated suspension of microorganisms corresponding to approximately 100 CFU.

Filter immediately the entire volume and wash the membrane using defined volumes of water (see 5.1), diluent (see 5.2.3) or neutralizing diluent (see 5.2.2). Transfer the membrane on to the surface of a suitable agar medium (see 5.4.2).

In parallel, prepare a control under the same conditions as above, but without the product. Filter and wash the control under the same conditions.

After incubation for 3 d to 5 d at $25 \text{ °C} \pm 2,5 \text{ °C}$ or using the alternative condition (see Notes in 4.2 and 4.3), count the colonies on the membranes and compare the counts obtained for the test and for the control. The membrane filtration method and the diluent are validated if the validation count is at least 50 % (0,3 log) of the control count.

13 Test report

The test report shall contain the following:

- a) all information necessary for the complete identification of the product;
- b) the method used;
- c) the results obtained;
- d) all operating details for the preparation of the initial suspension;
- e) the description of the method with the neutralizers and media used;
- f) the validation of the method, even if the test has been performed separately;
- g) any point not specified in this International Standard, or regarded as optional, together with details of any incidents which may have influenced the results.

Annex A (informative)

Other neutralizing diluents

A.1 General

Any neutralizing diluent may be used to prepare the initial suspension if it has been checked and validated. The following neutralizing diluents are examples of suitable formulae. General information on neutralization is given in Annex D.

A.2 Eugon LT100 liquid broth

A.2.1 General

This medium contains ingredients that neutralize inhibitory substances present in the sample (lecithin and polysorbate 80); and dispersing agent (octoxynol 9).

A.2.2 Composition

— pancreatic digest of casein	15,0 g
— papaic digest of soybean meal	5,0 g
— L-cystine	0,7 g
— sodium chloride	4,0 g
— sodium sulfite	0,2 g
— glucose	5,5 g
— egg lecithin	1,0 g
— polysorbate 80	5,0 g
— octoxynol 9	1,0 g
— water	1 000 ml

A.2.3 Preparation

Dissolve successively in hot water: polysorbate 80, octoxynol 9 and egg lecithin until their complete dilution. Dissolve the other components by mixing while heating. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min. After sterilization, the pH shall be equivalent to $7,0 \pm 0,2$ when measured at room temperature.

A.3 Lecithin polysorbate (LP) diluent

A.3.1 Composition

— polypeptone	1,0 g
— egg lecithin	0,7 g
— polysorbate 80	20,0 g
— water	980 ml

A.3.2 Preparation

Mix and dissolve the ingredients by heating. Cool to 25 °C before dispensing the solution into suitable containers. Sterilize in the autoclave at 121 °C for 15 min. After sterilization the pH shall be equivalent to $7,2 \pm 0,2$ when measured at room temperature.

Annex B (informative)

Other diluents

B.1 General

Any diluent may be used to prepare the initial suspension if it has been checked and validated. The following diluent is an example of a suitable formula.

B.2 Buffered peptone solution (pH 7)

B.2.1 Composition

— meat peptone	1,0 g
— sodium chloride	4,3 g
— monopotassium phosphate	3,6 g
— disodium phosphate dihydrate	7,2 g
— water	1 000 ml

B.2.2 Preparation

Dissolve the ingredients in hot water. Mix and cool to 25 °C before dispensing the solution into suitable containers. Sterilize in the autoclave at 121 °C for 15 min. After sterilization the pH shall be equivalent to $7,1 \pm 0,2$ when measured at room temperature.

B.3 Phosphate buffer (pH 7,2)

B.3.1 Composition

— potassium dihydrogen phosphate	34 g
— water	500 ml

B.3.2 Preparation

Dissolve the ingredients in water in a 1 000 ml volumetric flask. Adjust the pH value to $7,2 \pm 0,1$ with about 175 ml of sodium hydroxide (4,3 g/100 ml). Then water is added to a final volume of 1 000 ml. The final concentration of the solution is 0,05 mol/l. It is the stock solution. Store it under refrigeration.

Before use, dilute the stock solution with water at a ratio of 1:800 and sterilize it in the autoclave at 121 °C for 15 min.

Annex C (informative)

Other culture media

C.1 General

Any culture medium may be used if it has been checked and validated. The following media are examples of suitable formulae.

C.2 Agar for counting

C.2.1 Potato dextrose agar medium with antibiotics

C.2.1.1 Composition

— potato extract	4,0 g
— glucose	20,0 g
— agar	15,0 g
— chloramphenicol	0,05 g
— water	1 000 ml

C.2.1.2 Preparation

Mix all the components and dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min. After sterilization and cooling, the pH shall be equivalent to $5,6 \pm 0,2$ when measured at room temperature.

Alternatively, chloramphenicol may be replaced by use of 0,10 g of benzylpenicillin potassium and 0,10 g of tetracycline per litre of medium, added as sterile solution, just prior to use.

C.2.2 Glucose-peptone (GP) agar medium with antibiotics

C.2.2.1 Composition

— glucose	20,0 g
— yeast extract	2,0 g
— magnesium sulfate	0,5 g
— peptone	5,0 g
— monobasic potassium phosphate	1,0 g
— agar	15,0 g

— chloramphenicol	0,05 g
— water	1 000 ml

C.2.2.2 Preparation

Mix all the components and dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min. After sterilization and cooling, the pH shall be equivalent to $5,7 \pm 0,1$ when measured at room temperature.

Alternatively, chloramphenicol may be replaced by use of 0,10 g of benzylpenicillin potassium and 0,10 g of tetracycline per litre of medium, added as sterile solution, just prior to use.

C.3 Malt extract medium

C.3.1 Composition

— malt extract	30,0 g
— soya peptone, papaic digest of soybean meal	3,0 g
— agar	15,0 g
— chloramphenicol	0,05 g
— water	1 000 ml

C.3.2 Preparation

Dissolve the components (including the chloramphenicol) or the dehydrated complete medium in the water by heating. Dispense the medium into suitable containers. Sterilize in an autoclave at 115 °C for 10 min. After sterilization the pH shall be equivalent to $5,6 \pm 0,2$ when measured at room temperature.

Annex D (informative)

Neutralizers of antifungicidal activity of preservatives and rinsing liquids

Preservative	Chemical compounds able to neutralize preservative's antifungicidal activity	Examples of suitable neutralizers and of rinsing liquids (for membrane filtration methods)
Phenolic compounds: parabens, phenoxyethanol, phenylethanol, anilides	Lecithin Polysorbate 80 Ethylene oxide condensate of fatty alcohol Non-ionic surfactants	Polysorbate 80, 30 g/l + lecithin, 3 g/l. Ethylene oxide condensate of fatty alcohol, 7 g/l + lecithin, 20 g/l + polysorbate 80, 4 g/l. D/E neutralizing broth Rinsing liquid: distilled water; tryptone, 1 g/l + NaCl, 9 g/l; polysorbate 80, 5 g/l.
Quaternary ammonium compounds Cationic surfactants	Lecithin, saponin polysorbate 80, sodium dodecyl sulphate Ethylene oxide condensate of fatty alcohol	Polysorbate 80, 30 g/l + sodium dodecyl sulphate, 4 g/l + lecithin, 3 g/l. Polysorbate 80, 30 g/l + saponin, 30 g/l + lecithin, 3 g/l. D/E neutralizing broth Rinsing liquid: distilled water; tryptone, 1 g/l + NaCl, 9 g/l; polysorbate 80, 5 g/l.
Aldehydes Formaldehyde-release agents	Glycine, histidine	Lecithin, 3 g/l + polysorbate 80, 30 g/l + L-histidine, 1 g/l. Polysorbate 80, 30 g/l + saponin, 30 g/l + L-histidine, 1 g/l + L-cysteine, 1 g/l. D/E neutralizing broth Rinsing liquid: polysorbate 80, 3 g/l + L-histidine 0,5 g/l.
Oxidizing compounds	Sodium thiosulphate	Sodium thiosulphate, 5 g/l. Rinsing liquid: sodium thiosulphate, 3 g/l.
Isothiazolinones, imidazoles	Lecithin, saponin, amines, sulfates, mercaptans, sodium bisulfite, sodium thioglycollate	Polysorbate 80, 30 g/l + saponin, 30 g/l + lecithin, 3 g/l. Rinsing liquid: tryptone, 1 g/l + NaCl, 9 g/l; polysorbate 80, 5 g/l.
Biguanides	Lecithin, saponin, polysorbate 80	Polysorbate 80, 30 g/l + saponin, 30 g/l + lecithin, 3 g/l. Rinsing liquid: tryptone, 1 g/l + NaCl, 9 g/l; polysorbate 80, 5 g/l.
Metallic salts (Cu, Zn, Hg) Organo-mercuric compounds	Sodium bisulphate, L-cysteine Sulfhydryl compounds, thioglycollic acid	Sodium thioglycollate, 0,5 g/l or 5 g/l. L-cysteine, 0,8 g/l or 1,5 g/l. D/E neutralizing broth Rinsing liquid: sodium thioglycollate, 0,5 g/l.
NOTE According to the pH of the cosmetic product, the pH of the neutralizer can be adjusted at a suitable value.		

Bibliography

- [1] COLIPA, *Guidelines on Microbial Quality Management*, European Cosmetic, Toiletry and Perfumery Association (COLIPA), 1997
- [2] CTFA, *Microbiology Guidelines*, Toiletry and Fragrance Association, ISBN 1-882621-32-8, 2001
- [3] EP, *Microbiological Examination of Non-Sterile Products*, 4th edition, European Pharmacopoeia, 2002
- [4] FDA, *Bacteriological Analytical Manual*, 8th edition, U.S. Food and Drug Administration, 1995, <http://www.cfsan.fda.gov/~ebam/bam-23.html>
- [5] JP 14, *General Tests — Microbial Limit Test*, Japanese Pharmacopoeia, 2001
- [6] USP 28, *Microbial Limit Test (61)*, U.S. Pharmacopoeia, 2005
- [7] ATLAS, R.M., *Handbook of Microbiological Media*, CRC Press, ISBN 0-8493-2944-2, 1993
- [8] SINGER, S., *The Use of Preservative Neutralizers in Diluents and Plating Media*, *Cosmetics and Toiletries*, **102**, p. 55, December 1987
- [9] ISO 18415, *Cosmetics — Microbiology — Detection of specified and non-specified microorganisms*
- [10] ISO 18416, *Cosmetics — Microbiology — Detection of *Candida albicans**
- [11] EN 13624:2003, *Chemical disinfectants and antiseptics — Quantitative suspension test for the evaluation of fungicidal activity of chemical disinfectants for instruments used in the medical area — Test method and requirements (phase 2, step 1)*

ICS 71.100.70

Price based on 19 pages

BSI - British Standards Institution

BSI is the independent national body responsible for preparing British Standards. It presents the UK view on standards in Europe and at the international level. It is incorporated by Royal Charter.

Revisions

British Standards are updated by amendment or revision. Users of British Standards should make sure that they possess the latest amendments or editions.

It is the constant aim of BSI to improve the quality of our products and services. We would be grateful if anyone finding an inaccuracy or ambiguity while using this British Standard would inform the Secretary of the technical committee responsible, the identity of which can be found on the inside front cover. Tel: +44 (0)20 8996 9000. Fax: +44 (0)20 8996 7400.

BSI offers members an individual updating service called PLUS which ensures that subscribers automatically receive the latest editions of standards.

Buying standards

Orders for all BSI, international and foreign standards publications should be addressed to Customer Services. Tel: +44 (0)20 8996 9001. Fax: +44 (0)20 8996 7001 Email: orders@bsigroup.com You may also buy directly using a debit/credit card from the BSI Shop on the Website <http://www.bsigroup.com/shop>

In response to orders for international standards, it is BSI policy to supply the BSI implementation of those that have been published as British Standards, unless otherwise requested.

Information on standards

BSI provides a wide range of information on national, European and international standards through its Library and its Technical Help to Exporters Service. Various BSI electronic information services are also available which give details on all its products and services. Contact Information Centre. Tel: +44 (0)20 8996 7111 Fax: +44 (0)20 8996 7048 Email: info@bsigroup.com

Subscribing members of BSI are kept up to date with standards developments and receive substantial discounts on the purchase price of standards. For details of these and other benefits contact Membership Administration. Tel: +44 (0)20 8996 7002 Fax: +44 (0)20 8996 7001 Email: membership@bsigroup.com

Information regarding online access to British Standards via British Standards Online can be found at <http://www.bsigroup.com/BSOL>

Further information about BSI is available on the BSI website at <http://www.bsigroup.com>

Copyright

Copyright subsists in all BSI publications. BSI also holds the copyright, in the UK, of the publications of the international standardization bodies. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI.

This does not preclude the free use, in the course of implementing the standard, of necessary details such as symbols, and size, type or grade designations. If these details are to be used for any other purpose than implementation then the prior written permission of BSI must be obtained.

Details and advice can be obtained from the Copyright and Licensing Manager. Tel: +44 (0)20 8996 7070 Email: copyright@bsigroup.com