

Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — Nucleic acid extraction

ICS 67.050

National foreword

This British Standard is the UK implementation of EN ISO 21571:2005+A1:2013. It is identical to ISO 21571:2005, incorporating amendment 1:2013. It supersedes BS EN ISO 21571:2005, which is withdrawn.

The start and finish of text introduced or altered by amendment is indicated in the text by tags. Tags indicating changes to ISO text carry the number of the ISO amendment. For example, text altered by ISO amendment 1 is indicated by **A1** **A1**.

The UK participation in its preparation was entrusted to Technical Committee AW/275, Food analysis — Horizontal methods.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with a British Standard cannot confer immunity from legal obligations.

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 24 March 2005

© The British Standards Institution 2013. Published by BSI Standards Limited 2013.

Amendments/corrigenda issued since publication

Date	Comments
31 May 2013	Implementation of ISO amendment 1:2013 with CEN endorsement A1:2013

ICS 67.050

English version

Foodstuffs - Methods of analysis for the detection of genetically modified organisms and derived products - Nucleic acid extraction (ISO 21571:2005)

Produits alimentaires - Méthodes d'analyse pour la détection des organismes génétiquement modifiés et des produits dérivés - Extraction des acides nucléiques (ISO 21571:2005)

Lebensmittel - Verfahren zum Nachweis von gentechnisch modifizierten Organismen und ihren Produkten - Nukleinsäureextraktion (ISO 21571:2005)

This European Standard was approved by CEN on 14 February 2005.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the Central Secretariat or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the Central Secretariat has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: rue de Stassart, 36 B-1050 Brussels

Foreword

This document (EN ISO 21571:2005) has been prepared by Technical Committee CEN/TC 275 "Food analysis - Horizontal methods", the secretariat of which is held by DIN, in collaboration with Technical Committee ISO/TC 34 "Agricultural food products".

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by August 2005, and conflicting national standards shall be withdrawn at the latest by August 2005.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.

Foreword to amendment A1

This document (EN ISO 21571:2005/A1:2013) has been prepared by Technical Committee ISO/TC 34 "Food products" in collaboration with Technical Committee CEN/TC 275 "Food analysis - Horizontal methods" the secretariat of which is held by DIN.

This Amendment to the European Standard EN ISO 21571:2005 shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by September 2013, and conflicting national standards shall be withdrawn at the latest by September 2013.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Endorsement notice

The text of ISO 21571:2005/Amd 1:2013 has been approved by CEN as EN ISO 21571:2005/A1:2013 without any modification.

**Foodstuffs — Methods of analysis for
the detection of genetically modified
organisms and derived products —
Nucleic acid extraction**

*Produits alimentaires — Méthodes d'analyse pour la détection des
organismes génétiquement modifiés et des produits dérivés —
Extraction des acides nucléiques*

Contents

Page

Foreword	iv
Introduction	v
1 Scope.....	1
2 Normative references	1
3 Principle	1
3.1 General	1
3.2 DNA extraction	2
3.3 DNA quantitation.....	2
4 General laboratory requirements	2
5 Procedure.....	2
5.1 Preparation of the test portion.....	2
5.2 DNA extraction/purification.....	4
5.3 Quantitation of the extracted DNA	5
5.4 Stability of extracted DNA.....	6
6 Interpretation	6
7 Test report.....	6
Annex A (informative) Methods for DNA extraction.....	7
Annex B (informative) Methods for the quantitation of the extracted DNA.....	43
Bibliography	50

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

ISO 21571 was prepared by the European Committee for Standardization (CEN) Technical Committee CEN/TC 275, *Food analysis — Horizontal methods*, in collaboration with Technical Committee ISO/TC 34, *Food products*, in accordance with the Agreement on technical cooperation between ISO and CEN (Vienna Agreement).

Introduction

The search for genetically modified origin of ingredients is performed by means of the following successive (or simultaneous) steps. After sample collection, nucleic acids are extracted from the test portion. Extracted nucleic acids can be further purified, simultaneously or after the extraction process. Afterwards, they are quantified (if necessary), diluted (if necessary) and subjected to analytical procedures (such as PCR). These steps are detailed in this and the following International Standards:

A1 Text deleted **A1**

ISO 21569, *Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — Qualitative nucleic acid based methods.*

ISO 21570, *Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — Quantitative nucleic acid based methods.*

Further information about definitions and general items involving the steps cited above are collected in:

A1 ISO 24276, *Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — General requirements and definitions.* **A1**

The International Organization for Standardization (ISO) draws attention to the fact that it is claimed that compliance with this document may involve the use of a patent concerning the silica-based extraction method (No. EP 0389063/USP 5,234,809) given in Clause A.4.

ISO takes no position concerning the evidence, validity and scope of this patent right.

The holder of this patent right has assured the ISO that he/she is willing to negotiate licences under reasonable and non-discriminatory terms and conditions with applicants throughout the world. In this respect, the statement of the holder of this patent right is registered with ISO. Information may be obtained from:

Jean Deleforge,
BioMérieux
Chemin de l'Orme,
69280 Marcy-l'Étoile, France.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights other than those identified above. ISO shall not be held responsible for identifying any or all such patent rights.

Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — Nucleic acid extraction

1 Scope

This International Standard provides general requirements and specific methods for DNA extraction/purification and quantitation. These methods are described in Annexes A and B.

This International Standard has been established for food matrices, but could also be applicable to other matrices, such as grains and feed.

It has been designed as an integral part of nucleic-acid-based analytical methods, in particular ISO 21569 on qualitative analytical methods, and ISO 21570 on quantitative analytical methods.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

^{A1} ISO 24276:2006, *Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — General requirements and definitions*. ^{A1}

^{A1} Text deleted ^{A1}

3 Principle

3.1 General

The objective of nucleic acid extraction methods is to provide nucleic acids suitable for subsequent analysis (see ISO 24276).

NOTE The “quality” of DNA depends on the average length of the extracted DNA molecules, the chemical purity and the structural integrity of the DNA sequence and of the double helix (e.g. intra-, inter-strand linking between DNA bases, single-strand gaps, cross-linking with polyols, haemin, etc). Moreover, such alterations are often sequence-specific and consequently not randomly distributed all over the genome (see References [1], [2], [3] and [4]).

Users of this International Standard should note that some methods (e.g. all silica-based methods), might be covered by patent rights.

^{A1} Note deleted ^{A1}

3.2 DNA extraction

The basic principle of DNA extraction consists of releasing the DNA present in the matrix and further, concurrently or subsequently, purifying the DNA from polymerase chain reaction (PCR) inhibitors.

DNA extraction/purification methods are described in Annex A. Method-selection is an experience-based choice of the user, taking into account the scope and examples of matrices as given in each method.

Alternative protocols are suitable provided that the method has been validated on the respective matrix under investigation.

3.3 DNA quantitation

Quantitation of extracted DNA could be useful for subsequent PCR analysis.

It may be performed by either physical (e.g. measure of absorbance at a specific wavelength), chemical-physical (e.g. use of intercalating or binding agents able to emit fluorescence), enzymatic (e.g. bioluminescence detection) methods or by quantitative PCR. The latter method is especially suitable for composite matrices or for samples with a low DNA content or whose DNA is degraded.

There are several methods available to quantify the DNA present in a solution, as described in Annex B. It is for the user to choose the most appropriate one to be applied, depending on the amount and quality of DNA to be quantified and, consequently, on the matrix from which the DNA has been extracted.

Alternative protocols are suitable, provided that the method has been validated on the respective matrix under investigation.

4 General laboratory requirements

Accidental contamination of DNA can originate from dust and spreading aerosols. As a consequence, the organization of the work area in the laboratory is logically based on

- the systematic containment of the methodological steps involved in the production of the results, and
- a “forward flow” principle for sample handling.

The latter ensures that the DNA to be analysed and the amplified DNA generated by PCR remain physically segregated.

Further details can be found in ISO 24276.

5 Procedure

5.1 Preparation of the test portion

5.1.1 General

Commodity-specific variables (e.g. humidity) and processing can impact the amount and quality of DNA extracted from the material under investigation. Therefore the performance characteristics of a given DNA extraction method depend on the matrix.

Take appropriate measures to ensure that the test portion is representative of the laboratory sample.

The test portion shall be of sufficient size and shall contain a sufficient number of particles to be representative of the laboratory sample **[A1] text deleted [A1]**.

For practical/technical reasons, it is not recommended to exceed a size of 2 g.

Any restrictions that arise from the size of the test portion which prevent it from being representative shall be reported and taken into consideration in the interpretation of the analytical results. The methods for DNA extraction in Annex A describe test portions from 200 mg to 500 mg, which are usually adequate for DNA-rich raw materials (e.g. ground grains, flour). However, for certain matrices containing very low amounts or degraded DNA, insufficient DNA suitable for analysis can be extracted. In these cases, the test portion may be increased.

DNA extractions shall be carried out at least on two test portions.

Storage of standards, samples and test portions shall comply with ISO 24276 and shall be organized in such a way as to preserve the biochemical parameters to be analysed (for details, see ISO/IEC 17025).

5.1.2 Samples

All operations for the preparation of test samples (e.g. grinding, homogenization, division, drying) shall be carried out in accordance with the A_1 laboratory design specified in ISO 24276:2006, 5.3.2, A_1 taking care to prevent all contamination of the sample or modification of its composition.

A_1 Laboratory samples shall be sufficiently homogeneous before reducing or grinding and before taking the test portion. A_1

For liquid samples, shake the vessel containing the sample to improve the homogenization of the product. In the case of non-homogenous products like raw oils, check that the sediments have been completely removed from the walls of the vessel.

A_1 For solid matrices that cannot easily be suspended, the matrix shall be ground to reduce the particle size and/or facilitate the extractability of DNA. In such a case, attention shall be paid to the particle size. The test portion subjected to extraction shall contain a minimum number of particles. Milling/grinding devices should be capable of being thoroughly cleaned and shall be selected in order to achieve the expected particle number and particle size distribution within the test portion. A_1

If components of the laboratory sample have been removed prior to extraction, then such procedures shall be reported.

Final food products that are solid or paste and have high lipid contents are often not easy to grind to the desired particle size in a single step. Several procedures may therefore be added, such as lipid removal using hexane after intermediate grinding, freezing or freeze-drying before grinding.

In order to facilitate the grinding of paste or viscous products, it is possible to apply one of the following treatments to certain matrices:

- heating to a maximum temperature of 40 °C;
- dissolving in an appropriate liquid such as water;
- freezing at a temperature below or equal to -20 °C.

A_1 After such treatment, homogenize the whole laboratory sample. A_1 Sample the two test portions, taking into account possible dilutions or concentrations.

During milling/grinding, precautions should be taken to ensure that the heating of the sample is kept to a minimum since heating can have a negative impact on the quality of the extracted DNA.

Milling/grinding techniques with a high risk of cross-contamination (such as the combined use of liquid nitrogen and mortar) shall be avoided as far as possible. As a rule of good practice, any dust-producing methodological step should be contained from all other analytical steps.

If salts, spices, powdered sugars and/or other substances that could potentially interfere with the extraction or analytical method are present, appropriate purification steps should be considered according to the selected method (see Annex A).

For example, in samples from composite matrices, the target matrix (e.g. the breading layer of fish sticks) can be isolated for DNA extraction.

5.2 DNA extraction/purification

5.2.1 General

The following considerations apply for the design of extraction methods.

The quality and yield of nucleic acid extracted using a given method on a given matrix should be both repeatable and reproducible in terms of analysis, provided sufficient nucleic acid is present in the matrix from which it has been extracted.

In order to obtain a good quality DNA, it is advisable, where relevant, to remove the following:

- polysaccharides (pectin, cellulose, hemi-cellulose, starch, thickeners, etc.) using appropriate enzyme treatments (e.g. pectinase, cellulase, hemi-cellulase, α -amylase) or organic extraction (e.g. CTAB/chloroform);
- RNA and/or proteins using an appropriate treatment, such as enzymatic treatment by RNase and proteinase, respectively;
- the lipid fractions using for example enzyme treatments, or solvents (e.g. *n*-hexane);
- salts (e.g. from the extraction/lysis buffer, from the precipitation step) able to interfere with the subsequent analysis.

In particular for solid or dried samples, the volume of lysis/extraction buffer should be adapted to guarantee the DNA is dissolved.

NOTE 1 DNA purification can be performed by different means such as fractionated precipitation, using solvents like phenol, chloroform, ethanol, isopropanol, and/or by adsorption on solid matrices (anion exchange resin, silica or glass gel, diatomaceous earth, membranes, etc.). Several DNA purification principles may be combined. If appropriate, extraction and purification can be performed within the same step.

Should a DNA co-precipitant such as glycogen, PEG or t-RNA be used to improve the DNA recovery during the precipitation steps, it should neither contain any detectable level of nuclease activity or PCR inhibitors/competitors, nor bear any sequence similarity with the potential PCR target under study. For genetically modified plants, a carrier DNA may be used (e.g. salmon or herring sperm DNA).

When using vacuum freeze dryers to dry the DNA pellets obtained after a precipitation step, the risk of cross contamination should be taken into account.

Re-suspend the DNA in water or in a buffer solution that prevents DNA from degradation.

A₁ EXAMPLE A tris/EDTA (TE, 1× or 0,1×) buffer adjusted to pH 8,0 is appropriate for re-suspending or diluting DNA. **A₁**

When setting up a new type of DNA extraction, or when applying one of the methods described in Annex A to a new matrix, the potential quality and integrity of the extracted DNA using the chosen protocol should be estimated by the following approach. A known quantity of a tracer DNA is added to the lysis buffer plus sample used for DNA extraction. When the chosen tracer is a predetermined amount of DNA or represents a predetermined number of copies of a particular DNA-sequence mixed to a matrix at start of DNA extraction, attention shall be paid to ascertain the lack of DNA sequence similarity between the tracer DNA and the target DNA sequence under study.

NOTE 2 The use of a tracer DNA is a good approximation to a real situation where DNA of a given matrix, complexed to other components (e.g. proteins) is expected. Such a method may also be used to estimate the presence of soluble and *trans*-acting PCR inhibitors in the extracted DNA (see ISO 24276, ISO 21569 and ISO 21570).

However, tracer DNA may give a misleading impression of recovery, since tracer DNA may be much easier to separate from matrix than the target DNA.

5.2.2 Controls

The controls to be included are described in Table 1 of ISO 24276:— These should as a minimum include an extraction blank control and a positive extraction control, but may also include an environment control.

5.2.3 Control of DNA purity: Internal PCR control

When setting up a new type of extraction, the presence of PCR inhibitors in the extracted DNA may be estimated using DNA spikes (see ISO 24276, ISO 21569 and ISO 21570). The amount of added DNA shall not exceed the maximum level supported by PCR and shall contain a definite number of target sequence copies. This number should be determined individually for each target sequence and indicated as a multiple of the existing lower limit of detection. Ideally, the target concentration of the positive control PCR should correspond to the sensitivity needed in the analysis. Care shall be taken when using highly concentrated cloned target DNA. As far as possible, the positive controls shall conform to the conditions of the test material with regard to the nucleic acids they contain.

5.3 Quantitation of the extracted DNA

5.3.1 General

The quality, integrity and amount of the nucleic acid template influences the performance of the analytical method, and hence the analytical results obtained. The limit of detection of a specific method may therefore depend on the amount of nucleic acids used. A_1 The total amount of DNA used in the PCR should be determined as well as the total amount of target taxon DNA shall be considered, as non-target taxon DNA can affect the efficiency of the PCR. A_1

Quantitation of DNA is helpful

- to compare the efficiency of different DNA extraction protocols for a given matrix (repeatability), and
- to measure the concentration of nucleic acids prior to analysis.

5.3.2 Range of application

Each method of quantitation shall be applied within its dynamic range, also considering its level of precision.

5.3.3 Quantity standards

The accuracy of the quantitation methods depends on the nucleic acid standards used to calibrate the method.

If using a method that is sensitive to the size and/or quality of the nucleic acid fragments, then the nucleic acid standards that match the size and/or quality of the expected nucleic acid as extracted from the sample shall be used.

The reference material used should ensure traceability to stated references, usually national or international Standards, through an unbroken chain of comparison [see ISO Guide 30].

When a method using intercalating agents is employed, high molecular mass DNA standard should be used when high molecular mass DNA is to be quantified. Low molecular mass DNA should be used when low molecular mass DNA is to be quantified. High molecular mass nucleic acid usually also contains a certain amount of lower molecular mass fragments. This means that many methods for DNA quantitation suffer from a certain degree of inaccuracy, which should be taken into account.

NOTE Additionally, depending on the matrix and type of extraction method, a certain portion of the extracted DNA may be recovered as single-stranded DNA (with much poorer intercalation capacity), leading to an underestimation of the overall DNA content. In contrast, single-stranded DNA is equally well detected by physical measurements.

At least three points (preferably replicated) are required for the construction of a good calibration curve. The amount of standard DNA used for each calibration point depends on the sensitivity of the method and on the dynamic range under consideration.

5.4 Stability of extracted DNA

The DNA extracted shall be stored under such conditions that the stability is ensured to perform the subsequent analyses.

Repeated freezing and thawing of DNA solutions should be avoided.

6 Interpretation

The DNA extraction method employed shall be appropriate to obtain the quality and quantity of nucleic acid required for the subsequent analysis.

The quality of the extracted nucleic acid should be assessed using an analytical method that is influenced by the same quality parameters as the analysis to be performed on the extracted nucleic acid $[A_1]$ (e.g. if the analysis to be performed is PCR, the quality of the extracted DNA should be assessed using adequate PCR controls). $[A_1]$

Further parameters for method compatibility can be found in ISO 21569, ISO 21570 and ISO 24276.

7 Test report

When issuing the final test report in accordance with ISO 24276, the following additional information to document the activity of the laboratory shall be included:

- a statement describing the derivation of the test portions, and any preliminary processing of the sample before nucleic acid extraction;
- the size of the test portions used for the nucleic acid extraction;
- the nucleic acid extraction method used;
- any special observations made during testing;
- any operation not specified in the method or considered to be optional but that can have an effect on the results;
- the interpretation of the results;
- the experimenter's name.

Handling and storage of raw data are described in ISO/IEC 17025 and related quality assurance schemes. Consistency should be achieved.

Annex A (informative)

Methods for DNA extraction

A.1 Preparation of PCR-quality DNA using phenol/chloroform-based DNA extraction methods

A.1.1 Basic phenol/chloroform method

A.1.1.1 General

This routine method (see Reference [5]) is suitable for the extraction of DNA from a wide variety of matrices (see A.1.1.8).

Phenol is usually very suitable for nuclease destruction and protein denaturation.

When foliar or green material (e.g. chicory leaves, dried alfalfa) is investigated, many PCR inhibitors may also be co-precipitated together with DNA. For this reason, difficulties may be encountered in obtaining PCR-amplifiable DNA reproducibly.

The corrosive hazardous properties of phenol must be taken into consideration, thus the use of DNA extraction methods based on CTAB and/or PVP and/or silica adsorption are favoured as primary alternatives.

A.1.1.2 Validation status

The method has been widely applied in all areas of biology, agronomy and medicine, over the past 40 years, but has never been evaluated through interlaboratory studies for GMO detection in foodstuffs.

A.1.1.3 Principle

The method consists of a lysis step (thermal lysis in presence of sodium dodecyl sulfate and a high EDTA content) followed by the removal of contaminants (such as lipophylic molecules, polysaccharides and proteins) and nucleases from the DNA-containing aqueous phase using phenol and chloroform. A final DNA precipitation with ethanol concentrates the DNA and eliminates salts and residual chloroform. The critical step of the method is the lysis step [5].

A.1.1.4 Safety precautions

A fume hood is necessary for handling organic chemicals.

A.1.1.5 Reagents

A.1.1.5.1 Ethanol, volume fraction ϕ (C₂H₅OH) = 96 %

Store and use at -20 °C.

- A.1.1.5.2 Glacial acetic acid** (CH_3COOH).
- A.1.1.5.3 Potassium acetate** ($\text{C}_2\text{H}_3\text{O}_2\text{K}$).
- A.1.1.5.4 Hydrochloric acid**, ϕ (HCl) = 37 %.
- A.1.1.5.5 Isoamyl alcohol** [$(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{OH}$].
- A.1.1.5.6 Phenol** ($\text{C}_6\text{H}_5\text{OH}$).
- A.1.1.5.7 Chloroform** (CHCl_3).
- A.1.1.5.8 Tris(hydroxymethyl)-aminomethane** (Tris) ($\text{C}_4\text{H}_{11}\text{NO}_3$).
- A.1.1.5.9 Ethylenediaminetetraacetic acid dipotassium salt** (K_2EDTA) ($\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_8\text{K}_2$).
- A.1.1.5.10 Potassium hydroxide** (KOH).
- A.1.1.5.11 Potassium chloride** (KCl).
- A.1.1.5.12 Sodium dodecyl sulfate** (SDS) ($\text{C}_{12}\text{H}_{25}\text{O}_4\text{SNa}$).
- A.1.1.5.13 Proteinase K**, approximately 20 Units/mg lyophilisate.
- A.1.1.5.14 RNase-A**, DNase-free, from bovine pancreas, approximately 50 Kunitz Units/mg of lyophilisate.
- A.1.1.5.15 Equilibrated phenol**, pH > 7,8.

Use phenol equilibrated against extraction buffer (A.1.1.5.18) without SDS, or prepared according to Reference [5], or according to the manufacturers recommendations.

A.1.1.5.16 Chloroform-isoamyl alcohol

Mix 24 volume parts of chloroform (A.1.1.5.7) with 1 volume part of isoamyl alcohol (A.1.1.5.5).

A.1.1.5.17 Phenol-chloroform-isoamyl alcohol

Mix 1 volume part of equilibrated phenol (A.1.1.5.15) with 1 volume part of the chloroform-isoamyl alcohol solution (A.1.1.5.16).

A.1.1.5.18 Extraction/lysis buffer, substance concentration $c(\text{Tris}) = 0,050 \text{ mol/l}$, $c(\text{K}_2\text{EDTA}) = 0,050 \text{ mol/l}$, mass concentration $\rho(\text{SDS}) = 30 \text{ g/l}$.

Adjust the pH to 8,0 with HCl or KOH.

A.1.1.5.19 TE buffer, $c(\text{Tris}) = 0,010 \text{ mol/l}$, $c(\text{K}_2\text{EDTA}) = 0,001 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or KOH.

A.1.1.5.20 Proteinase-K solution, $\rho = 20 \text{ mg/ml}$, dissolved in sterile water.

Do not autoclave. Store at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.1.1.5.21 RNase-A solution, $\rho = 10$ mg/ml lyophilisate.

Store at -20 °C, but avoid repeated freezing and thawing.

A.1.1.5.22 Ethanol solution, ϕ (C₂H₅OH) = 70 %.

Store and use at -20 °C.

A.1.1.5.23 Potassium acetate solution, c (C₂H₃O₂K) = 3 mol/l.

Adjust the pH to 5,2 with glacial acetic acid. Do not autoclave. If necessary, filter through a 0,22 μ m filter.

A.1.1.6 Apparatus and equipment

Usual laboratory equipment and, in particular, the following.

A.1.1.6.1 Centrifuge, capable of achieving a minimum acceleration of 10 000 *g*.

In some steps a refrigerated centrifuge is required.

A.1.1.6.2 Water bath or incubator, working in a temperature range from 60 °C to 70 °C.

A.1.1.6.3 Vacuum dryer (optional).

A.1.1.6.4 Freeze dryer (optional).

A.1.1.6.5 Mixer, e.g. Vortex^{®1)}

A.1.1.6.6 Reaction vessels, resistant to freezing in liquid nitrogen.

A.1.1.7 Procedure

A.1.1.7.1 General

Once the matrix test portion has been prepared, apply the following DNA extraction/purification protocol. Scale-adaptation of masses and buffer volumes is required as a function of the selected size of the test portion.

A.1.1.7.2 Extraction procedure

Weigh 0,25 g of the test sample into a microtube.

Add 1,6 ml of extraction buffer (A.1.1.5.18) and, when necessary (e.g. in protein-rich matrices), 50 μ l of proteinase K solution (A.1.1.5.20.) Incubate at 60 °C to 70 °C, usually for between 30 min to 2 h (overnight incubation is also possible). Add RNase A (A.1.1.5.21) up to a final concentration of 0,1 μ g/ml. Centrifuge at 5 000 *g* for 30 min and recover the supernatant in a fresh tube. Add 1 volume of equilibrated phenol (A.1.1.5.15) to the supernatant, then mix gently and thoroughly. Centrifuge at 5 000 *g* for 15 min and recover the upper aqueous phase in a fresh tube. Add 1 volume of phenol-chloroform isoamyl alcohol (A.1.1.5.17) to the supernatant, then mix gently and thoroughly. Centrifuge at 5 000 *g* for 15 min and recover the aqueous phase in a fresh tube. Repeat this step once or more times (depending on the matrix) until the interface between the phases is clean.

1) Vortex is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product. Equivalent products may be used if they can be shown to lead to the same results.

Add 1 volume of chloroform/isoamyl alcohol (A.1.1.5.16) to the supernatant, then mix gently and thoroughly. Centrifuge at 5 000 *g* for 10 min and recover the upper aqueous phase in a fresh tube. Repeat, if necessary, until the interface between the phases is clear. Mix the supernatant with 0,1 volume of potassium acetate solution (A.1.1.5.23.) and 2,5 volumes of 96 % ethanol (A.1.1.5.1), then mix thoroughly by inversion. Incubate for at least 5 min in liquid nitrogen, or 1 h at –80 °C, or overnight at –20 °C. Centrifuge at 10 000 *g* (or up to 13 000 *g*) at 4 °C for at least 15 min, then carefully discard the supernatant.

Carefully wash the DNA pellet with 2 volumes of 70 % ethanol solution (A.1.1.5.22). Centrifuge at 10 000 *g* to 13 000 *g* at 4 °C for 15 min, then discard carefully the supernatant. This step is essential for the removal of the precipitating salts that could interfere with the subsequent analysis (e.g. PCR).

Dry the pellet and re-dissolve it in 100 µl of water or appropriate buffer, e.g. TE buffer (A.1.1.5.19). This is the DNA master stock. Add RNase-A (A.1.1.5.21) up to a final concentration of 0,1 µg/ml.

A.1.1.8 List of examples

The method has been successfully applied to extract DNA²⁾ from the following matrices:

acidified soya beans²⁾, dehydrated alfalfa, baby biscuits²⁾, baby milk²⁾, bacteria and spores thereof, barley seeds, beef/pork paté²⁾, beer²⁾, blue cheese, brownies²⁾, canned corn, carrot seeds, cereal bars²⁾, cheese, chicken nuggets, chicory leaves, chicory roots, chocolate cookies²⁾, chocolate paste²⁾, cinnamon cookies²⁾, compotes, cornflakes²⁾, cracked rice, dessert cream²⁾, dried pea seeds, maize biscuits²⁾, maize feeding oil cakes, maize flour, maize gluten feed, maize seeds, manioc hard pellets for feed, manioc tapioca meat, fresh and cooked²⁾ meats (beef, pork, chicken and turkey), melon fruit pulp, melon seeds, minced meat, muesli ingredients²⁾, muesli²⁾, mung bean sprouts²⁾, oat seeds, potato tuber, rapeseed feeding oil cake, glupacolza, rapeseed seeds, sausages (slicing²⁾ and cocktail²⁾ (see A.1.2 for an improved extraction method), schnitzel, soeie hoi sin²⁾, soup balls, soya protein in meat preparations²⁾, soya lecithin (raw brown and yellow refined²⁾), soya sprouts²⁾, soya drinks, soya beans, soya bean cream, soya bean feeding oil cake, soya bean tofu, spaghetti sauce²⁾, spelt seeds, sugar beet (dried pulp), sugar beet (fresh root), sugar beet seeds, sunflower seeds, tofu, tomato fresh fruit, tomato purée²⁾, tomato seeds, vegetarian hamburger, waffles (with²⁾ and without²⁾ chocolate), wheat bran, wheat flour, wheat gluten feed, wheat seeds, wheat semolina, yoghurt²⁾ (see A.1.3 for an improved extraction method).

A.1.2 Phenol/chloroform method: Protocol for starter cultures of fermented sausages

A.1.2.1 General

This method is designed to isolate the total DNA, including bacterial genomic DNA, from sausages. The applicability of the method to obtain DNA of high quality suitable for the specific detection of recombinant DNA by PCR has been demonstrated for fermented sausages^[6] as well as for thermally treated fermented sausages, so-called summer sausages^[7]. In addition, the extraction method was shown to be suitable to isolate total DNA from cream to detect specifically *Staphylococcus aureus* in this food matrix^[8]. (For a summary of the matrices for which the method is suitable, see A.1.2.8.)

A.1.2.2 Validation status

This method has been validated by an interlaboratory study on a test portion of 0,4 g (see A.1.2.9).

2) Repeatability may depend on the batch of the matrix and/or to its production technology. In some cases, DNA could not be found or was degraded in such a way that PCR results were below the limit of detection of the method, irrespective of the PCR primers/protocols used. This may be a source of low reproducibility between laboratories.

A.1.2.3 Principle

The method consists of the recovery of bacterial cells from the food matrix by homogenization of the sausage samples, followed by a centrifugation step. The sediment contains not only bacterial cells but also meat particles. For specific lysis of the bacterial cells, the cell walls are degraded by addition of lysozyme. To improve the degradation of cell walls of meat lactobacilli, mutanolysin can be added. Complete lysis of the cells is performed by addition of the detergent SDS (sodium dodecyl sulfate) and proteinase-K, followed by several extractions of the aqueous phase with phenol and/or chloroform. The phenol/chloroform extraction step is important to eliminate any nuclease activities and PCR-inhibiting substances including those arising from the food matrix (e.g. haematin). A final precipitation of DNA by ethanol is performed.

A.1.2.4 Safety precautions

A fume hood is necessary for handling organic chemicals.

A.1.2.5 Reagents

A.1.2.5.1 Isopropanol [$\text{CH}_3\text{CH}(\text{OH})\text{CH}_3$].

A.1.2.5.2 Ethanol, $\varphi(\text{C}_2\text{H}_5\text{OH}) = 96 \%$.

Store and use at $-20 \text{ }^\circ\text{C}$.

A.1.2.5.3 Glacial acetic acid (CH_3COOH).

A.1.2.5.4 Hydrochloric acid, $\varphi(\text{HCl}) = 37 \%$.

A.1.2.5.5 Sodium hydroxide (NaOH).

A.1.2.5.6 Isoamyl alcohol [$(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{OH}$].

A.1.2.5.7 Phenol ($\text{C}_6\text{H}_5\text{OH}$).

A.1.2.5.8 Chloroform (CHCl_3).

A.1.2.5.9 Tris(hydroxymethyl)-aminomethane (Tris) ($\text{C}_4\text{H}_{11}\text{NO}_3$).

A.1.2.5.10 Ethylenediaminetetraacetic acid disodium salt (Na_2EDTA) ($\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_8\text{Na}_2$).

A.1.2.5.11 Sodium dodecyl sulfate (SDS) ($\text{C}_{12}\text{H}_{25}\text{O}_4\text{SNa}$).

A.1.2.5.12 Lysozyme

50 000 U/mg protein (1 U will produce a ΔA_{450} of 0,001 per minute at pH 6,24 and $25 \text{ }^\circ\text{C}$, using a suspension of *Micrococcus lysodeikticus* as substrate, in a 2,6 ml reaction mixture using a 1 cm light path).

A.1.2.5.13 Sucrose ($\text{C}_{12}\text{H}_{22}\text{O}_{11}$).

A.1.2.5.14 Proteinase-K, approximately 20 Units/mg lyophilisate.

A.1.2.5.15 Sodium acetate ($\text{C}_2\text{H}_3\text{O}_2\text{Na}$).

A.1.2.5.16 Equilibrated phenol, saturated with Tris/HCl (pH > 7,8) buffer, or prepared according to Reference [5], or according to the manufacturer's recommendations.

A.1.2.5.17 Chloroform-isoamyl alcohol

Mix 24 volume parts of chloroform (A.1.2.5.8) with 1 volume part of isoamyl alcohol (A.1.2.5.6).

A.1.2.5.18 Phenol-chloroform-isoamyl alcohol

Prepare by mixing 25 volume parts of equilibrated phenol (A.1.2.5.16) with 24 volume parts of chloroform (A.1.2.5.8) and 1 volume part of isoamyl alcohol (A.1.2.5.6).

A.1.2.5.19 Mutanolysin-solution, containing 500 U/ml or 5 000 U/ml mutanolysin, dissolved in sterile water.

Do not autoclave. Store at $-20\text{ }^{\circ}\text{C}$, but avoid repeated freezing and thawing.

A.1.2.5.20 Lysozyme solution, containing 10 mg/ml, dissolved in sterile water.

Do not autoclave. Store at $-20\text{ }^{\circ}\text{C}$, but avoid freezing and thawing.

A.1.2.5.21 Sucrose solution, $\rho(\text{C}_{12}\text{H}_{22}\text{O}_{11}) = 400\text{ g/l}$.

A.1.2.5.22 Buffer solution A, $c(\text{Tris}) = 0,020\text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,020\text{ mol/l}$, $c(\text{NaCl}) = 0,1\text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.1.2.5.23 Extraction/lysis buffer, containing one volume part of buffer solution A (A.1.2.5.22) and one volume part of sucrose solution (A.1.2.5.21).

A.1.2.5.24 SDS solution, $\rho(\text{SDS}) = 250\text{ g/l}$.

A.1.2.5.25 Proteinase-K solution, containing 20 mg/ml, stored at $-20\text{ }^{\circ}\text{C}$, dissolved in sterile water.

Do not autoclave. Store at $-20\text{ }^{\circ}\text{C}$, but avoid repeated freezing and thawing.

A.1.2.5.26 Ethanol solution, $\phi(\text{C}_2\text{H}_5\text{OH}) = 70\text{ }%$.

Store and use at $-20\text{ }^{\circ}\text{C}$.

A.1.2.5.27 Sodium acetate solution, $c(\text{C}_2\text{H}_3\text{O}_2\text{Na}) = 3\text{ mol/l}$.

Adjust the pH to 5,2 with glacial acetic acid.

A.1.2.5.28 TE buffer, $c(\text{Tris}) = 0,010\text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,001\text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.1.2.6 Apparatus and equipment

Usual laboratory equipment and, in particular, the following.

A.1.2.6.1 Instrument and/or material for chopping tissue (e.g. scalpel).

A.1.2.6.2 Centrifuge, capable of achieving an acceleration of 12 000 g.

In some steps a refrigerated centrifuge is required.

A.1.2.6.3 Water bath or incubator.

A.1.2.6.4 Vacuum dryer (optional).

A.1.2.6.5 Mixer, e.g. Vortex^{®2}.

A.1.2.7 Procedure

A.1.2.7.1 General

Once the matrix test portion has been prepared, apply the following DNA extraction/purification protocol. Scale-adaptation of masses and buffer volumes is required as a function of the selected size of the test portion.

A.1.2.7.2 Sample preparation

Chop the meat sausage, homogenize it and add 200 mg to 500 mg to 3 volumes of water (up to 1,5 ml). Keep at room temperature for about 10 min.

A.1.2.7.3 Extraction procedure

Carefully transfer 500 µl of the aqueous phase (suspension) to a new reaction vessel. Centrifuge for 10 min at 12 000 *g*.

Discard the supernatant and re-suspend the pellet in 500 µl of extraction/lysis buffer (A.1.2.5.23).

Add 50 µl of lysozyme solution (A.1.2.5.20). Incubate at 37 °C for 1 h. If the results are not satisfactory, lysozyme may be combined with 10 U of mutanolysin (A.1.2.5.19), but the matrix-specific effect of this should be tested prior to routine application.

Add 25 µl of SDS solution (A.1.2.5.24) and 25 µl of proteinase-K solution (A.1.2.5.25), then incubate for 10 min at 60 °C.

Add 1 volume of phenol/chloroform/isoamyl alcohol (A.1.2.5.18) and mix.

Centrifuge the mixture for 3 min at about 12 000 *g*. Transfer the upper aqueous phase to a new vial.

Add 1 volume of chloroform-isoamyl alcohol (A.1.2.5.17) and mix.

Centrifuge for 3 min at about 12 000 *g*. Transfer the upper phase to a new reaction vessel.

Add 0,1 volume of sodium acetate solution (A.1.2.5.27) and 1 volume of isopropanol (A.1.2.5.1). Mix gently several times by inversion.

Keep at room temperature for at least 30 min. Centrifuge for 15 min at about 12 000 *g*. Decant the supernatant and discard it.

Wash the pellet carefully with at least 500 µl of ethanol solution (A.1.2.5.26), shaking gently. Centrifuge the mixture for 10 min at about 12 000 *g*. This step is essential for the removal of the precipitating salts that could interfere with the subsequent analysis (e.g. PCR).

Discard the supernatant.

Dry the pellet and re-dissolve it in 100 µl of water or appropriate buffer, e.g. TE buffer (A.1.2.5.28). This is the DNA master stock.

A.1.2.8 List of examples

See Table A.1.

Table A.1 — List of matrices for which the method has been successfully applied

Matrices successfully analysed	Microorganism	Reference
Fermented sausage	<i>Lactobacillus curvatus</i>	[6]
Summer sausage (thermally treated)	<i>Lactobacillus curvatus</i>	[7]
Cream	<i>Staphylococcus aureus</i>	[8]

A.1.2.9 Validation

The validation data in Table A.2 have been elaborated in a collaborative study carried out by the working group “Development of methods for identifying foodstuffs produced by genetic engineering techniques” of the Commission of the former German Federal Health Board for the implementation of methods according to article 35 of the German Foodstuffs Act^[6].

In this collaborative study, two samples were false positive, probably caused by incorrect packing. Mutanolysin was not used for this study.

Table A.2 — Validation data

Number of laboratories participating	Number of sausage samples per laboratory	Number of total samples	Number of samples correctly identified
15	10	150	148 (control samples and GMO samples)

A.1.3 Phenol/chloroform method: Protocol for starter cultures of yoghurt

A.1.3.1 General

This method describes a procedure for the extraction of bulk DNA from starter cultures used for the fermentation of the milk product yoghurt. This procedure was successfully carried out with plain yoghurts, yoghurts containing different ingredients such as fruits, additives and stabilizers and with products of different fat content (see A.1.3.8 and References [9], [10], [11]). DNA of PCR quality was also extracted from yoghurts that were thermally treated ^[12].

A.1.3.2 Validation status

This method has been validated in an interlaboratory study; see A.1.3.9.

A.1.3.3 Principle

This method is basically a phenol-chloroform extraction procedure that is adapted for a special food matrix. The Gram-positive starters of the yoghurts are harvested after resolving the coagulated casein at alkaline pH. The recovered cells are re-suspended in a buffered aqueous solution and treated with lysozyme (and mutanolysin) in order to damage the cell walls. Cell lysis is carried out by adding an ionic detergent such as sodium dodecylsulfate (SDS). Proteins are removed by proteinase-K treatment and several subsequent phenol/chloroform and chloroform steps. DNA is finally precipitated with alcohol.

A.1.3.4 Safety precautions

A fume hood is necessary for handling organic chemicals.

A.1.3.5 Reagents

A.1.3.5.1 Isopropanol [CH₃CH(OH)CH₃].

A.1.3.5.2 Ethanol, $\phi(\text{C}_2\text{H}_5\text{OH}) = 96 \%$.

Store and use at $-20 \text{ }^\circ\text{C}$.

A.1.3.5.3 Glacial acetic acid (CH₃COOH).

A.1.3.5.4 Sodium chloride (NaCl).

A.1.3.5.5 Sodium citrate (C₆H₅Na₃O₇).

A.1.3.5.6 Hydrochloric acid, $\phi(\text{HCl}) = 37 \%$.

A.1.3.5.7 Sodium hydroxide (NaOH).

A.1.3.5.8 Isoamyl alcohol [(CH₃)₂CHCH₂CH₂(OH)].

A.1.3.5.9 Phenol (C₆H₅OH).

A.1.3.5.10 Chloroform (CHCl₃).

A.1.3.5.11 Tris(hydroxymethyl)-aminomethane (Tris) (C₄H₁₁NO₃).

A.1.3.5.12 Ethylenediaminetetraacetic acid disodium salt (Na₂EDTA) (C₁₀H₁₄N₂O₈Na₂).

A.1.3.5.13 Sodium dodecyl sulfate (SDS) (C₁₂H₂₅O₄SNa).

A.1.3.5.14 Lysozyme, 50 000 U/mg protein (1 U will produce a ΔA_{450} of 0,001 per minute at pH 6,24 and 25 °C, using a suspension of *Micrococcus lysodeikticus* as substrate, in a 2,6 ml reaction mixture with 1 cm light path).

A.1.3.5.15 Sucrose (C₁₂H₂₂O₁₁).

A.1.3.5.16 Proteinase-K, approximately 20 Units/mg lyophilisate.

A.1.3.5.17 Sodium acetate (C₂H₃O₂Na).

A.1.3.5.18 Sodium citrate solution, $\rho(\text{C}_6\text{H}_5\text{Na}_3\text{O}_7) = 400 \text{ g/l}$.

A.1.3.5.19 Sodium hydroxide solution, $c(\text{NaOH}) = 0,4 \text{ mol/l}$.

Dissolve in sterile water, but do not autoclave. Prepare freshly before use.

A.1.3.5.20 Sodium chloride/sodium citrate solution (SSC, concentrated 5×), $c(\text{NaCl}) = 0,75 \text{ mol/l}$, $c(\text{C}_6\text{H}_5\text{Na}_3\text{O}_7) = 0,075 \text{ mol/l}$.

It is advisable to prepare the SSC solution as a concentrated stock solution (e.g. SSC 20×), because solutions with a high salt concentration are usually more stable. Dilute before use.

A.1.3.5.21 Equilibrated phenol, adjusted to pH 8, saturated with Tris/HCl (pH > 7,8) buffer, or for example, prepared according to Reference [5], or according to the manufacturer's recommendations.

A.1.3.5.22 Chloroform-isoamyl alcohol

Mix 24 volume parts of chloroform (A.1.3.5.10) with 1 volume part of isoamyl alcohol (A.1.3.5.8).

A.1.3.5.23 Phenol-chloroform-isoamyl alcohol

Prepare by mixing 25 volume parts of equilibrated phenol (A.1.3.5.21) with 24 volume parts of chloroform (A.1.3.5.10) and 1 volume part of isoamyl alcohol (A.1.3.5.8).

A.1.3.5.24 Mutanolysin solution, containing 500 U/ml or 5 000 U/ml mutanolysin, dissolved in sterile water.

Do not autoclave. Store at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.1.3.5.25 Lysozyme solution, containing 10 mg/ml of lysozyme, dissolved in sterile water.

Do not autoclave. Store at $-20 \text{ }^\circ\text{C}$, but avoid freezing and thawing.

A.1.3.5.26 Sucrose solution, $c(\text{C}_{12}\text{H}_{22}\text{O}_{11}) = 400 \text{ g/l}$.

A.1.3.5.27 Buffer solution A, $c(\text{Tris}) = 0,020 \text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,020 \text{ mol/l}$, $c(\text{NaCl}) = 0,100 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.1.3.5.28 Extraction/lysis buffer, containing 1 volume part of buffer solution A (A.1.3.5.27) and 1 volume part of sucrose solution (A.1.3.5.26).

A.1.3.5.29 SDS Solution, $\rho(\text{SDS}) = 250\text{g/l}$.

A.1.3.5.30 Proteinase-K solution, containing 20 mg/ml, dissolved in sterile water.

Do not autoclave. Store at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.1.3.5.31 Ethanol solution, $\phi(\text{C}_2\text{H}_5\text{OH}) = 70 \text{ } \%$.

Store and use at $-20 \text{ }^\circ\text{C}$.

A.1.3.5.32 Sodium acetate solution, $c(\text{C}_2\text{H}_3\text{O}_2\text{Na}) = 3 \text{ mol/l}$.

Adjust the pH to 5,2 with glacial acetic acid.

A.1.3.5.33 TE buffer, $c(\text{Tris}) = 0,010 \text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,001 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.1.3.6 Apparatus and equipment

Usual laboratory equipment and, in particular, the following.

A.1.3.6.1 Centrifuge, capable of achieving an acceleration of 12 000 *g*. In some steps a refrigerated centrifuge is required.

A.1.3.6.2 Water bath or incubator.

A.1.3.6.3 Vacuum dryer (optional).

A.1.3.6.4 Mixer, e.g. Vortex^{®2}).

A.1.3.7 Procedure

A.1.3.7.1 General

Once the matrix test portion is prepared, apply the following DNA extraction/purification protocol. Some scale-adaptation of masses and buffer volumes is required as a function of the selected size of the test portion.

A.1.3.7.2 Extraction procedure

Shake the yoghurt well. Transfer 250 µl of yoghurt to a 2-ml reaction vessel. Add 80 µl of sodium citrate solution (A.1.3.5.18). Add 150 µl of NaOH solution (A.1.3.5.19) and mix well. Centrifuge at about 12 000 *g* for 2 min.

The pellet should not be much bigger than 0,7 cm in diameter and about 100 µl in volume. Otherwise these steps (adding 80 µl of sodium citrate solution and 150 µl of NaOH) should be repeated.

Discard the upper layer of fat and the aqueous supernatant and re-suspend the pellet in 500 µl of 5× SSC solution (A.1.3.5.20). Centrifuge for at least 2 min at about 12 000 *g* and discard the supernatant. Resuspend the pellet in 500 µl of 5× SSC solution. Centrifuge for 2 min at about 12 000 *g* and discard the supernatant.

Resuspend the pellet in 500 µl of extraction/lysis buffer (A.1.3.5.28). Add 50 µl of lysozyme solution (A.1.3.5.25). Incubate at 37 °C for 1 h. If results are not satisfying, lysozyme may be combined with 10 U of mutanolysin (A.1.3.5.24), but the matrix-specific effect of this should be tested prior to routine application.

Add 25 µl of SDS solution (A.1.3.5.29) and 25 µl of proteinase-K solution (A.1.3.5.30). Incubate for 10 min at 60 °C. Add 500 µl of phenol/chloroform/isoamyl alcohol (A.1.3.5.23) and mix. Centrifuge for 3 min at about 12 000 *g*. Transfer the upper aqueous phase to a new reaction vessel. Add 1 volume of chloroform/isoamyl alcohol (A.1.3.5.22) and mix. Centrifuge for 3 min at about 12 000 *g*.

Transfer the upper phase to a new reaction vessel. Add 0,1 volume of sodium acetate solution (A.1.3.5.32) and 1 part by volume of isopropanol (A.1.3.5.1). Keep for at least 30 min at room temperature. Centrifuge for 15 min at about 12 000 *g*. Discard the supernatant. Wash the pellet carefully in at least 500 µl of ethanol solution (A.1.3.5.31). Centrifuge for 10 min at about 12 000 *g*. This step is essential for the removal of the precipitating salts that could interfere with the subsequent analysis (e.g. PCR). Discard the supernatant.

Dry the pellet and re-dissolve it in 100 µl of water or appropriate buffer, e.g. TE buffer (A.1.2.5.33). This is the "DNA master stock".

A.1.3.8 List of examples

See Table A.3.

Table A.3 — List of matrices for which the method has been successfully applied

Matrices successfully analysed	Contents, additives etc.	Micro-organism	Reference
Plain yoghurt	0,3 % fat, 3,5 % fat	<i>Streptococcus thermophilus</i>	[9], [11]
Fruit yoghurts	1,5 % fat, modified starch, hazelnuts, gelatine 1,5 % fat, 3,8 % protein, aspartam, acesulfam, pineapple 3,5 % fat, flavour, gelatine, peach, coconut 10 % fat, modified starch, lemon, flavour, almonds, pectin, carotin, riboflavin	<i>Streptococcus thermophilus</i>	[9]
Thermally treated plain yoghurt	3,5 % fat	<i>Streptococcus thermophilus</i>	[12]

A.1.3.9 Validation

The validation data in Table A.4 have been elaborated in a collaborative study carried out by the working group “Development of methods for identifying foodstuffs produced by genetic engineering techniques” of the Commission of the former German Federal Health Board for the implementation of methods according to article 35 of the German Foodstuffs Act [11].

At this collaborative trial, two laboratories did not carry out the verification by hybridization. *Mutanolysin* was not used for the trial.

Table A.4 — Validation data

Number of laboratories participating	Number of yoghurt samples per laboratory	Number of total samples	Number of samples correctly identified
20	10	200	200 (99 control samples and 101 GMO samples)

A.1.4 Phenol/chloroform method: Protocol for yeasts and/or filamentous fungi harvested from foodstuffs

A.1.4.1 General

This method describes a one-step extraction and purification of PCR-quality DNA from yeast or filamentous fungi^[13], or isolated microbial populations. It is well suited for the tracing of the DNA from genetically modified microorganisms in highly complex matrices^[14], [15]. It can be used either to extract total DNA from the matrix^[14], [15], or from the microbial fraction either directly segregated from a matrix or harvested from starter cultures (liquid or agar colonies).

NOTE Separation of the microbial fraction from the test portion yields the most reliable results if done prior to DNA extraction.

Total DNA from the matrix may also be extracted by the alternative protocols of Annex A. However, these protocols do not guarantee that DNA from all microorganisms (especially from lysis-resistant fungi, or Gram-negative bacteria) would be randomly isolated with a reliable yield. Although the present protocol may be used for total DNA extraction in matrices such as yoghurt, milk or cheese, the extraction efficiency has been shown to be reliable for extensively ground or milled solid matrices only and shall always be checked on a matrix basis.

A.1.4.2 Validation status

This DNA extraction method has been applied and validated^[13] for 25 fungal genera representing 325 species (including yeasts used for bakery or wine production, and *Penicillia* spp.^[16] used by blue cheese producers) under mycelium or spore forms (among which the most breakage- or lysis-resistant species such as *Aspergillus fumigatus* and *Cryptococcus neoformans*). The method has been designed to avoid inter-portion and laboratory contamination making it suitable for large-scale routine DNA extractions and PCR applications. No variation in template quality in qualitative PCR was observed after long-term storage at $-20\text{ }^{\circ}\text{C}$ after up to 5 years, or between different DNA preparations from the same organism. Although the quality of the DNA is suitable for qualitative PCR, it can be poorly digestible with some restriction enzymes. For purposes such as quantitative PCR, the DNA obtained with the present method shall be further purified using another biochemical principle such as the one described in A.4 as an example.

A1 Text deleted **A1**

A.1.4.3 Principle

Basically, bacteria, yeasts or mycelia are disrupted and the DNA is simultaneously extracted by high-speed agitation in the presence of glass beads in a mixture of Tris-phenol-chloroform-EDTA-SDS followed by precipitation with ethanol.

A.1.4.4 Safety precautions

A fume hood is necessary for handling organic chemicals.

A.1.4.5 Reagents

A.1.4.5.1 Ethanol, $\phi(\text{C}_2\text{H}_5\text{OH}) = 96\%$.

Store and use at $-20\text{ }^{\circ}\text{C}$.

A.1.4.5.2 Glacial acetic acid (CH_3COOH).

A.1.4.5.3 Sulfuric acid, $\phi(\text{H}_2\text{SO}_4) > 90\%$.

A.1.4.5.4 Potassium bicarbonate (KHCO_3).

A.1.4.5.5 Potassium acetate ($\text{C}_2\text{H}_3\text{O}_2\text{K}$).

A.1.4.5.6 Hydrochloric acid, $\phi(\text{HCl}) = 37\%$.

A.1.4.5.7 Isoamyl alcohol [$(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{OH}$].

A.1.4.5.8 Phenol ($\text{C}_6\text{H}_5\text{OH}$).

A.1.4.5.9 Chloroform (CHCl_3).

A.1.4.5.10 Tris(hydroxymethyl)-aminomethane (Tris) ($\text{C}_4\text{H}_{11}\text{NO}_3$).

A.1.4.5.11 Ethylenediaminetetraacetic acid dipotassium salt (K_2EDTA) ($\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_8\text{K}_2$).

A.1.4.5.12 Potassium hydroxide (KOH).

A.1.4.5.13 Sodium dodecyl sulfate (SDS) ($\text{C}_{12}\text{H}_{25}\text{O}_4\text{SNa}$).

A.1.4.5.14 RNase-A, DNase-free, from bovine pancreas, approximately 50 Kunitz Units/mg lyophilisate.

A.1.4.5.15 Equilibrated phenol, adjusted to pH > 7,8.

Phenol (A.1.4.5.8) prepared according to, for example, Reference [5] and (optional) finally equilibrated against extraction buffer (A.1.4.5.18) without SDS, or according to the manufacturer's recommendations.

A.1.4.5.16 Chloroform-isoamyl alcohol

Mix 24 volume parts of chloroform (A.1.4.5.9.) with 1 volume part of isoamyl alcohol (A.1.4.5.7).

A.1.4.5.17 Phenol-chloroform-isoamyl alcohol

Prepare by mixing 1 volume part of equilibrated phenol (A.1.4.5.15) with 1 volume part of chloroform-isoamyl alcohol (A.1.4.5.16).

A.1.4.5.18 Extraction/lysis buffer, $c(\text{Tris}) = 0,050 \text{ mol/l}$, $c(\text{K}_2\text{EDTA}) = 0,050 \text{ mol/l}$, $\rho(\text{SDS}) = 30 \text{ g/l}$.

Adjust the pH to 8,0 with HCl or KOH.

A.1.4.5.19 TE buffer, $c(\text{Tris}) = 0,010 \text{ mol/l}$, $c(\text{K}_2\text{EDTA}) = 0,001 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or KOH.

A.1.4.5.20 RNase-A solution, $\rho(\text{RNase-A}) = 10 \text{ mg/ml}$ lyophilisate.

Store at $-20 \text{ }^\circ\text{C}$.

A.1.4.5.21 Ethanol solution, $\phi(\text{C}_2\text{H}_5\text{OH}) = 70 \%$.

Store and use at $-20 \text{ }^\circ\text{C}$.

A.1.4.5.22 Potassium acetate solution, $c(\text{C}_2\text{H}_3\text{O}_2\text{K}) = 3 \text{ mol/l}$.

Adjust to pH 5,2 with glacial acetic acid. Do not autoclave. If necessary, filter through a $0,22 \mu\text{m}$ filter.

A.1.4.5.23 Conditioned glass beads

Incubate glass beads of 0,2 mm to 0,5 mm diameter overnight in concentrated H_2SO_4 (A.1.4.5.3). Wash them with autoclaved water, boil in KHCO_3 solution (A.1.4.5.24), wash again with autoclaved water, and dry at $80 \text{ }^\circ\text{C}$ under vacuum.^{[13], [14]}

A.1.4.5.24 Potassium bicarbonate solution, $\rho(\text{KHCO}_3) = 50 \text{ g/l}$.

Prepare freshly in water.

A.1.4.6 Apparatus and equipment

Usual laboratory equipment and, in particular, the following.

A.1.4.6.1 Cell disruptor, for 2 ml screw-capped polyethylene microtubes, with a beating frequency of at least 100 beats/min [e.g. Mini-BeadBeater™⁴].

A.1.4.6.2 Microtubes, O-ring-secured, screw-capped 2 ml polyethylene tubes.

A.1.4.6.3 Filtering device, for 25 mm diameter-glass fibre filters.

A.1.4.6.4 Centrifuge, capable of holding 2 ml microtubes at a minimum acceleration of 10 000 *g*.

In some steps a refrigerated centrifuge is required.

A.1.4.6.5 Water bath or incubator.

A.1.4.6.6 Vacuum dryer (optional). It is recommended for the preparation of item A.1.4.5.23.

A.1.4.6.7 Mixer, e.g. Vortex®²).

A.1.4.7 Procedure

A.1.4.7.1 Preparation of the test portion and the microbial fraction

Protocols for the isolation of the microbial fraction, occasionally coupled to an enrichment step on agar or in liquid media for food microbiological quality controls, can be exploited for further DNA extraction from the microbial population, being defined as an ingredient.

Starting from 1 ml to 2 ml of the laboratory sample as standard portion, the microbial population is isolated appropriately (see A.1.3). Alternatively, yeasts or filamentous fungi isolated from the test portion may be grown as starter cultures. In both cases, the microorganisms are harvested and further processed according to A.1.4.7.2 or stored at –20 °C until processing is carried out.

A.1.4.7.2 DNA extraction

A.1.4.7.2.1 For mycelia collected on glass fibre filters, wash the mycelial mat twice with lysis buffer (A.1.4.5.18) not containing SDS. Peel off the fungal mat from the filter and transfer it to a 2 ml screw-capped microtube (A.1.4.6.2) containing conditioned glass beads (A.1.4.5.23) (half a tube), 600 µl of lysis buffer (A.1.4.5.18) and 600 µl of phenol-chloroform-isoamyl alcohol (A.1.4.5.17). Further processing is described in A.1.4.7.2.3.

A.1.4.7.2.2 For pellets of either total microbial population, bacteria, mycelium, yeasts or yeast-like microorganisms, wash the cells once with 1 ml of lysis buffer (A.1.4.5.18) not containing SDS, centrifuge at 10 000 *g* to 13 000 *g* for 10 min, repeat at least once then resuspend in 600 µl of lysis buffer (A.1.4.5.18) and transfer to a microtube containing glass beads and phenol-chloroform as in A.1.4.7.2.1.

A.1.4.7.2.3 After step A.1.4.7.2.1 or A.1.4.7.2.2, agitate the microtube at least at 100 beats/min on a cell-disrupter (A.1.4.6.1) for 1 min to 2 min, then incubate immediately at 65 °C for 30 min to 120 min. Centrifuge at 10 000 *g* to 13 000 *g* for 10 min. Transfer the supernatant to a new microtube.

Optionally for further quantitative PCR, after 30 min of incubation, centrifuge at 10 000 *g* to 13 000 *g* for 15 min. Transfer the supernatant into a common microtube, add RNase-A (A.1.4.5.20) at a final concentration of 0,001 mg/ml and incubate further for 30 min to 90 min at 65 °C.

4) Mini-BeadBeater is an example of a suitable product available commercially from Biospec products. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product. Equivalent products may be used if they can be shown to lead to the same results.

Add potassium acetate solution (A.1.4.5.22) at 0,3 mol/l final concentration. Mix, add 1,2 ml of ethanol (A.1.4.5.1) and incubate at –20 °C overnight or for 1 h at –80 °C. Pellet the DNA at 10 000 *g* to 13 000 *g* for 15 min at 4 °C.

Wash carefully the pelleted DNA with ethanol solution (A.1.4.5.21). Drain the microtube upside down on paper and dry the microtube under vacuum. Dissolve the DNA in 50 µl to 100 µl of water. Long-term (up to 5 years) storage at –20 °C is permitted. The use of water instead of TE buffer (A.1.4.5.19) has been validated [13]. This is the DNA master stock.

A.1.4.8 List of examples

The number of species/strains investigated is indicated within brackets:

Absidia corymbifera (1), *Acremonium* spp. (2), *Aspergillus* spp. (119), *Candida* spp. (7), *Cladosporium* spp. (2), *Cryptococcus* spp. (6), *Epidermophyton floccosum* (1), *Fusarium solani* (1), *Malbranchea pulchella* (1), *Geotrichum* spp. (2), *Microsporium canis* (1), *Paecilomyces* spp. (2), *Penicillium* spp. (20), *Pityrosporum ovale* (1), *Rhizopus* spp. (2), *Saccharomyces cerevisiae* (1), *Schizosaccharomyces pombe* (1), *Scopulariopsis brevicaulis* (1), *Trichoderma* spp. (124), *Trichophyton* spp. (2), *Trichosporon* spp. (2), *Ulocladium botrytis* (1), *Verticillium tenerum* (1).

A.1.4.9 Validation

The effectiveness of the method has been validated in the case of fungi^[13]. Quantitative analysis of the extraction efficiency has shown that the use of glass milling was most effective^[18].

A.2 Preparation of PCR-quality DNA using polyvinyl-pyrrolidone (PVP)-based DNA extraction methods

A.2.1 Basic PVP method

A.2.1.1 General

This simple, fast and cheap method^[19] is suitable for a wide range of matrices, in particular the ones containing high amounts of polyphenolic compounds.

A.2.1.2 Validation status

The method has been in-house validated and is used for routine DNA preparations in many laboratories, although it has not yet been evaluated in official interlaboratory studies.

A.2.1.3 Principle

The method consists of a lysis step (thermal lysis in presence of sodium dodecyl sulfate and a high EDTA content), followed by the removal of contaminants such as polyphenolic molecules, polysaccharides, metabolites and soluble proteins, from the DNA-containing liquid phase in combination with PVP and ammonium acetate. A final alcohol-precipitation step allows concentration of the DNA and the elimination of salts, see also References [19], [20], [21], [22] and [23].

A.2.1.4 Safety precautions

A fume hood is necessary for handling organic chemicals.

A.2.1.5 Reagents

A.2.1.5.1 Ethanol, $\phi(\text{C}_2\text{H}_5\text{OH}) = 96 \%$.

Store and use at $-20 \text{ }^\circ\text{C}$.

A.2.1.5.2 Isopropanol ($\text{CH}_3\text{CHOHCH}_3$).

A.2.1.5.3 Polyvinylpyrrolidone (PVP), molecular mass $M = 360\,000 \text{ D}$; intrinsic viscosity (K value) = 80 to 100^5 .

A.2.1.5.4 Glacial acetic acid (CH_3COOH).

A.2.1.5.5 Hydrochloric acid, $\phi(\text{HCl}) = 37 \%$.

A.2.1.5.6 Sodium chloride (NaCl).

A.2.1.5.7 Tris(hydroxymethyl)-aminomethane (Tris) ($\text{C}_4\text{H}_{11}\text{NO}_3$).

A.2.1.5.8 Ethylenediaminetetraacetic acid disodium salt (Na_2EDTA).

A.2.1.5.9 Sodium dodecyl sulfate (SDS) ($\text{C}_{12}\text{H}_{25}\text{O}_4\text{SNa}$).

A.2.1.5.10 Ammonium acetate ($\text{C}_2\text{H}_3\text{O}_2 \text{NH}_4$).

A.2.1.5.11 Ethanol solution, $\phi(\text{C}_2\text{H}_5\text{OH}) = 70 \%$.

Store and use at $-20 \text{ }^\circ\text{C}$.

A.2.1.5.12 Extraction buffer, pH 8,0, $c(\text{Tris}) = 0,2 \text{ mol/l}$, $c(\text{NaCl}) = 0,250 \text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,025 \text{ mol/l}$, $\rho(\text{SDS}) = 50 \text{ g/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.2.1.5.13 Ammonium acetate solution, $c(\text{NH}_4\text{C}_2\text{H}_3\text{O}_2) = 7,5 \text{ mol/l}$.

Dissolve in sterile water and possibly sterilize by filtration through a $0,22 \text{ } \mu\text{m}$ filter.

A.2.1.5.14 TE buffer, $c(\text{Tris}) = 0,010 \text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,001 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.2.1.6 Apparatus and equipment

Usual laboratory equipment and, in particular, the following.

A.2.1.6.1 Centrifuge, capable of achieving an acceleration of $10\,000 \text{ g}$.

In some steps a refrigerated centrifuge is required.

A.2.1.6.2 Water bath or incubator.

5) SIGMA P-5288 is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product. Equivalent products may be used if they can be shown to lead to the same results.

A.2.1.6.3 Vacuum dryer (optional).

A.2.1.6.4 Freeze dryer (optional).

A.2.1.6.5 Mixer, e.g. Vortex^{®2}.

A.2.1.7 Procedure

A.2.1.7.1 General

Once the matrix test portion has been prepared, apply the following DNA extraction/purification protocol. Scale-adaptation of masses and buffer volumes is required as a function of the selected size of the test portion.

A.2.1.7.2 Extraction procedure

Weigh 0,25 g of milled, crushed or liquid material into a vial. Add 1 ml of extraction buffer (A.2.1.5.12). Agitate the suspension at 65 °C for 1 h, cool to room temperature. Mix sequentially the suspension with 60 mg of PVP powder (A.2.1.5.3) and with 0,5 volume of ammonium acetate solution (A.2.1.5.13). Incubate on ice for 30 min.

Centrifuge at 10 000 *g* for 10 min and transfer the supernatant to a fresh tube. Mix the lysate with 1 volume of isopropanol (A.2.1.5.2) and incubate at –20 °C for 30 min. Centrifuge at 10 000 *g* at 4 °C for 10 min and carefully discard the supernatant.

Wash the DNA pellet with 2 volumes of ethanol solution (A.2.1.5.11). This step is essential for the removal of any salts that could interfere with the subsequent analysis (e.g. PCR). Carefully discard the supernatant (in the case of a loose pellet, centrifuge at 10 000 *g* at 4 °C for 10 min). Dry the pellet and redissolve it in 100 µl of water or appropriate buffer, e.g. TE buffer (A.2.1.5.14). This is the DNA master stock.

A.2.1.8 List of examples

The method has been successfully applied to extract DNA⁶⁾ from the following matrices:

baby biscuits⁶⁾, baby milk⁶⁾, Belgian paté, breadcrumbs (wheat or corn) from fish sticks, brownies⁶⁾, canned maize, cereal bars⁶⁾, cheese croquettes, chicken nuggets, chicken, chocolate cookies⁶⁾, chocolate paste⁶⁾, corn flakes⁶⁾, crisps, dessert creams⁶⁾, infant formula, maize biscuits⁶⁾, maize flour, fresh and cooked meat (beef, pork, chicken and turkey), minced meat, muesli⁶⁾, pop corn, powder milk, sausages (slicing⁶⁾ and cocktail⁶⁾), schnitzel, soya sprouts⁶⁾, soup balls, soya protein in meat preparations⁶⁾, soya lecithin⁶⁾, soya drinks⁶⁾, soya bean cream, spaghetti sauces⁶⁾, speculoos, tofu, vegetarian hamburgers, waffles with chocolate⁶⁾, waffles⁶⁾, yoghurts⁶⁾.

A.3 Preparation of PCR-quality DNA using the CTAB-based DNA extraction methods

A.3.1 Basic CTAB method

A.3.1.1 General

The method is applicable to the extraction of DNA from plants and plant-derived matrices, in particular because of its ability to remove polysaccharides and polyphenolic compounds that would otherwise affect the DNA quality. It is also useful for some other matrices (see A.3.1.8).

6) Repeatability may depend on the batch of the matrix and/or to its production technology. In some cases, DNA could not be found or was degraded in such a way that PCR results were below the limit of detection of the method, irrespective of the PCR primers/protocols used. This may be a source of low reproducibility between laboratories.

A.3.1.2 Validation status

This method has been ring tested (see A.3.1.9).

The method is commonly used for routine DNA preparations in many laboratories.

A.3.1.3 Principle

The method consists of a lysis step (thermal lysis in the presence of CTAB), followed by several extraction steps in order to remove contaminants, such as polysaccharides and proteins^[24].

For some matrices, it is helpful to perform different enzymatic steps as outlined in A.3.1.7. Alpha-amylase is added to the lysis buffer to digest the starches in case of amylaceous matrices. Treatment of samples with proteinase-K is necessary in a variety of matrices to eliminate proteins. Also treatment with RNase is usually recommended for those matrices where RNA co-precipitation may disturb the subsequent analytical test.

The salt concentration during the extraction steps is very important for the removal of the contaminants, since a CTAB-nucleic acid precipitate will occur if the salt concentration drops below approximately 0,5 mol/l at room temperature and/or if the temperature drops below 16 °C. By increasing the salt concentration (e.g. addition of sodium chloride), the removal of denaturated proteins and polysaccharides complexed to CTAB is achieved, while the nucleic acids are solubilized. Chloroform is used to further separate the nucleic acids from CTAB and polysaccharide/protein complexes.

Finally, the nucleic acids are purified by isopropanol precipitation and washing with ethanol.

A.3.1.4 Safety precautions

A fume hood is necessary for handling organic chemicals.

A.3.1.5 Reagents

A.3.1.5.1 α -Amylase (optional), type IIa from *Bacillus* species, 1 500 to 3 000 units/mg of protein.

A.3.1.5.2 Chloroform (CHCl₃).

A.3.1.5.3 Ethanol, ϕ (C₂H₅OH) = 96 %.

A.3.1.5.4 Ethylenediaminetetraacetic acid disodium salt (Na₂-EDTA) (C₁₀H₁₄N₂O₈Na₂).

A.3.1.5.5 Hexadecyl-trimethyl-ammonium-bromide (CTAB) (C₁₉H₄₂BrN).

A.3.1.5.6 Hydrochloric acid, ϕ (HCl) = 37 %.

A.3.1.5.7 Isopropanol [CH₃CH(OH)CH₃].

A.3.1.5.8 Proteinase-K (optional), approximately 20 Units/mg of lyophilisate.

A.3.1.5.9 RNase A, DNase-free, (optional) from bovine pancreas, approximately 50 Units/mg of lyophilisate.

A.3.1.5.10 Sodium chloride (NaCl).

A.3.1.5.11 Sodium hydroxide (NaOH).

A.3.1.5.12 Tris(hydroxymethyl)-aminomethane (Tris) (C₄H₁₁NO₃).

A.3.1.5.13 α -Amylase solution (optional), $c(\alpha\text{-amylase}) = 10 \text{ mg/ml}$.

Do not autoclave. Store at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.3.1.5.14 CTAB extraction buffer, $\rho(\text{CTAB}) = 20 \text{ g/l}$, $c(\text{NaCl}) = 1,4 \text{ mol/l}$, $c(\text{Tris}) = 0,1 \text{ mol/l}$,
 $c(\text{Na}_2\text{EDTA}) = 0,02 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.3.1.5.15 CTAB-precipitation buffer, $\rho(\text{CTAB}) = 5\text{g/l}$, $c(\text{NaCl}) = 0,04 \text{ mol/l}$.

A.3.1.5.16 Sodium chloride solution, $c(\text{NaCl}) = 1,2 \text{ mol/l}$.

A.3.1.5.17 Ethanol solution, $\phi(\text{C}_2\text{H}_5\text{OH}) = 70 \%$.

A.3.1.5.18 Proteinase-K solution (optional), $\rho = 20 \text{ mg/ml}$, dissolved in sterile water.

Do not autoclave. Store at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.3.1.5.19 RNase-A solution (optional), $\rho(\text{RNase A}) = 10 \text{ mg/ml}$.

Store in aliquots at $-20 \text{ }^\circ\text{C}$.

A.3.1.5.20 TE buffer, $c(\text{Tris}) = 0,01 \text{ mol/l}$, $c(\text{Na}_2\text{-EDTA}) = 0,001 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.3.1.6 Apparatus and equipment

Usual laboratory equipment and, in particular, the following.

A.3.1.6.1 Oven or incubator, preferably with a shaker.

A.3.1.6.2 Centrifuge, e.g. microcentrifuge, capable of achieving an acceleration of up to $12\,000 g$.

In some steps a refrigerated centrifuge is required.

A.3.1.6.3 Mixer, e.g. Vortex[®] 2).

A.3.1.6.4 Vacuum dryer (optional).

A.3.1.7 Procedure

A.3.1.7.1 General

Once the matrix test portion has been prepared, apply the following DNA extraction/purification protocol. Scale-adaptation of masses and buffer volumes is required as a function of the selected size of the test portion.

A.3.1.7.2 Sample extraction

Weigh 200 mg to 300 mg of the test sample into a tube.

Add 1,5 ml of pre-warmed ($65 \text{ }^\circ\text{C}$) CTAB extraction buffer (A.3.1.5.14) and mix. (In some cases a higher amount of buffer may be required to suspend the matrix.) Add $10 \mu\text{l}$ of α -amylase solution (A.3.1.5.13, optional), $10 \mu\text{l}$ of RNase A solution (A.3.1.5.19, optional) and mix gently. Incubate for 30 min at $65 \text{ }^\circ\text{C}$, under

agitation. Add 10 µl of proteinase-K solution (A.3.1.5.18, optional), smoothly mix the tubes and incubate for 30 min at 65 °C, under agitation (optional). Centrifuge for 10 min at approximately 12 000 g. Transfer the supernatant to a new tube, add 0,7 to 1 volume of chloroform (A.3.1.5.2) and mix thoroughly.

Centrifuge for 15 min at approximately 12 000 g. Transfer the upper phase (aqueous) to a new tube.

A.3.1.7.3 CTAB-precipitation

Add 2 volumes of the CTAB precipitation buffer (A.3.1.5.15). Incubate for 60 min at room temperature without agitation. Centrifuge for 15 min at 12 000 g. Discard the supernatant. Dissolve the precipitated DNA by adding 350 µl of NaCl solution (A.3.1.5.16). Add 350 µl of chloroform (A.3.1.5.2) and mix thoroughly. Centrifuge for 10 min at 12 000 g. Transfer the aqueous phase into a new tube.

NOTE CTAB-precipitation is not necessary for all matrices, only for protein- and polysaccharide-rich matrices. Alternatively, a solid-phase purification of the DNA (e.g. by the use of spin columns) is possible assuming the results are equivalent.

A.3.1.7.4 DNA precipitation

Add 0,6 volume of isopropanol (A.3.1.5.7), mix smoothly by inverting the tube and keep the tube at room temperature for 20 min. Centrifuge for 15 min at 12 000 g. Discard the supernatant. Add 500 µl of ethanol solution (A.3.1.5.17) to the tube and invert several times. This is the critical step ensuring the complete removal of CTAB. Centrifuge for 10 min at 12 000 g. Discard the supernatant. Dry the DNA pellet and redissolve it into 100 µl of water or an appropriate buffer, e.g. TE buffer (A.3.1.5.20). This is the DNA master stock.

A.3.1.8 List of examples

The method has been successfully applied to extract DNA⁷⁾ from the following matrices:

baby food (powder), baby food, baking mix, biscuits, bouillon cubes⁷⁾, sweet and sour candies, canned corn, caramel cream⁷⁾, cattle cakes, cereal grains (rice, wheat, oat, rye, buckwheat, millet), chocolate bars⁷⁾, chocolate cream⁷⁾, chocolates⁷⁾, chocolate cookies⁷⁾, cookies, corn beer⁷⁾, cornflakes⁷⁾, dessert cream, dextrose⁷⁾, fillings of pralines, fine pastries, fish⁷⁾, fish fingers⁷⁾, flakes of whole soya bean, French fries, gravy⁷⁾, boiled ham, honey⁷⁾, instant meals, maize ear, maize flour, maize germs⁷⁾, maize gluten feed, maize grids, maize leaves, maize native starch⁷⁾, maize oil (native)⁷⁾, maize proteins⁷⁾, maize seeds/grains, maize semolina, margarine⁷⁾, fresh meat, milk powder, milk, mixed pet food, muesli⁷⁾, mungbean seeds, mustard leaves, popcorn (raw), potato chips, potato starch (native), potato tubers, rape leaves, rape press cake, rape seed oil (crude/native)⁷⁾, rape seed seeds, raw soya lecithin⁷⁾, ready-meals, salami (high fat content), salty snacks (maize), sausages, seasoning agents⁷⁾, modified starches (some types of)⁷⁾, sour cream with onion⁷⁾, soya flour, soya germs (preserved, frozen), soya protein⁷⁾, soya drinks⁷⁾, soya bean seeds/grains, soya bean tofu, soya beans (acidified)⁷⁾, sugar-beet leaves, sugar-beet seeds, sunflower seeds, surimi with soya⁷⁾, sweet corn, taco shells, taramas (paste of fish roes), tobacco, tomato ketchup⁷⁾, tomato-concentrate⁷⁾, tomatoes (fruits), tortilla chips⁷⁾, vegetarian hamburgers, waffles⁷⁾, wheat starch (native), yoghurts⁷⁾.

A.3.1.9 Validation

The validation data in Table A.5 have been elaborated in collaborative studies carried out by the working group "Development of methods for identifying foodstuffs produced by genetic engineering techniques" of the Commission of the former German Federal Health Board for the implementation of methods according to article 35 of the German Foodstuffs Act (see References [25], [26], [27]). The matrices tested were potatoes, soya beans and tomatoes. In these interlaboratory studies, the procedure was carried out with 100 mg of sample material. The CTAB precipitation step was necessary for the analysis of soya bean and soya flour. The enzymatic steps were not carried out in these interlaboratory studies.

7) Repeatability may depend on the batch of the matrix and/or to its production technology. In some cases, DNA could not be found or was degraded in such a way that PCR results were below the limit of detection of the method, irrespective of the PCR primers/protocols used. This may be a source of low reproducibility between laboratories.

For the collaborative study on soya beans, two participants used strongly modified protocols and in one laboratory the testing of the five samples was discontinued. So 22 out of 25 participants identified all 110 samples correctly.

For the collaborative study on potatoes, three samples were identified false negative and one sample false positive. Three samples were not assessed due to ambiguous results between both replicates.

Table A.5 — Validation data

Matrix	Number of laboratories participating	Number of samples per laboratory	Number of total samples	Number of samples correctly identified
Soya beans ^[25]	25	5	125	110
Potatoes ^[26]	18	10	180	173
Tomatoes ^[27]	18	5	90	90

A.4 Preparation of PCR quality DNA using the silica-based DNA extraction methods

A.4.1 Basic silica method

A.4.1.1 General

The method is suitable for DNA extraction from a wide range of matrices (see examples in A.4.1.8). This method can also be employed as a further purification step of DNA solutions obtained after DNA extraction with other methods.

The method is adapted from a published protocol^[28]. It has some advantages for the matrices for which it is suitable, in particular for the avoidance of heavily toxic reagents. Furthermore, the procedure can be easily adapted for manual high-throughput analyses and for automation, in particular because of the lack of labile interfaces (such as water-chloroform) and because low-speed centrifugation is required.

The method is usually not recommended to extract DNA from fat-rich matrices.

A.4.1.2 Validation status

The method has been in-house validated and is used for routine DNA preparations in many laboratories, although it has not yet been evaluated by official interlaboratory studies. The principle of this method has also been implemented in various kit-based methods that were successfully ring tested (see References [29], [30] and [31]).

A.4.1.3 Principle

The method consists of a lysis step (thermal lysis in the presence of sodium dodecyl sulfate in a buffered solution), followed by a purification step realised by means of a silica resin, in the presence of the chaotropic reagent guanidine-hydrochloride. The principle of the method is the binding of nucleic acids to silica under low water activity due to an entropic effect^[32]. Contaminants are washed from the resin by isopropanol, while the DNA stays attached. A final elution step with a low-salt buffer solution permits DNA recovery.

Users of this International Standard are made aware that silica-based methods may be covered by patent rights^[33].

A.4.1.4 Safety precautions

A fume hood is necessary for handling organic chemicals.

A.4.1.5 Reagents

A.4.1.5.1 Sodium chloride (NaCl).

A.4.1.5.2 Tris(hydroxymethyl)-aminomethane (Tris) (C₄H₁₁NO₃).

A.4.1.5.3 Ethylenediaminetetraacetic acid-disodium salt (Na₂EDTA) (C₁₀H₁₄N₂O₈Na₂).

A.4.1.5.4 Hydrochloric acid, $\phi(\text{HCl}) = 37 \%$.

A.4.1.5.5 Sodium hydroxide (NaOH).

A.4.1.5.6 Sodium dodecyl sulfate (SDS) (C₁₂H₂₅O₄SNa).

A.4.1.5.7 Proteinase-K, approximately 20 U/mg of lyophilisate.

A.4.1.5.8 Guanidine hydrochloride (CH₅N₃-HCl).

A.4.1.5.9 Potassium chloride (KCl).

A.4.1.5.10 Disodium hydrogen phosphate (Na₂HPO₄).

A.4.1.5.11 Potassium dihydrogen phosphate (KH₂PO₄).

A.4.1.5.12 Isopropanol [CH₃CH(OH)CH₃].

A.4.1.5.13 Silica (SiO₂), silicon dioxide with particle size distribution between 0,5 and 10 μm (80 % between 1 μm to 5 μm)⁸.

A.4.1.5.14 RNase-A, DNase-free, approximately 100 Kunitz Units/mg of lyophilisate.

A.4.1.5.15 Proteinase-K solution, $\rho = 20 \text{ mg/ml}$.

Dissolve the enzyme in autoclaved water or buffer as described in Reference [34]. Do not autoclave this solution. Store in aliquots at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.4.1.5.16 Guanidine hydrochloride solution I, $c(\text{CH}_5\text{N}_3\text{-HCl}) = 5 \text{ mol/l}$.

Autoclave for a maximum of 15 min at $121 \text{ }^\circ\text{C}$.

A.4.1.5.17 Guanidine-HCl solution II, $c(\text{CH}_5\text{N}_3\text{-HCl}) = 6 \text{ mol/l}$.

Autoclave for a maximum of 15 min at $121 \text{ }^\circ\text{C}$.

A.4.1.5.18 PBS-buffer solution, $c(\text{NaCl}) = 0,157 \text{ mol/l}$, $c(\text{KCl}) = 0,0027 \text{ mol/l}$, $c(\text{Na}_2\text{HPO}_4) = 0,010 \text{ mol/l}$, $c(\text{KH}_2\text{PO}_4) = 0,0018 \text{ mol/l}$.

Adjust the pH to 7,5 with HCl.

8) SIGMA S-5631 is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product. Equivalent products may be used if they can be shown to lead to the same results.

A.4.1.5.19 Silica suspension

Weigh 5 g of silica (A.4.1.5.13) into a 50 ml tube and add 50 ml of PBS-buffer (A.4.1.5.18). Mix well and allow to settle for 2 h. Remove the supernatant by aspiration with a pipette. Add another 50 ml of the PBS-buffer, mix well and allow to settle again for 2 h. Remove the supernatant by aspiration. Centrifuge for 2 min at 2 000 g. Discard the remaining supernatant. Resuspend the pellet up to 50 ml in guanidine-HCl solution II (A.4.1.5.17). Use within 2 to 5 months. Mix well before use.

A.4.1.5.20 TNE-SDS extraction buffer, $c(\text{NaCl}) = 0,150 \text{ mol/l}$, $c(\text{Tris}) = 0,002 \text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,002 \text{ mol/l}$, $\rho(\text{SDS}) = 10\text{g/l}$.

Adjust the pH to 8,0 with HCl or NaOH and autoclave before adding the SDS.

A.4.1.5.21 Isopropanol solution, $\phi[\text{CH}_3\text{CH}(\text{OH})\text{CH}_3] = 80 \%$.

A.4.1.5.22 TE-buffer solution, $c(\text{Tris}) = 0,010 \text{ mol/l}$ and $c(\text{Na}_2\text{EDTA}) = 0,001 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.4.1.5.23 RNase-A solution, $\rho(\text{RNase}) = 10 \text{ mg/ml}$.

Store in aliquots at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.4.1.6 Apparatus and equipment

Usual laboratory equipment and, in particular, the following.

A.4.1.6.1 Centrifuge, capable of achieving an acceleration of at least 2 000 g.

In some steps a refrigerated centrifuge is required.

A.4.1.6.2 Oven or incubator, with a working temperature of $60 \text{ }^\circ\text{C}$.

A.4.1.6.3 Shaker, to be put inside the oven/incubator.

A.4.1.6.4 Mixer, e.g. Vortex^{®2}.

A.4.1.6.5 Centrifugation vials, 50 ml, for preparation of the silica suspension.

A.4.1.7 Procedure

A.4.1.7.1 General

Once the matrix test portion has been prepared, apply the following DNA extraction/purification protocol. Scale-adaptation of masses and buffer volumes is required as a function of the selected size of the test portion.

A.4.1.7.2 Extraction procedure

Weigh 200 mg to 300 mg of milled or crushed material into a vial. Add 2 ml of extraction buffer (A.4.1.5.20) and 20 μl of Proteinase-K solution (A.4.1.5.15). Incubate for 1 h to 5 h in the oven set at $60 \text{ }^\circ\text{C}$. During the incubation time, shake the samples vigorously (approximately 250 min^{-1}). Centrifuge for 15 min at 2 000 g. Transfer 550 μl of the supernatant to a new tube.

Treat the transferred supernatant with 2 µl of RNase solution (A.4.1.5.23) for 5 min at 37 °C (this hydrolysing RNA step is recommended before the silica binding, otherwise the hydrolysed RNA and the resulting nucleotides may interfere with subsequent UV-spectrometry measurements). Add 55 µl of the guanidine-HCl solution I (A.4.1.5.16) plus 100 µl of the silica suspension (A.4.1.5.19) to the supernatant. Mix carefully several times. Leave the tubes on the bench for approximately 1 min.

Centrifuge for 2 min at approximately 800 *g*. Discard the supernatant and add 500 µl of the isopropanol solution (A.4.1.5.21). Close the tubes and mix, possibly with the aid of the mixer (A.4.1.6.4), to completely resuspend the pellet.

Centrifuge for 2 min at approximately 1 500 *g*. Discard the supernatant and dry the pellet. Add 100 µl of TE-buffer solution (A.4.1.5.22). Mix carefully in order to resuspend the pellet. Incubate the samples at 60 °C for 5 min. Centrifuge for 5 min at 2 000 *g*. Transfer 80 % of the supernatant to a new tube. Be careful not to transfer any silica particles because of their inhibiting activity on enzymes (e.g. DNA polymerase, restriction endonucleases).

Treat the transferred supernatant with 2 µl of RNase solution (A.4.1.5.23) for 1 h at 37 °C, or overnight at room temperature. This is the DNA master stock.

A.4.1.8 List of examples

The method has been successfully applied to extract DNA⁹⁾ from the following matrices:

germs from maize⁹⁾, maize flour, maize gluten feed⁹⁾, maize leaves, maize-modified starches⁹⁾, maize native starch⁹⁾, maize seeds, maize semolina, protein from soya beans⁹⁾, soya beans, soya bean leaves, sugar beet (fresh root), sugar beet (frozen pulp), sugar beet leaves.

A.5 Preparation of PCR quality DNA using guanidinium–chloroform-based DNA extraction methods

A.5.1 Basic guanidine-chloroform method

A.5.1.1 General

This method is suitable for DNA extraction from a wide range of food and feed matrices (see A.5.1.8). A further purification step may be necessary depending on the sample.

A.5.1.2 Validation status

The method has been validated in-house.

A.5.1.3 Principle

The method consists of a thermal and enzymatic lysis in presence of sodium dodecyl sulfate in a denaturing guanidinium buffer, occasionally followed by a purification step depending on the matrix.

Contaminants such as lipids and proteins are eliminated by a chloroform-extraction step after the lysis, then DNA is precipitated by isopropanol before purification.

A.5.1.4 Safety precautions

A fume hood is necessary for manipulation of organic chemicals.

9) Repeatability may depend on the batch of the matrix and/or to its production technology. In some cases, DNA could not be found or was degraded in such a way that PCR results were below the limit of detection of the method, irrespective of the PCR primers/protocols used. This may be a source of low reproducibility between laboratories.

A.5.1.5 Reagents

A.5.1.5.1 α -Amylase, type IIa from *Bacillus* spp., 1 500 to 5 000 Units/mg of lyophilisate.

A.5.1.5.2 Glacial acetic acid (CH_3COOH).

A.5.1.5.3 Chloroform (CHCl_3).

A.5.1.5.4 Ethanol, $\phi(\text{C}_2\text{H}_5\text{OH}) = 96 \%$.

A.5.1.5.5 Ethylenediaminetetraacetic acid disodium salt (Na_2EDTA) ($\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_8\text{Na}_2$).

A.5.1.5.6 Guanidine hydrochloride ($\text{CH}_5\text{N}_3\text{-HCl}$).

A.5.1.5.7 Hydrochloric acid, $\phi(\text{HCl}) = 37 \%$.

A.5.1.5.8 Isopropanol [$\text{CH}_3\text{CH}(\text{OH})\text{CH}_3$].

A.5.1.5.9 Proteinase-K, approximately 20 Units/mg lyophilisate.

A.5.1.5.10 RNase-A, DNase-free, from bovine pancreas, approximately 50 Kunitz Units/mg of lyophilisate.

A.5.1.5.11 Sodium chloride (NaCl).

A.5.1.5.12 Sodium dodecyl sulfate (SDS) ($\text{C}_{12}\text{H}_{25}\text{O}_4\text{SNa}$).

A.5.1.5.13 Sodium hydroxide (NaOH).

A.5.1.5.14 Tris(hydroxymethyl)-aminomethane (Tris) ($\text{C}_4\text{H}_{11}\text{NO}_3$),

A.5.1.5.15 α -Amylase solution, $\rho = 10 \text{ mg/ml}$, dissolved in sterile water.

Do not autoclave. Store at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.5.1.5.16 Ethanol solution, $\phi(\text{C}_2\text{H}_5\text{OH}) = 70 \%$.

A.5.1.5.17 Extraction buffer, $c(\text{Tris}) = 0,1 \text{ mol/l}$, $c(\text{NaCl}) = 0,15 \text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,05 \text{ mol/l}$,
 $\rho(\text{SDS}) = 10\text{g/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.5.1.5.18 Guanidine hydrochloride solution, $c(\text{CH}_5\text{N}_3\text{-HCl}) = 5 \text{ mol/l}$.

Autoclave after preparation (maximum 15 min at $121 \text{ }^\circ\text{C}$).

A.5.1.5.19 Proteinase-K solution, $\rho = 20 \text{ mg/ml}$, dissolved in sterile water.

Do not autoclave. Store at $-20 \text{ }^\circ\text{C}$, but avoid repeated freezing and thawing.

A.5.1.5.20 RNase-A solution, $\rho = 10 \text{ mg/ml}$.

Store in aliquots at $-20 \text{ }^\circ\text{C}$.

A.5.1.5.21 TE buffer, $c(\text{Tris}) = 0,01 \text{ mol/l}$, $c(\text{Na}_2\text{EDTA}) = 0,001 \text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH.

A.5.1.6 Apparatus and equipment

A.5.1.6.1 Centrifuge, capable of achieving an acceleration of 8 000 *g*. In some steps a refrigerated centrifuge is required.

A.5.1.6.2 Oven or incubator, preferably with a shaker.

A.5.1.6.3 Mixer, e.g. Vortex^{®2}).

A.5.1.7 Procedure

A.5.1.7.1 General

Once the matrix test portion has been prepared, apply the following DNA extraction/purification protocol. Scale-adaptation of masses and buffer volumes is required as a function of the selected size of the test portion.

A.5.1.7.2 Extraction procedure

Weigh 200 mg to 500 mg of milled or crushed material into a microtube. Add 1,6 ml of prewarmed (60 °C) extraction buffer (A.5.1.5.17).

Add 10 µl of the RNase-A solution (A.5.1.5.20) and 10 µl of the α -amylase solution (A.5.1.5.15), and mix gently by inverting manually the tube. Incubate for 50 min at 60 °C under gentle agitation. Add 1/10 volume of guanidine-hydrochloride solution (A.5.1.5.18) solution, then mix with the mixer (A.5.1.6.3).

Add 20 µl of the Proteinase-K solution (A.5.1.5.19), mix smoothly by inverting the tube manually and incubate for at least 2 h at 60 °C under gentle agitation. Leave the tubes on the bench for 15 min then centrifuge for 15 min at a minimum acceleration of 8 000 *g*.

Transfer the supernatant into a new tube. Add 1 volume of chloroform (A.5.1.5.3) and mix with a mixer (A.5.1.6.3). Centrifuge for 15 min at a minimum acceleration of 8 000 *g*. Transfer the supernatant to a new tube. Add 0,6 volumes of isopropanol (A.5.1.5.8), mix inverting and put the tubes on ice for 50 min.

Centrifuge for 20 min at a minimum acceleration of 8 000 *g*. Wash the pellet with at least 2 ml of ethanol solution (A.5.1.5.16) and centrifuge for 10 min at a minimum acceleration of 8 000 *g*. This step is essential for the removal of any salts that could interfere with the subsequent analysis (e.g. PCR). Discard the supernatant. Dry the pellet and redissolve it in 100 µl of water or appropriate buffer, e.g. TE buffer (A.5.1.5.21). This is the DNA master stock. If a further purification step is required, it shall be performed on the DNA master stock.

A.5.1.8 List of examples

The method has been successfully applied to extract DNA¹⁰⁾ from the following matrices:

acidified soya bean, canned corn, cattle cakes, chocolate bars¹⁰⁾, dessert cream¹⁰⁾, flakes of whole soya bean, maize ears, maize flour, maize germs¹⁰⁾, maize gluten feed¹⁰⁾, maize grids, maize modified starches¹⁰⁾, maize proteins¹⁰⁾, maize seeds/grains, maize semolina, maize native starches¹⁰⁾, rapeseed seeds/grains, sauces¹⁰⁾, soya flour, soya bean tofu, soya lecithin (raw brown¹⁰⁾ and refined yellow¹⁰⁾), soya protein, soya seeds/grains, soya tonyu, tortilla chips¹⁰⁾.

The method has not yet been successfully applied even on a 1 g test sample size of oils, maltodextrine, D-glucose, maltitol, mannitol or xylitol.

10) Repeatability may depend on the batch of the matrix and/or to its production technology. In some cases, DNA could not be found or was degraded in such a way that PCR results were below the limit of detection of the method, irrespective of the PCR primers/protocols used. This may be a source of low reproducibility between laboratories.

A.5.2 Guanidine chloroform method: Protocol for soybean lecithin

A.5.2.1 Purpose, relevance and scientific basis

Soybean lecithin is a frequent ingredient and is used in many food products as an emulsifier. Soybean lecithin can be either produced using genetically modified (GM) or non-modified soybeans. The method described here can be used to extract DNA present in the sample in order to perform subsequent PCR analyses for detection of genetically modified DNA sequences derived from GM soybeans.

The method is based on Reference [44], and a very similar procedure has been validated in a collaborative trial in Switzerland. In that study, for the purpose of interpretation, the quantity of extracted DNA was spectrometrically determined (Reference [35], section 1.3). The Chemical and Veterinary Institute Freiburg, Germany, conducted an additional collaborative trial with 12 participating laboratories where the amount of extractable and amplifiable soybean DNA was determined by means of quantitative real-time PCR. For the results, see Reference [45].

A.5.2.2 Scope

This method describes a procedure for DNA extraction from soybean lecithins in raw and cold-pressed vegetable oils, respectively. If the DNA content of the sample material is low, the real-time PCR approach is used for quantitation of the amount of DNA isolated from a test portion and for the calculation of the practical limit of detection achievable in PCR analysis of the extracted DNA, respectively.

A.5.2.3 Validation status and performance criteria

A.5.2.3.1 Validation criteria

The method described in this subclause has been validated in a collaborative trial by determining the quantity of extractable, amplifiable soybean DNA by means of quantitative real-time PCR. The collaborative study was performed in accordance with the IUPAC protocol (Reference [46]).

A.5.2.3.2 Robustness of the method

The method has been routinely used in enforcement and private GMO testing laboratories in Germany and Switzerland for more than 10 years and no problems have been reported. Although specific robustness data (e.g. by modifying method parameters) are not available, the experiences of the laboratories showed that small variations in conditions does not interfere with the performance of the method.

A.5.2.3.3 Intralaboratory trial

The materials used in the collaborative trial study were tested in the method developers' laboratory regarding the intralaboratory precision of the method.

For estimation of the precision, five DNA extractions from each of five soybean lecithins were prepared and measured by real-time PCR under repeatability conditions using a soybean lectin gene specific method according to ISO 21570:2005,^[41] C.2. The results are given in Table A.6. **A1**

Table A.6 — Intralaboratory validation of the DNA extraction method with five soybean lecithin samples ($n = 5$ extractions each)

Lecithin sample No.	Mean lectin copy No.	Repeatability standard deviation, s_r copy No.	Coefficient of variation of repeatability, $C_{V,r}$ %
1	20 464	5 367	26
2	3 203	620	19
3	2 005	306	15
4	6 978	331	5
5	14	8	61

Prior to the collaborative trial, DNA was extracted from five commercial soybean lecithins and tested for PCR inhibition. The DNA prepared from each sample was further diluted using TE buffer (1 volume DNA diluted with 4 volumes of TE; 1 volume of first DNA dilution diluted with 4 volumes of TE, 1 volume of second DNA dilution diluted with 4 volumes of TE). The difference between the calculated (extrapolated) C_T -value of the undiluted DNA sample and the measured C_T -value of each DNA dilution was below 0,1, showing that DNA preparations were free of PCR-inhibitors.

A.5.2.3.4 Collaborative trial

A collaborative trial (validation study) was carried out in 12 laboratories (Reference [45]). Five commercial soybean lecithin samples were used. Each laboratory received 15 coded samples and a standard DNA for calibration. The samples were distributed in a way that each participant received three identical samples of each of the five soybean lecithins. A single DNA extraction was performed per sample by each laboratory. The returned results were assigned to the five different lecithin samples for evaluation of the collaborative trial.

The extracted DNAs were tested in a real-time PCR amplifying a DNA sequence of the soybean lectin gene according to the method in ISO 21570:2005,[41] C.2. Standard-DNA for calibration was prepared from soybean flour [ERM BF410a¹⁾] by using the Plant Mini Kit³⁾ (Qiagen, Hilden/Germany) starting with a CTAB extraction (see A.3). The concentration of DNA was estimated fluorimetrically with the PicoGreen³⁾ dsDNA method (Reference [17]). DNA-standards were defined by the copy number (cp) of haploid genome equivalents per microlitre. For the calculation, a haploid genome mass for soybean of 1,13 pg was assumed. A dilution series was prepared ranging from about 50 000 cp/5 μ l to 80 cp/5 μ l. Seven laboratories used ABI³⁾ real-time PCR equipment (ABI 7000, 7500, 7700), and five laboratories used the Light Cycler³⁾ (Roche) with the following modifications to the protocol described in ISO 21570:2005,[41] C.2: PCR was performed in a 20 μ l final volume containing QuantiTect Probe PCR Master Mix³⁾ (Qiagen), primers GM1-F and GM1-R at 500 nmol/l each and 150 nmol/l of probe GM1. Real-time PCR program was: initial step with 900 s at 95 °C, then 45 cycles with 10 s at 95 °C, 30 s at 60 °C and 30 s at 72 °C. Fluorescence signal acquisition was done within the elongation step. The temperature ramping rate was set to 2°C/s.

1) Product available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement of this product by ISO. Equivalent products may be used if they can be shown to lead to the same results.

A1) As a criterion for the suitability of the method, the practical limit of detection, LOD_{prac} , for genetically modified soya was determined in accordance with ISO 24276. The value was calculated individually for each sample using Formula (A.1):

$$LOD_{\text{prac}} \% = \frac{LOD_{\text{abs}}}{c_{s, \text{DNA}}} \times 100 \quad (\text{A.1})$$

where

LOD_{abs} is the LOD of the event-specific real-time PCR method used for quantitation, in copies per PCR;

NOTE For the calculations in Table A.7, an LOD of 10 copies is assumed.

$c_{s, \text{DNA}}$ is the amount of amplifiable soybean DNA in the sample, in copies per PCR, determined by means of real-time PCR method specific for a soybean reference gene.

Table A.7 summarizes the results of the collaborative trial. In four out of five lecithin samples, mean values of LOD_{prac} below 0,9 % (example for an existing legislative threshold) were obtained. For lecithin sample 2, the LOD_{prac} was above 0,9 % in only one laboratory, for lecithin sample 3 in two laboratories. For the lecithin sample 5, all laboratories amplified fewer than 80 copies of the lectin gene, and thus an LOD_{prac} of 0,9 % or less could not be achieved. In addition, the coefficient of variation of reproducibility for the lectin copy numbers reported for the five lecithin samples were calculated. In total, real-time PCR data of 36 extractions per lecithin sample were used for the evaluation. No outliers were eliminated for the calculations of the coefficient of variation of reproducibility, $C_{V, R}$. Precision data for sample 5 cannot be given because of the low amount of lectin copies below the LOQ of the method. It is noted, that the precision data reflect the coefficient of variation of reproducibility of the copy numbers extracted from the lecithin samples under reproducibility conditions in the collaborative trial.

Table A.7 — Summary of the validation data

Lecithin sample No.	Mean lectin copy No.	Coefficient of variation of reproducibility, $C_{V, R}$ %	Mean practical limit of detection, LOD_{prac} %	No. laboratories with $LOD_{\text{prac}} < 0,9$ %
1	17 044	67	0,06	12/12
2	3 630	63	0,28	11/12
3	2 318	57	0,43	10/12
4	8 325	52	0,12	12/12
5	<80	not determined	>10	0/12

A1)

A₁ A.5.2.4 Principle and summary

Viscous sample material is homogenized after heating and extracted with hexane after addition of a guanidine thiocyanate buffer. Interfering accompanying substances are separated by extraction with chloroform, and the RNA present in the solution is digested with RNase A. The DNA is precipitated with isopropanol in the presence of glycogen, then washed with ethanol and dissolved in water. For purification of the DNA, a subsequent gel filtration step (using a cross-linked dextran gel for size exclusion chromatography) is carried out.

A.5.2.5 Terms and definitions

For the purposes of this subclause, the terms and definitions given in ISO 5725-1^[43] and ISO 24276 apply.

A.5.2.6 Sample type and amounts

Ensure that the test sample is representative of the laboratory sample. Measures and operational steps to be taken into consideration are described in 5.1. Sample material should be as homogeneous as possible.

A.5.2.7 Estimation of measurement uncertainty

The measurement uncertainty of the lectin real-time PCR data assessed in the collaborative trial is given as coefficient of variation of reproducibility. The results are given in Table A.7.

A.5.2.8 Interferences

The degree of refinement of the lecithin can interfere with the ability to extract DNA from lecithin samples due to the DNA degradation accompanied by this processing.

A.5.2.9 Physical or environmental conditions

No specific conditions required. See ISO 24276 for details.

A.5.2.10 Apparatus

Usual laboratory equipment and in particular the following.

A.5.2.10.1 Table-top centrifuge for 1,5 ml or 2 ml centrifugation vessels, at least 12 000 × *g*.

A.5.2.10.2 Centrifuge for 50 ml centrifuge tube, at least 4 000 × *g*.

A.5.2.10.3 Polypropylene centrifugation vessels of capacities 1,5 ml, 2,0 ml, and 50 ml, for use in centrifuges at 12 000 × *g* and 4 000 × *g*.

A.5.2.10.4 Heating block with shaking device.²⁾

A.5.2.10.5 Vacuum drier, optional.

A.5.2.10.6 Multi mixer, e.g. Vortex.

2) The Eppendorf Thermomixer is an example of a suitable product available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of this product. **A₁**

A.5.2.10.7 Real-time PCR thermal cycler equipped with an energy source suitable for the excitation of fluorescent molecules and an optical detection system suitable for the detection of the fluorescence signals generated during PCR.

A.5.2.10.8 Reaction vessels and caps or closures which can be repeatedly heated to 100 °C and cooled to 4 °C without damage and which do not influence the fluorescence signal generated during the amplification.

A.5.2.10.9 UV-spectrophotometer or fluorometer to determine the concentration of DNA.

A.5.2.11 Reagents and materials

For the quality of the reagents used, see ISO 24276.

A.5.2.11.1 Guanidine thiocyanate.

A.5.2.11.2 *Tris*(hydroxymethyl)aminomethane (tris) or ***tris*(hydroxymethyl)aminomethane hydrochloride** (tris·HCl).

A.5.2.11.3 Ethylenediaminetetraacetic acid disodium salt (Na₂EDTA).

A.5.2.11.4 Pancreatic RNase A.

A.5.2.11.5 Glycogen.

A.5.2.11.6 Polyethyleneglycol tert-octylphenyl ether [4-(1,1,3,3-tetramethylbutyl)phenylpolyethyleneglycol] [Triton X 100³].

A.5.2.11.7 *n*-Hexane.

A.5.2.11.8 Chloroform.

A.5.2.11.9 Sodium chloride.

A.5.2.11.10 Sodium hydroxide.

A.5.2.11.11 Hydrochloric acid, $c(\text{HCl}) = 0,1 \text{ mol/l}$.

A.5.2.11.12 Sodium hydroxide solution, $c(\text{NaOH}) = 0,1 \text{ mol/l}$. Weigh 10 g of sodium hydroxide into a 250 ml volumetric flask, and fill the flask with water to the calibration mark.

A.5.2.11.13 Isopropanol (2-propanol).

A.5.2.11.14 Ethanol, $\varphi(\text{C}_2\text{H}_5\text{OH}) = 70 \text{ \%}$ volume fraction. Mix 70 ml of ethanol with 30 ml of water. **A1**

A.5.2.11.15 Tris·HCl buffer, pH = 6,4, $c(\text{tris}\cdot\text{HCl}) = 0,1 \text{ mol/l}$. Dissolve 1,57 g of tris·HCl in approximately 70 ml of water; adjust to pH 6,4 using hydrochloric acid, transfer to a 100 ml volumetric flask, and make up to volume with water.

A.5.2.11.16 Tris·HCl + NaCl buffer, pH 7,5, $c(\text{tris}\cdot\text{HCl}) = 10 \text{ mmol/l}$, $c(\text{NaCl}) = 15 \text{ mmol/l}$. Dissolve 0,166 g of tris·HCl and 88 mg of sodium chloride in approximately 70 ml of water; adjust to pH 7,5 using hydrochloric acid, transfer to a 100 ml volumetric flask, and make up to volume with water.

A.5.2.11.17 EDTA solution, pH 8,0, $c = 0,5 \text{ mol/l}$. Dissolve 18,6 g of EDTA disodium salt in approximately 70 ml of water; adjust to pH 8,0 using first sodium hydroxide pellets and then sodium hydroxide solution, transfer to a 100 ml volumetric flask, and make up to volume with water.

A.5.2.11.18 TE buffer solution, $c(\text{tris}) = 0,010 \text{ mol/l}$ and $c(\text{Na}_2\text{EDTA}) = 0,001 \text{ mol/l}$.

A.5.2.11.19 Guanidine thiocyanate buffer, $c(\text{tris}\cdot\text{HCl}) \approx 4,6 \text{ mol/l}$, $c(\text{Na}_2\text{EDTA}\cdot 2\text{H}_2\text{O}) \approx 0,02 \text{ mol/l}$, $\rho[\text{Triton X } 100^3] \approx 11,8 \text{ g/l}$. Mix 2,6 g of Triton X 100³ (A.5.2.11.6) and 120 g of guanidine thiocyanate (A.5.2.11.1) and dissolve in 100 ml of tris·HCl buffer (A.5.2.11.15), 8,8 ml of EDTA solution (A.5.2.11.17) and 13,2 ml of water. The buffer has a final volume of approximately 220 ml at ~pH 7,2. The solution can be kept for at least 12 months at room temperature.

A.5.2.11.20 RNase A solution, $\rho(\text{RNase A}) \approx 10 \text{ mg/ml}$, in accordance with the manufacturer's instructions or Reference [3], B.17.

Dissolve 10 mg of RNase A (A.5.2.11.4) in 1,0 ml of tris·HCl + NaCl buffer (A.5.2.11.16). Temper the solution for 15 min to 20 min at 95 °C, allow to cool slowly to room temperature, and then divide into 50 µl aliquots. Store the solution frozen, avoiding repeated freezing and thawing.

A.5.2.11.21 Glycogen solution, $\rho(\text{glycogen}) \approx 20 \text{ mg/ml}$. Weigh 200 mg of glycogen into a 15 ml plastics centrifuge tube and dissolve in 10 ml of deionized water. The aliquoted solution can be kept for 24 months at 5 °C.

A.5.2.11.22 Elution buffer, (0,2 TE), $c(\text{tris}\cdot\text{HCl}) = 2 \text{ mmol/l}$, $c(\text{EDTA}) = 0,2 \text{ mmol/l}$, pH 8,0. Dissolve 158 mg of tris·HCl and 37 mg of EDTA disodium salt in approximately 400 ml of water; adjust to pH 8,0. Transfer the solution to a 500 ml volumetric flask and make up to volume with water. Autoclave the solution. The aliquoted solution can be kept for at least 12 months at room temperature.

A.5.2.11.23 MicroSpin S-300 HR column for additional DNA purification (Amersham Pharmacia Biotech)³⁾

A.5.2.11.24 Quant-iT PicoGreen dsDNA Quantitation Kit for fluorometric quantitation of DNA (Invitrogen), optional.⁵⁾

A.5.2.12 Sample collection, transport, preservation and storage

No specific requirements.

A.5.2.13 Test sample preparation

Apply the following DNA extraction and purification protocol. Scale adaptation of masses and buffer volumes is required as a function of the selected size of the test portion.

3) The column and test kit indicated are examples of suitable products available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of these products. Equivalent products may be used if they can be shown to lead to the same results. **A1**

A1) Usually, lecithin or plant oils which have not been refined can be weighed in directly. Viscous lecithin samples are heated to 60 °C first and shaken up well prior to being weighed in.

A.5.2.14 Instrument calibration

Instruments should be calibrated as per ISO/IEC 17025.^[39]

A.5.2.15 Method steps

A.5.2.15.1 General

For solvent extraction, the viscous sample material is treated with hexane and guanidine thiocyanate buffer. Interfering material is removed by extraction with chloroform and the remaining RNA is digested with RNase A. The DNA is precipitated in the presence of glycogen with isopropanol, washed with ethanol, and dissolved in an aqueous solution. For removal of PCR-inhibitors, the DNA is purified by means of gel filtration.

A.5.2.15.2 DNA extraction procedure

Ensure that the sample material is as homogeneous as possible and transfer 2,5 g to a 50 ml polypropylene centrifugation vessel.

An extraction blank control shall be processed in parallel.

Add 15 ml of *n*-hexane and dissolve lecithin.

Add 2 ml of the guanidine thiocyanate buffer and mix thoroughly (e.g. for 10 s to 20 s using a mixer).

Centrifuge for 10 min at approximately 4 000 × *g*.

Decant the major part of the upper hexane phase.

Transfer the lower, aqueous phase (without sediment) to a sterile 2 ml polypropylene reaction vessel.

Centrifuge for 10 min at 10 000 × *g*.

Transfer 1 ml of the lower, aqueous phase to a sterile 2 ml polypropylene reaction vessel.

Add 0,5 ml of chloroform and mix thoroughly for approximately 2 min.

Centrifuge for 10 min at 10 000 × *g*.

Transfer 500 µl to 750 µl of the aqueous supernatant to a new 1,5 ml polypropylene reaction vessel and add 5 µl of RNase A solution. (The RNase A digestion was performed in a collaborative trial and may be omitted, if applicable, in which case, proceed directly to the addition of glycogen.)

Incubate the mixture for 10 min at room temperature in order to hydrolyse any co-isolated RNA.

Add 4 µl of glycogen solution and 0,8 volumes of isopropanol (relative to the supernatant volume used), mix gently, and incubate for 30 min at room temperature.

Centrifuge for 10 min at no less than 12 000 × *g*.

Discard the supernatant; wash the DNA precipitate with 500 µl of ethanol.

Centrifuge for 5 min at no less than 12 000 × *g*.

Carefully draw off the supernatant, dry the DNA precipitate at room temperature or under vacuum.

Dissolve the DNA precipitate in 50 µl of elution buffer (e.g. in the refrigerator over night; heat slightly if not fully dissolved). **A1)**

A.5.2.15.3 DNA purification by means of gel filtration

A ready-to-use gel filtration column is used for additional DNA purification. Other columns or purification principles can be used if they are suitable for this purpose and perform equivalently.

Resuspend the resin in a Microspin S300 HR³) column by intensive vortexing.

Open the screw cap slightly by twisting by a quarter-turn and break off the lower column closure.

Place the Microspin S300 HR³) column into a 3 ml polypropylene reaction vessel.

Centrifuge at $770 \times g$ for 1 min.

Place the Microspin S300 HR³) column into a new 1,5 ml polypropylene reaction vessel; remove the screw cap.

Slowly load the DNA solution (50 μ l) on to the column.

Centrifuge at $770 \times g$ for 2 min.

The DNA is present in the eluate. DNA solutions should be kept refrigerated. If stored for 1–2 weeks, storage at 4 °C is sufficient; for longer storage periods, a storage temperature of –20 °C is required.

A.5.2.15.4 DNA quantitation

If the amount of DNA isolated from the sample is sufficient, DNA quantitation can be done spectrometrically using a UV-spectrometer as described in B.1. Alternatively, the concentration of DNA can be estimated fluorimetrically by the PicoGreen³) dsDNA method (Reference [17]) using a reagent kit system (A.5.2.11.24) or an equivalent product.

Since lecithin often has a low DNA content, the quantitation of the amplifiable soybean DNA is preferably done using real-time PCR (see B.3). A method for quantitation of the soybean lectin gene is indicated in ISO 21570:2005,^[41] C.2.

A.5.2.15.5 DNA integrity evaluation

The integrity of DNA obtained from the extractions is not evaluated further, since the quality criteria for the extracted DNA is its amplifiability in subsequent PCR analyses using soybean specific methods where the target DNA sequences are ≤ 100 bp.

A.5.2.15.6 Preparation of standards

Standard-DNA for calibration is prepared from soybean flour [ERM BF410a³]) using a CTAB protocol. The concentration of DNA is estimated fluorimetrically with the PicoGreen³) dsDNA method (Reference [17]).

A.5.2.15.7 Accept or reject criteria

The amount of amplifiable species DNA in the sample (copies per PCR), determined by means of real-time PCR, shall be sufficient to achieve a practical LOD of 0,9 % or less for a GMO-specific DNA.

A.5.2.15.8 Identification

The result of the DNA extraction method can be verified via soybean-specific reference gene methods as indicated in ISO 21570:2005,^[41] C.2. Other soybean-specific methods can be used, if they show equivalent or better performance.

A.5.2.16 Sample identification

All samples shall be identified unambiguously. $\text{\textcircled{A}}$

A.5.2.17 Calculations

For calculation of the practical LOD, see A.5.2.3.4.

A.5.2.18 Record keeping

Record keeping shall conform to ISO/IEC 17025.^[39]

A.5.2.19 Reporting

Reporting should be carried out as specified in ISO 24276 and other applicable standards (ISO/IEC 17025^[39]).

A.5.2.20 Safety measures

A hood is required when handling organic chemicals. If gloves are used, ensure that they are powder free. Cross-contamination can be prevented by using aerosol-protected pipette tips.

A.5.2.21 Pollution prevention and waste disposal

No specific requirements. Refer to ISO 24276. **A1**

Annex B (informative)

Methods for the quantitation of the extracted DNA

B.1 Basic ultraviolet spectrometric method

B.1.1 General

This annex describes a routine method to determine the concentration of DNA in solutions.

B.1.2 Validation status

The method has been widely ring-tested and results published^[35]. As an example, it was successfully applied within an interlaboratory study on GMO detection organized by the Federal Office of Public Health, Bern, and the Cantonal Laboratories of Basel and Zurich, Switzerland.

B.1.3 Principle

Nucleic acids in solution absorb ultraviolet (UV) light in the range from 210 nm to 300 nm with an absorption maximum at 260 nm. Since DNA, RNA and nucleotides have their absorption maximum at 260 nm, RNA and nucleotide contamination of DNA solutions cannot be determined by UV spectrometry. For this reason, RNA must be removed enzymatically during DNA extraction before DNA determination. Also, oligonucleotides and nucleotides derived from RNA hydrolysis should be eliminated (e.g. by silica treatment, as outlined in A.4.1.7.2). Oligonucleotides and nucleotides generated by RNase treatment, if not removed (e.g. by silica treatment) can lead to an overestimation of the DNA content of the sample. Moreover, double-stranded DNA absorbs less UV light compared to single-stranded DNA. Since the proportion of single-stranded DNA in the solution is unknown, to avoid overestimation of the DNA content, all the DNA in the test sample is converted to its single-stranded form by using the denaturing agent sodium hydroxide. Since nucleic acids do not absorb at 320 nm, reading at 320 nm is informative for the determination of background absorption due to light scattering and UV-active compounds.

The production of a calibration curve is not necessary, provided that an appropriate molar extinction coefficient is chosen as a function of the type of nucleic acid under study and/or its integrity.

However, the calibration of the spectrometer should be verified periodically by measuring the concentration of reference DNA solutions.

B.1.4 Application range

The method is applicable to DNA concentrations in the range from 2 µg/ml to 50 µg/ml. Before quantitation, suitable dilutions of the extracted DNA to be quantified should be made, in order to be in the linear range of the spectrometric measurement (optical density between 0,05 to 1).

NOTE Occasionally, residual compounds (e.g. CTAB from the DNA extraction procedure) may interfere with the UV spectrometric detection at 260 nm, because they absorb at this wavelength.

B.1.5 Reagents

B.1.5.1 Tris(hydroxymethyl)-aminomethane (Tris) ($C_4H_{11}NO_3$).

B.1.5.2 Sodium hydroxide (NaOH).

B.1.5.3 Hydrochloric acid, $\phi(\text{HCl}) = 37\%$.

B.1.5.4 Carrier DNA, e.g. Herring Sperm DNA¹¹⁾, or Calf Thymus DNA¹¹⁾.

B.1.5.5 DNA reference solution

Prepare a DNA stock solution with 10 mg/ml by dissolving 100 mg carrier DNA (B.1.5.4) in 10 ml of dilution buffer (B.1.5.7). DNA dissolves at this concentrations only slowly and the resulting solution is very viscous. Afterwards dilute this prepared stock reference DNA-solution further with dilution buffer up to the desired working concentration (e.g. 25 $\mu\text{g}/\text{ml}$).

B.1.5.6 Sodium hydroxide solution, $c(\text{NaOH}) = 2 \text{ mol/l}$.

B.1.5.7 Dilution buffer, $c(\text{Tris}) = 0,01 \text{ mol/l}$.

Adjust the pH to 9,0 with HCl.

B.1.6 Apparatus and equipment

B.1.6.1 UV-spectrometer, single-beam, double-beam or photodiode array instruments are suitable.

B.1.6.2 Mixer/shaker, e.g. Vortex^{® 2)}.

B.1.6.3 Measurement vessels, for example quartz cells/cuvettes or plastic cells/cuvettes suitable for UV detection at a wavelength of 260 nm.

The size of the measurement vessels used determines the volume for measurement: half-micro cells (1 000 μl), micro cells (400 μl), ultra-micro cells (100 μl) and quartz capillaries (3 μl to 5 μl). The optical path of standard cell is usually 1 cm.

B.1.7 Procedure

B.1.7.1 Measurement of a reference DNA solution

The correct calibration of the spectrometer can be verified by the use of a reference DNA solution, as follows:

- for blank measurement only dilution buffer (B.1.5.7) is used to fill the measurement vessel;
- the measurement vessel is filled with the reference DNA solution (B.1.5.5).

Absorption is measured for both the blank and reference DNA solutions at wavelengths of 260 nm and 320 nm.

11) These are available from Sigma as D-7290 and D-1501, respectively. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product. Equivalent products may be used if they can be shown to lead to the same results.

B.1.7.2 Measurement of a test DNA solution of unknown concentration

For the blank solution, mix dilution buffer (B.1.5.7) plus sodium hydroxide solution (B.1.1.5.13), so that a final NaOH substance concentration of 0,2 mol/l is reached. This mixture is used to fill the measurement vessel.

Mix the test DNA solution with sodium hydroxide solution and, if needed, with dilution buffer, to obtain a final NaOH substance concentration of 0,2 mol/l. This mix is used to fill the measurement vessel.

Measure the absorption after 1 min incubation time for both the blank and reference DNA solutions at wavelengths of 260 nm and 320 nm. The reading is stable for at least 1 h.

EXAMPLE 1 For the blank solution, mix 90 µl of dilution buffer and 10 µl of sodium hydroxide solution and transfer to a 100 µl measurement vessel.

EXAMPLE 2 For the test DNA solution, mix 80 µl of dilution buffer or water, 10 µl of sodium hydroxide solution and 10 µl of a DNA solution of unknown concentration and transfer to a 100 µl measurement vessel.

B.1.8 Evaluation

The absorption (OD) at 320 nm (background) is subtracted from the absorption at 260 nm, resulting in the corrected absorption at 260 nm.

If the corrected OD at 260 nm equals 1, then the estimated DNA concentration is 50 µg/ml for double-stranded DNA, or 37 µg/ml for single-stranded DNA (i.e. denatured with sodium hydroxide), respectively.

Reliable measurements require OD values at a wavelength of 260 nm to be greater than 0,05.

Finally, calculate the mass concentration, ρ , of the double-stranded test DNA solution, taking into consideration the denaturation and the dilution factor applied according to Equation (1):

$$\rho_{\text{DNA}} = F \times (\text{OD}_{260} - \text{OD}_{320}) \times 37 \quad (1)$$

where

F is the dilution factor;

OD_{260} is the absorbance at 260 nm;

OD_{320} is the absorbance at 320 nm;

37 is the conversion factor, in micrograms per millilitre.

EXAMPLE For a calculation with a dilution factor of 10 and an OD_{260} of 0,658 and an OD_{320} of 0,040:

$$\rho_{\text{DNA}} = 10 \times (0,658 - 0,040) \times 37 \text{ µg/ml} = 229 \text{ µg/ml}.$$

B.2 Agarose gel electrophoresis and ethidium bromide staining method

B.2.1 General

This annex describes a routine method to determine the concentration of DNA in solutions. Agarose gel electrophoresis of DNA and ethidium bromide (EtBr) staining provide a way of estimating the quantity of DNA and to analyse at the same time its physical state (e.g. degree of degradation, presence of residual RNA and of some contaminants). The method also applies if insufficient DNA is available for spectrometric detection, or if the DNA is not sufficiently purified and may contain substances that absorb ultraviolet radiation^[36]. Gel electrophoresis is usually not recommended to quantify DNA if degraded. If this is the case, other methods should be employed.

B.2.2 Validation status

The method has been known for many years and has been widely applied, although it has never been tested in interlaboratory studies for GMO detection in foodstuffs.

B.2.3 Principle

DNA separates electrophoretically, on the basis of its charge and molecular mass, when loaded onto a molecular sieve (agarose gel) and subjected to an electric field in the presence of a buffer solution^[37].

EtBr intercalates into the DNA and, when excited by ultraviolet light, emits orange fluorescence. Because the amount of fluorescence is proportional to the total mass of DNA, the quantity of DNA in the sample can be estimated by comparing the fluorescence produced by the unknown sample with that of a series of quantity standards. The molecular mass of such standards must be similar to that of the DNA under quantitation, because EtBr intercalation, and thus fluorescence emission, also depends on the length of the DNA fragments. EtBr also stains single-stranded DNA and RNA. For a more precise estimation of the DNA content, RNA must be removed enzymatically.

B.2.4 Application range

The method is applicable to DNA concentrations in the approximate range from 5 ng to 500 ng when photographic image acquisition systems are used. CCD-based systems for video documentation may be more sensitive.

B.2.5 Safety precautions

EtBr is a powerful mutagen and a carcinogen and must be handled with care. The use of gloves is compulsory. All solutions and gels containing EtBr should be decontaminated before disposal, see Reference [36].

Ultraviolet light (UV-C) is dangerous especially for eye retina. Always wear UV protection devices (facial mask in particular) when using it.

B.2.6 Reagents

The agarose gel electrophoresis may be carried out as TAE buffer electrophoresis or as TBE buffer electrophoresis.

Usually, molecular biology grade is not necessary for the reagents described in this method. Solutions as described in this method do not usually need to be autoclaved.

B.2.6.1 Agarose, suitable for DNA electrophoresis and for the intended size separation of the DNA molecules.

B.2.6.2 Boric acid (H_3BO_3), for the TBE buffer system only.

B.2.6.3 Bromophenol blue ($\text{C}_{19}\text{H}_9\text{Br}_4\text{O}_5\text{SNa}$) and/or xylene cyanole FF ($\text{C}_{25}\text{H}_{27}\text{N}_2\text{O}_6\text{S}_2\text{Na}$).

B.2.6.4 DNA quantity standard, of suitable molecular mass (e.g linearized Lambda Phage DNA for high molecular mass DNA and restriction-digested Lambda Phage DNA for lower molecular mass DNA).

B.2.6.5 DNA molecular mass standard, for example a commercial preparation containing DNA fragments from very high to very low molecular mass.

B.2.6.6 Glacial acetic acid (CH_3COOH), for the TAE buffer system only.

B.2.6.7 Ethylenediaminetetraacetic acid disodium salt ($\text{Na}_2\text{-EDTA}$) ($\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_8\text{Na}_2$).

B.2.6.8 Ethidium bromide (EtBr) ($C_{21}H_{20}N_3Br$).

B.2.6.9 Glycerol ($C_3H_8O_3$).

B.2.6.10 Sodium acetate ($C_2H_3O_2Na$), for the TAE buffer system only.

B.2.6.11 Hydrochloric acid, $\phi(HCl) = 37\%$.

B.2.6.12 Sodium hydroxide (NaOH).

B.2.6.13 Tris(hydroxymethyl)-aminomethane (Tris) ($C_4H_{11}NO_3$).

B.2.6.14 TAE buffer solution (1x), $c(Tris) = 0,050\text{ mol/l}$, $c(C_2H_3O_2Na) = 20\text{ mmol/l}$, $c(Na_2-EDTA) = 0,001\text{ mol/l}$.

Adjust the pH to 8,0 with glacial acetic acid or NaOH. It is advisable to prepare the TAE buffer solution as a concentrated stock solution (maximum 50-fold concentrated). Discard it if a precipitate is visible. Dilution of the concentrated electrophoresis buffers can be carried out, immediately before its use, with non-sterile, (mono)-distilled or deionised water.

B.2.6.15 Tris/borate (TBE) buffer solution (0,5x), $c(Tris) = 0,055\text{ mol/l}$, $c(\text{boric acid}) = 0,055\text{ mol/l}$, $c(Na_2EDTA) = 0,001\text{ mol/l}$.

Adjust the pH to 8,0 with HCl or NaOH. It is advisable to prepare the TBE buffer solution as a concentrated stock solution (maximum 10-fold concentrated). Discard it if precipitation is visible. Dilution of the concentrated electrophoresis buffers can be carried out, immediately before its use, with non-sterile, (mono)-distilled or deionised water.

B.2.6.16 Sample loading buffer solution (5x), $\phi(\text{glycerol}) = 50\%$, $\rho(\text{bromophenol blue}) = 2,5\text{ g/l}$ and/or $\rho(\text{xylene cyanol}) = 2,5\text{ g/l}$, dissolved in electrophoresis buffer solution (B.2.6.14 or B.2.6.15).

B.2.6.17 Ethidium bromide solution, $c(EtBr) = 0,5\text{ mg/l}$.

It is advisable to store the ethidium bromide solution as a concentrate (e.g. 10mg/ml) at 5° C in the dark (EtBr is light-sensitive). It is also advisable to avoid weighing EtBr. The stock solution should be prepared by dissolving an appropriate amount of water in the vessel already containing the EtBr powder, or alternatively, by employing pre-weighed EtBr tablets. Solubilization of EtBr should be carried out protected from light, under agitation at room temperature. This usually takes approximately 1 h.

B.2.7 Apparatus and equipment

B.2.7.1 Microwave oven or boiling water bath

B.2.7.2 Equipment for agarose gel electrophoresis, with accessories and power supply.

B.2.7.3 Ultraviolet (UV) trans-illuminator or lamp, preferably with wavelength of 312 nm.

Alternatively, equipment for column chromatography of nucleic acids and the according detection system or other similar suitable systems may be used.

B.2.7.4 Recording instrument, for example a photo documentation system with 3 000 ASA films and UV filter adequate for EtBr-emitted fluorescence.

As an alternative, a video-documentation system with CCD camera, adequate UV filter and (optional) quantitative analysis software may be used.

B.2.8 Procedure

B.2.8.1 General

The agarose gel electrophoresis may be carried out as TAE buffer electrophoresis or as TBE buffer electrophoresis. Use the same buffer to dissolve the agarose and to fill the electrophoresis tank.

B.2.8.2 Agarose gel preparation

Gels should not be thicker than 1 cm.

The agarose concentration and quality determines the resolution capacity of the gel. For high molecular mass DNA quantitation, agarose concentrations between 8 g/l and 10 g/l are used. For low molecular mass DNA (e.g. degraded or restricted) higher agarose concentrations are used (up to 40 g/l)^[38].

Weigh an appropriate amount of agarose (B.2.6.1) and add it to the electrophoresis buffer solution (B.2.6.14 or B.2.6.15). Allow the solution to boil in a microwave oven or in a water bath (B.2.7.1), until the agarose is completely dissolved. Replace the volume lost by evaporation with an equivalent amount of water, mix by swirling (avoid air bubbles trapping), cool down the solution to about 60° C and keep it at this temperature until usage. Prepare a gel support (gel tray) with a suitable sample comb placed in right position. Pour the agarose solution onto the gel tray and allow the gel to solidify at room temperature (1 h is usually recommended).

B.2.8.3 DNA sample preparation

Mix the sample DNA solutions (e.g. 5 µl to 10 µl) with approximately 20 % (with respect to the final sample volume) of loading buffer (B.2.6.16) (e.g. add 2,5 µl of loading buffer to 10 µl of DNA sample), mix and apply the mixture to the sample slots (wells) with a micropipette. If the unknown samples are suspected to be too concentrated, also provide some dilutions of them to be loaded onto the gel.

To determine the size of the extracted DNA fragments, add the sample loading buffer (B.2.6.16) in the proportion of 20 % with respect to the sample volume) to a suitable amount of the DNA molecular mass standard (B.2.6.5) and carry out electrophoresis in parallel.

To estimate the concentration of the unknown sample, run standard DNA quantity samples in parallel. Such samples contain known amounts (within the dynamic range of the method, i.e. 5 ng to 500 ng) of the DNA quantity standard (B.2.6.4) diluted in water or in electrophoresis buffer (B.2.6.14 or B.2.6.15). It is recommended to use quantitation standards containing at least 5 calibration points (i.e. different amounts of DNA).

B.2.8.4 Submarine electrophoresis

Carefully remove the samples comb from the gel. Transfer the gel (with its gel tray) to the electrophoresis cell, so that the wells reside closer to the cathode (negative electrode). Fill the cell with the electrophoresis buffer (B.2.6.14 or B.2.6.15). Overlay the gel with approximately 2 mm of the same buffer and load the samples using a micropipette.

Carry out the electrophoresis at room temperature at the appropriate voltage and power intensity (generally a maximum constant voltage of 5 V/cm, with respect to the distance between the electrodes, is recommended). Under the described conditions, DNA is negatively charged, so it migrates from the cathode to the anode. The electrophoresis time depends on the migration distance required, on the current generated by the power supply, the buffer used, the electro-endosmosis and the concentration of the agarose in the gel.

B.2.8.5 Staining

After completing the electrophoresis, incubate the gel for 15 min to 50 min in the ethidium bromide solution (B.2.6.17) at room temperature, possibly in the dark (and/or in a stainless steel tank with a cover) with gentle shaking.

If necessary, reduce the background staining by de-staining the gel in water for 10 min to 30 min.

As an alternative to post-electrophoresis staining, EtBr can be added to the gel before pouring it. In this case, EtBr is added to the gel to a final concentration of 0,01 mg per millilitre of gel when the gel has been cooled to a temperature of 60 °C.

If the gel is cast with ethidium bromide, load the unknown sample and the DNA quantity standard (B.2.6.4) into separate slots produced with the same comb on the same gel. Otherwise the quantity of ethidium bromide will be different for the two, so yielding erroneous quantitation results. To minimize the problems of EtBr movement in the gel, some EtBr can also be added to the electrophoresis (tank) buffer. After the gel electrophoresis, no de-staining step is usually required.

B.2.8.6 Gel recording

Transfer the gel to the trans-illuminator surface, switch on the UV light and record the DNA fluorescence by photography or video-documentation.

B.2.9 Evaluation/interpretation

The DNA content of the sample is estimated by comparing the unknown samples with the DNA quantity standard samples that underwent electrophoresis in parallel. This evaluation can be carried out visually or, better, with the aid of a quantitation software able to calculate an adequate calibration curve.



B.3 Real time PCR method for quantitation of extracted DNA

When DNA-poor matrices are extracted, the quantitation of the obtained DNA is not always possible with the usual physical methods (e.g. the ones described in this annex), because of their insufficient sensitivity. Where such quantitation is necessary, a real time PCR approach can be followed. This method also provides information on the PCR amplificability of isolated DNA molecules that cannot be provided by physical measurements of DNA concentrations. For details, see ISO 21570.

Bibliography

- [1] AUSUBEL, F.M., BRENT, R., KINGSTON, R.E., MOORE, D.D., SEIDMAN, J.G., SMITH, J.A., STRUHL, K.: *Current Protocols in Molecular Biology*. John Wiley & Sons (1988) (ISBN 0-471-50338-X)
- [2] SAMBROOK, J., RUSSEL D "Molecular Cloning, a Laboratory Manual" 3rd Edition, Cold Spring Harbor Laboratory Press, NY (2001) (ISBN 0-87969-576-5)
- [3] ^{A1} SAMBROOK J., FRITSCH E.F., MANIATIS, T., eds. *Molecular cloning: A laboratory manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 2nd edition, 1989 [Superseded by GREEN M.R., SAMBROOK J., eds. *Molecular cloning: A laboratory manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 4th Edition, 2012] ^{A1}
- [4] PROKISCH, J. ZELNY, R., TRAPMANN, S., LE GUERN, L., SCHIMMEL, H., KRAMER, G.N, PAUWELS, J. Estimation of the minimum uncertainty of DNA concentration in a genetically modified maize sample candidate certified reference material. *Fresenius J. Analy. Chem.* (2001) **370**(7) pp. 935-9
- [5] SAMBROOK, J., FRITSCH, E.F and MANIATIS, T. In: *Molecular Cloning: a Laboratory Manual*. 2nd ed. Vol. 3, Appendix E.3, *Purification of nucleic acids*. (1987) (ISBN 0-87969-309-6)
- [6] Detection of a genetic modification of *Lactobacillus curvatus* in uncooked-meat sausage by amplification of the modified DNA sequence using the polymerase chain reaction (PCR) and hybridization of the PCR product with a DNA probe. L 08.00-44 In: *Collection of official methods under article 35 of the German Federal Foods Act; Methods of sampling and analysis of foods, tobacco products, cosmetics and commodity goods*. Federal Health Office, September 1998, revised version: state 05/2001; Berlin, Köln, Beuth Verlag GmbH
- [7] STRAUB, J. A., HERTEL, C. and HAMMES, W.P. The fate of recombinant DNA in thermally treated fermented sausages. *Eur. Food Res. Technol.* (1999) **210**, pp. 62-67
- [8] STRAUB, J. A., HERTEL, C. and HAMMES, W. P. A 23S rDNA-targeted polymerase chain reaction-based system for detection of *Staphylococcus aureus* in meat starter cultures and dairy products. *J. Food Protect.* (1999) **62**, pp. 1150-1156
- [9] LICK, S., KELLER, M., BOCKELMANN, W., HELLER, K.J. Optimized DNA extraction method for starter cultures from yoghurt. *Milk Sci. Int.* (1998) **51**, pp. 183-186
- [10] LICK, S., HELLER, K.J. Quantitation by PCR of yoghurt starters in a model yoghurt produced with a genetically modified *Streptococcus thermophilus*. *Milk Sci. Int.* (1998) **53**, pp. 671-675
- [11] Detection of a genetic modification of *Streptococcus thermophilus* in yoghurt by amplification of the modified DNA sequence by means of the polymerase chain reaction (PCR) and hybridization of the PCR product with a DNA probe. L 02.02-4 In: *Collection of official methods under article 35 of the German Federal Foods Act; Methods of sampling and analysis of foods, tobacco products, cosmetics and commodity goods*. Federal Health Office, September 1998, revised version: state 05/2001; Berlin, Köln, Beuth Verlag GmbH
- [12] LICK, S., KELLER, M., KRUSCH, U., HELLER, K.J. Identification of starter cultures in thermally treated plain yoghurt using geneprobos and PCR. *J. Dairy Res.* (1998) **63**, pp. 607-613
- [13] VAN VAERENBERGH, B., GROOTAERT, B., MOENS, W. Validation of a method for the preparation of fungal genomic DNA for Polymerase chain reaction (PCR) and Random Amplification of Polymorphic DNA (RAPD). *J. Mycol. Méd.* (1995) **5**, pp. 133-139
- [14] HAYNES, K.A., WESTERNENG, T.J., FELL J.W., MOENS, W. Rapid detection and identification of pathogenic fungi by polymerase chain amplification of large subunit ribosomal DNA. *J. Med. & Vet. Mycology* (1995) **33**, pp. 319-325

- [15] GUILLAUME, G.; VERBRUGGE, D., CHASSEUR-LIBOTTE, M-L., MOENS, W., COLLARD, J-M.: PCR typing of tetracycline determinants (Tet A-E) in *Salmonella enterica* Hadar and in the microbial community of activated sludges from hospital and urban wastewater treatment facilities in Belgium. In: *FEMS Microbiology Ecology*, 2000, **32**, pp. 77-85
- [16] PEDERSEN, LH., SKOUBOE, P., BOYSEN, M., SOULE, J., ROSSEN, L: Detection of *Penicillium* species in complex food samples using the polymerase chain reaction. *Int J Food Microbiol*, 1997, **35**(2), pp. 169-77
- [17] ^{A1}AHN S.J., COSTA J., EMANUEL J.R. PicoGreen quantitation of DNA: Effective evaluation of samples pre- or post-PCR. *Nucleic Acids Res.* 1996, **13** pp. 2623–2625 ^{A1}
- [18] HAUGLAND, R.A., HECKMAN, J.L., WYMER, L.J. Evaluation of different methods for the extraction of fungal conidia by quantitative competitive PCR analysis. In: *J.Microbiol. Methods*, 1999, **37**(2), pp. 165-176
- [19] KIM, C.S., LEE, C. H., SHIN, J.S., CHUNG, Y.S., and HYUNG, N.I. A simple and rapid method for isolation of high quality genomic DNA from fruit trees and conifers using PVP. *Nucleic Acids Research*, 1997, **25**, No. 5, pp. 1085-1086
- [20] BERTHELET, M., WHYTE, L.G., and GREER, C. W.: Rapid, direct extraction of DNA from soils for PCR analysis using polyvinylpolypyrrolidone spin columns. *FEMS Microbiology Letters*, 1996, **138**, pp. 17-22
- [21] PETIT, F., CRAQUELIN, S., GUESPIN-MICHEL, J., and BUFFET-JANVRESSE, C. Nucleic acid extraction from polluted estuarine water for detection of viruses and bacteria by PCR and RT-PCR analysis. *Res. Microbiol.*, 1999, **150**, pp. 143-151
- [22] WATSON, R.J., and BLACKWELL, B.: Purification and characterization of a common soil component which inhibits the polymerase chain reaction. *Can J. Microbiol.*, 2000, **46**, pp. 633-642
- [23] GRIFFITHS, R. J., WHITELEY, A.S., O'DONNELL, A.G., and BAILEY, M.J.: Rapid Method for Coextraction of DNA and RNA from Natural Environments for Analysis of Ribosomal DNA- and rRNA-Based Microbial Community Composition. *Appl. Environ. Microbiol.*, 2000, **66**, No. 12, pp. 5488-5491
- [24] ROGERS, S.O. and BENDICH, A.J.. *Extraction of DNA from plant tissues. Plant Molecular Biology Manual A6* (1988) pp. 1-10
- [25] Analysis of foodstuffs: Detection of a genetic modification of soya beans by amplification of the modified DNA sequence using the polymerase chain reaction (PCR) and hybridisation of the PCR product with a DNA probe. L 23.01.22-1, March 1998. In: Amtliche Sammlung von Untersuchungsverfahren nach § 35 LMBG; Verfahren zur Probenahme und Untersuchung von Lebensmitteln, Tabakerzeugnissen, kosmetischen Mitteln und Bedarfsgegenständen/ BgVV (In: *Official Collection of methods under article 35 of the German Federal Food Act; Methods of sampling and analysis of foods, tobacco products, cosmetics and commodity goods/BgVV*). March 1998 Bd. 1 1999-03 Vol. 1) Berlin, Köln: Beuth Verlag GmbH
- [26] Analysis of foodstuffs: Detection of a genetic modification of potatoes by amplification of the modified DNA sequence using the polymerase chain reaction (PCR) and hybridization of the PCR product with a DNA probe. L 24.01-01, February 1996, revised in January 1997. In: Amtliche Sammlung von Untersuchungsverfahren nach § 35 LMBG; Verfahren zur Probenahme und Untersuchung von Lebensmitteln, Tabakerzeugnissen, kosmetischen Mitteln und Bedarfsgegenständen/ BgVV (In: *Official Collection of methods under article 35 of the German Federal Food Act; Methods of sampling and analysis of foods, tobacco products, cosmetics and commodity goods/BgVV*). Jan 1997 Bd. 1 (1997-01 Vol. 1) Berlin, Köln: Beuth Verlag GmbH

- [27] Analysis of foodstuffs: Detection of a genetic modification of tomatoes bean by amplification of the modified DNA sequence by means of the polymerase chain reaction (PCR) and hybridisation of the PCR product with a DNA probe. L 25.03.01-1, November 1999. In: Amtliche Sammlung von Untersuchungsverfahren nach § 35 LMBG; Verfahren zur Probenahme und Untersuchung von Lebensmitteln, Tabakerzeugnissen, kosmetischen Mitteln und Bedarfsgegenständen/ BgVV (In: *Official Collection of methods under article 35 of the German Federal Food Act; Methods of sampling and analysis of foods, tobacco products, cosmetics and commodity goods/BgVV*). Nov 1999 Bd. 1 (1999-11 Vol. 1) Berlin, Köln: Beuth Verlag GmbH
- [28] BOYLE, J.S., and LEW, A.M. An inexpensive alternative to glassmilk for DNA purification, *Trends in Genetics*, Vol. **11**(1), p. 8, 1995
- [29] BRODMANN, P., EUGSTER, A., HÜBNER, Ph., MEYER, R., PAULI, U., VÖGELI, U. and LÜTHY, J. Nachweis gentechnisch veränderter Roundup Ready™ Sojabohnen mittels der Polymerase - Kettenreaktion (PCR). *Methodenvorprüfung im Rahmen der Subkommission 29a des Schweizerischen Lebensmittelbuches. Mitt. Gebiete Lebensm. Hyg.* 1997, **88**, pp. 722-731
- [30] HÜBNER, P., STUDER, E., and LÜTHY, J. Quantitative Competitive PCR for the Detection of Genetically Modified in Food. *Food Control*, 1999, **10**, pp. 353-358
- [31] HÜBNER, P., STUDER, E., and LÜTHY, J. Quantitation of genetically modified organisms in food. *Nat Biotechnol.* 1999, **17**, pp. 1137-1138
- [32] MELZAK, K.A., SHERWOOD, C.S., TURNER, R.F.B. and HAYNES, C.A. Driving Forces for DNA Adsorption to Silica in Perchlorate Solutions. *J. Colloid and Interface Science*, 1996, **181**, pp. 635-644
- [33] Patents: EP 0389063, USP5,234,809
- [34] SAMBROOK, J., FRITSCH, E.F, and MANIATIS, T. In: *Molecular Cloning: a Laboratory Manual*. 2nd ed., Vol. 5, Appendix B 16, 1989 *Proteinase-K*. (ISBN 0-87969-509-6)
- [35] Swiss Food Manual, Chapter 52B, Section 1 to 5.(2000) *Eidgenössische Drucksachen und Materialzentrale*, CH-5005 Bern, available on CD.
- [36] SAMBROOK, J., FRITSCH, E.F and MANIATIS, T. In: *Molecular Cloning: a Laboratory Manual*. 2nd ed., Vol. 5, Appendix 5, (1989) *Quantitation of DNA and RNA* (ISBN 0-88989-509-8)
- [37] SAMBROOK, J., FRITSCH, E.F and MANIATIS, T. In: *Molecular Cloning: a Laboratory Manual* 2nd ed., Vol. 1, Chapter 6, (1989) *Gel electrophoresis of DNA* (ISBN 0-88989-509-8,)
- [38] SAMBROOK, J., FRITSCH, E.F and MANIATIS, T. In: "Molecular Cloning: a Laboratory Manual" 2nd ed., Vol. 1, Section 6.5, table 8.1 (1989) (ISBN 0-88989-509-8)
- [39] ISO/IEC 17025, *General requirements for the competence of testing and calibration laboratories*.
- [40] ISO 21569, *Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — Qualitative nucleic acid based methods*
- [41] ISO 21570, *Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — Quantitative nucleic acid based methods*
- [42] ISO Guide 30, *Terms and definitions used in connection with reference materials*
- [43]  ISO 5725-1, *Accuracy (trueness and precision) of measurement methods and results — Part 1: General principles and definitions*
- [44] WURZ A., RÜGGERBERG H., BRODMANN P., WAIBLINGER H.-U., PIETSCH K. DNA-Extraktionsmethode für den Nachweis gentechnisch veränderter Soja in Sojalecithin [DNA extraction method for the detection of genetically modified soy in soy lecithin]. *Dtsch. Lebensm. Rundsch.* 1998, **94** pp. 159–161 

- [45] ^{A1} WAIBLINGER H.-U., ERNST B., GRAF N., PIETSCH K. Ringtrial validation of a method for the extraction of DNA from soy lecithins. *J. Verbrauch. Lebensm.* 2007, **2** pp. 113–115
- [46] HORWITZ W. Protocol for the design, conduct and interpretation of method-performance studies. *Pure Appl. Chem.* 1995, **67**, pp. 331–343 ^{A1}

British Standards Institution (BSI)

BSI is the independent national body responsible for preparing British Standards and other standards-related publications, information and services. It presents the UK view on standards in Europe and at the international level.

BSI is incorporated by Royal Charter. British Standards and other standardization products are published by BSI Standards Limited.

Revisions

British Standards and PASs are periodically updated by amendment or revision. Users of British Standards and PASs should make sure that they possess the latest amendments or editions.

It is the constant aim of BSI to improve the quality of our products and services. We would be grateful if anyone finding an inaccuracy or ambiguity while using British Standards would inform the Secretary of the technical committee responsible, the identity of which can be found on the inside front cover. Similar for PASs, please notify BSI Customer Services.

Tel: +44 (0)20 8996 9001 Fax: +44 (0)20 8996 7001

BSI offers BSI Subscribing Members an individual updating service called PLUS which ensures that subscribers automatically receive the latest editions of British Standards and PASs.

Tel: +44 (0)20 8996 7669 Fax: +44 (0)20 8996 7001

Email: plus@bsigroup.com

Buying standards

You may buy PDF and hard copy versions of standards directly using a credit card from the BSI Shop on the website www.bsigroup.com/shop. In addition all orders for BSI, international and foreign standards publications can be addressed to BSI Customer Services.

Tel: +44 (0)20 8996 9001 Fax: +44 (0)20 8996 7001

Email: orders@bsigroup.com

In response to orders for international standards, BSI will supply the British Standard implementation of the relevant international standard, unless otherwise requested.

Information on standards

BSI provides a wide range of information on national, European and international standards through its Knowledge Centre.

Tel: +44 (0)20 8996 7004 Fax: +44 (0)20 8996 7005

Email: knowledgecentre@bsigroup.com

BSI Subscribing Members are kept up to date with standards developments and receive substantial discounts on the purchase price of standards. For details of these and other benefits contact Membership Administration.

Tel: +44 (0)20 8996 7002 Fax: +44 (0)20 8996 7001

Email: membership@bsigroup.com

Information regarding online access to British Standards and PASs via British Standards Online can be found at

www.bsigroup.com/BSOL

Further information about British Standards is available on the BSI website at **www.bsigroup.com/standards**

Copyright

All the data, software and documentation set out in all British Standards and other BSI publications are the property of and copyrighted by BSI, or some person or entity that owns copyright in the information used (such as the international standardization bodies) has formally licensed such information to BSI for commercial publication and use. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI. This does not preclude the free use, in the course of implementing the standard, of necessary details such as symbols, and size, type or grade designations. If these details are to be used for any other purpose than implementation then the prior written permission of BSI must be obtained. Details and advice can be obtained from the Copyright & Licensing Department.

Tel: +44 (0)20 8996 7070

Email: copyright@bsigroup.com

BSI

389 Chiswick High Road London W4 4AL UK

Tel +44 (0)20 8996 9001

Fax +44 (0)20 8996 7001

www.bsigroup.com/standards