

Soil improvers and growing media — Determination of nitrogen —

Part 1: Modified Kjeldahl method

The European Standard EN 13654-1:2001 has the status of a
British Standard

ICS 65.080

National foreword

This British Standard is the official English language version of EN 13654-1:2001.

The UK participation in its preparation was entrusted to Technical Committee AW/20, Top soil and other growing media, which has the responsibility to:

- aid enquirers to understand the text;
- present to the responsible European committee any enquiries on the interpretation, or proposals for change, and keep the UK interests informed;
- monitor related international and European developments and promulgate them in the UK.

A list of organizations represented on this committee can be obtained on request to its secretary.

Cross-references

The British Standards which implement international or European publications referred to in this document may be found in the BSI Standards Catalogue under the section entitled “International Standards Correspondence Index”, or by using the “Find” facility of the BSI Standards Electronic Catalogue.

A British Standard does not purport to include all the necessary provisions of a contract. Users of British Standards are responsible for their correct application.

Compliance with a British Standard does not of itself confer immunity from legal obligations.

This British Standard, having been prepared under the direction of the Consumer Products and Service Sector Policy and Strategy Committee, was published under the authority of the Standards Policy and Strategy Committee on 09 November 2001

Summary of pages

This document comprises a front cover, an inside front cover, the EN title page, pages 2 to 10, an inside back cover and a back cover.

The BSI copyright date displayed in this document indicates when the document was last issued.

Amendments issued since publication

Amd. No.	Date	Comments

© BSI 09 November 2001

ISBN 0 580 38268 0

ICS 65.080

English version

Soil improvers and growing media - Determination of nitrogen - Part 1: Modified Kjeldahl method

Amendements du sol et supports de culture -
Détermination de l'azote - Partie 1: Méthode de Kjeldahl
modifiée

Bodenverbessungsmittel und Kultursubstrate - Bestimmung
von Stickstoff - Teil 1: Modifiziertes Verfahren nach Kjeldahl

This European Standard was approved by CEN on 11 August 2001.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Czech Republic, Denmark, Finland, France, Germany, Greece, Iceland, Ireland, Italy, Luxembourg, Netherlands, Norway, Portugal, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: rue de Stassart, 36 B-1050 Brussels

Contents

	page
Foreword	3
1 Scope	4
2 Normative references	4
3 Terms and definitions	4
4 Principle	4
5 Reagents	4
6 Apparatus	5
7 Test sample	5
8 Procedure	5
9 Calculation	7
10 Expression of results	7
11 Precision	7
12 Test Report	7
Annex A (informative) Results of an interlaboratory trial to determine nitrogen (modified Kjeldahl method)	8
Annex B (informative) Methods of analysis used in the interlaboratory trial	9
Bibliography	10

Foreword

This European Standard has been prepared by Technical Committee CEN/TC 223 "Soil improvers and growing media", the secretariat of which is held by BSI.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by March 2002, and conflicting national standards shall be withdrawn at the latest by March 2002.

The annexes A and B are informative.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Czech Republic, Denmark, Finland, France, Germany, Greece, Iceland, Ireland, Italy, Luxembourg, Netherlands, Norway, Portugal, Spain, Sweden, Switzerland and the United Kingdom.

SAFETY PRECAUTIONS — Care should be taken when handling samples that may contain sharps or is of a dusty nature.

1 Scope

This European Standard specifies a method for the determination of nitrogen in soil improvers and growing media. The Kjeldahl method determines ammonium-N, nitrate-N, nitrite-N and organic N content of soil improvers and growing media. Nitrogen in N-N-linkages, N-O-linkages and some heterocyclics (especially pyridine) is only partially determined. [6], [7], [8]

The method is not applicable to liming materials and preformed materials such as mineral wool slabs and foam slabs.

NOTE The requirements of the standard may differ from the national legal requirements for the declaration of the products concerned.

2 Normative references

This European Standard incorporates by dated or undated reference, provisions from other publications. These normative references are cited at the appropriate places in the text, and the publications are listed hereafter. For dated references, subsequent amendments to or revisions of any of these publications apply to this European Standard only when incorporated in it by amendment or revision. For undated references the latest edition of the publication referred to applies (including amendments).

EN ISO 3696, *Water for analytical laboratory use - Specification and test methods (ISO 3696:1987)*.

EN 13040:1999, *Soil improvers and growing media - Sample preparation for chemical and physical test, determination of dry matter content, moisture content and laboratory compacted bulk density*.

3 Terms and definitions

For the purposes of this standard the terms and definitions given in EN 13040 apply.

4 Principle

The nitrogen content of the sample is determined using a method based on a sulfuric acid/potassium sulfate digestion. Copper sulfate is used as the catalyst.

5 Reagents

5.1 General

All reagents shall be of recognized analytical grade. Use water of grade 2 complying with EN ISO 3696.

5.2 Salicylic acid/sulfuric acid, dissolve 25 g of salicylic acid ($\text{HO.C}_6\text{H}_4\text{COOH}$) in 1000 ml of concentrated sulfuric acid ($\rho = 1,84 \text{ g/ml}$).

5.3 Potassium sulfate catalyst mixture, grind and thoroughly mix 200 g of potassium sulfate, 6 g of copper (II) sulfate pentahydrate. (This mixture may be available commercially).

5.4 Sodium thiosulfate pentahydrate, crush the crystals to form a powder that passes through a sieve with an aperture of 0,25 mm.

5.5 Calibration substances, pure substances of known nitrogen content, for example acetanilide (C_8H_9NO), L-aspartic acid ($C_4H_7NO_4$), or amino acids of known composition.

NOTE The total content of nitrogen of the calibration substance should be as similar to the sample nitrogen content as possible.

6 Apparatus

Usual laboratory apparatus, and in particular the following :

6.1 Analytical balance, capable of weighing accurately to 1,0 mg.

6.2 Kjeldahl digestion flasks or tubes, of nominal value 50 ml, suitable for the digestion system (6.3)

NOTE Larger flasks (300 ml to 500 ml) may be required for bulky materials or where a larger sample mass is used.

6.3 Digestion or mineralisation system with variable heating controls.

6.4 Sieve, aperture 5 mm

6.5 Stand, capable of holding the digestion flasks or tubes.

7 Test sample

Prepare the test sample in accordance with EN 13040:1999, clause 9.

Losses of nitrogen can occur with samples of high ammonium-N content in alkaline situations and particularly when these samples are dried. If the sample contains (or is suspected to contain) > 500 mg/l NH_4 -N fresh basis, as determined by prEN 13651:2001 [4] or prEN 13652:2001 [5], analyze the fresh material as soon as possible.

8 Procedure

8.1 WARNING — Digestions with sulfuric acid are potentially hazardous and laboratory coats, gloves and safety spectacles or goggles shall be used.

The digestion shall be carried out in a well-ventilated fume cupboard with the reflux digestion on a temperature - controlled heating apparatus. Antibumping granules (or roughened glass beads) may be added both to the blank and the samples to prevent bumping and loss of solution. It is important to maintain gentle reflux, both of the blank and the test samples, to avoid temperature fluctuations, which could cause local superheating.

The toxicity or carcinogenicity of each reagent used in this method has not been precisely defined; however, each chemical compound needs to be treated as a potential health hazard. From this viewpoint, reduce exposure to these chemicals to the lowest possible level by whatever means available.

Toxic fumes are evolved by boiling sulfuric acid. Always use the concentrated acid in a fume cupboard.

EN 13654-1:2001 (E)

8.2 Dried sample

Place a portion of the dried ground sample (**clause 7**) of between 0,2 g (expected nitrogen content \approx 0,5 %) and 1,0 g (expected nitrogen content \approx 0,1 %) in the digestion flask (**6.2**). Add 4 ml of salicylic/sulfuric acid (**5.2**) and swirl until the acid is thoroughly mixed with the sample. Allow the mixture to stand for several hours (overnight). Add 0,5 g of sodium thiosulfate (**5.4**) through a dry funnel with a long stem that reaches down into the bulb, or just above the acid if a tube is used. Heat the mixture cautiously on the digestion stand (**6.3**) until frothing has ceased.

Cool the flask; add 1,1 g of the crystalline mixture (**5.3**) and heat until the digestion mixture becomes clear. Boil the mixture gently for up to 5 h so that the sulfuric acid condenses about 1/3 of the way up the neck of the flask. Ensure that the temperature of the solution does not exceed 400 °C.

NOTE In many cases a boiling time of about 2 h is sufficient.

8.3 Residual moisture

The moisture in the analyzed sample (**8.2**) is determined in accordance with EN 13040:1999, clause 10, using 10 g of the dried sample obtained from EN 13040:1999, clause 9.

8.4 Fresh samples

Fresh samples shall be reduced in size to pass a 5 mm sieve or milled in the presence of dry ice (solid carbon dioxide). Weigh up to 20 g sample into a large Kjeldahl flask. Add sufficient salicylic/sulfuric acid (**5.2**) until the acid is thoroughly mixed with the sample. Allow the mixture to stand for several hours (overnight). Add sodium thiosulfate (**5.4**) (0,5 g for every 4 ml of acid used) through a dry funnel with a long stem that reaches down into the bulb. Heat the mixture cautiously on the digestion stand (**6.3**) until frothing has ceased.

Cool the flask, add the crystalline mixture (**5.3**) (1,1 g for every 4 ml of acid used) and heat until the digestion mixture becomes clear. Boil the mixture gently for up to 5 h so that the sulfuric acid condenses about 1/3 of the way up the neck of the flask. Ensure that the temperature of the solution does not exceed 400 °C.

8.5 Determination of ammonium nitrogen in the digestion

The nitrogen in the digest, as ammonium, may be determined by distillation, spectrophotometrically or any other suitable technique (See annex B).

8.6 Moisture content

Determine the moisture content of the fresh sample in accordance with EN 13040:1999, clause 10.

8.7 Blank determination

The reagent blank test shall be carried out in parallel with the determination, by the same procedure, outlined in (**8.2**) and (**8.4**) above using the same quantities of all the reagents as in the determination but omitting the test portion.

NOTE The measurement of a blank is introduced to determine the contribution of the extracting solution and glassware used.

8.8 Calibration

Weigh out a quantity of the calibration standard (**5.5**) equivalent in nitrogen content to the samples being analyzed and proceed as in (**8.2**). The nitrogen determined should exceed 95 % of the calculated value.

8.9 Laboratory compacted bulk density

Determine the laboratory compacted bulk density in accordance with EN 13040:1999, annex A.

9 Calculation

The nitrogen content in milligrams per gram dry matter or milligrams per litre as received basis is calculated according to the final method of determination used.

10 Expression of results

Subtract the values determined for reagent blank from those obtained for the samples. Report all results on a mass/mass basis calculated to a dry matter basis or mass/volume on as received basis.

11 Precision

The repeatability and reproducibility of the nitrogen content in separately prepared samples should be in accordance with Table A.1.

A summary of the results of an interlaboratory trial to determine the precision of the method, in accordance with ISO 5725 [1], is given in annex A.

NOTE The values derived from this interlaboratory trial may not be applicable to concentrations and matrices other than those tested.

12 Test Report

The test report can be prepared separately or in conjunction with the test report of the subsequent analytical method.

The test report shall include the following information:

- a) a reference to the European Standard;
- b) a complete identification of the sample;
- c) all the analytical methods used;
- d) a statement whether ground or un-ground sample has been used;
- e) the results of the determination expressed as mass/mass on dry matter basis for dried samples; for fresh samples on a mass/mass or mass/volume basis as received or calculated to mass/mass dry matter basis;
- f) moisture content;
- g) the laboratory compacted bulk density;
- h) any details not specified in the European Standard, or which are optional, as well as any other factor, which may have affected the results.

Annex A (informative)

Results of an interlaboratory trial to determine nitrogen (modified Kjeldahl method)

An interlaboratory trial was organised in 1997 under the auspices of the European Committee for Standardization, to test the procedures specified in this European Standard.

In this trial the number of laboratories given in Table A.1 determined the nitrogen content using the modified Kjeldahl method in six sample types.

All results are reported on a dry matter basis.

Table A.1 - Summary of the results of the interlaboratory trial for the determination of nitrogen (modified Kjeldahl method)

Sample	No. of labs after Eliminating outliers	No. of outliers (labs)	Mean value Mg/g	Repeatability s_r	Repeatability Std Dev %	Repeatability limit $r = 2,8 s_r$	Reproducibility s_R	Reproducibility Std Dev %	Reproducibility limit $R = 2,8 s_R$
Composted bark	15	0	19,41	0,63	9,11	1,77	2,70	38,98	7,57
Biowaste	15	2	12,88	0,32	7,03	0,91	0,93	20,27	2,61
Clay peat (fertilized)	15	0	9,44	0,31	9,21	0,87	0,66	19,61	1,85
Coarse peat (fertilized)	15	1	11,17	0,22	5,51	0,62	1,10	27,57	3,08
Composted sludge	15	1	29,90	0,96	8,97	2,68	2,97	27,77	8,30
Composted wood fibre	14	1	11,77	0,24	5,69	0,67	0,61	14,62	1,72

Annex B (informative)

Methods of analysis used in the interlaboratory trial

ELEMENT

METHOD

Methods based on or used in the interlaboratory trial

- 1 ISO 11261: 1995 Distillation method [2]
- 2 ISO 11732: 1997 Flow analysis (CFA and FIA) and spectrometric detection [3]

Alternative methods may be suitable for the concentration range and extract used.

Bibliography

- [1] ISO 5725:1994, *Accuracy (trueness and precision) of measurement methods and results.*
- [2] ISO 11261: 1995, *Soil Quality - Determination of total nitrogen - Modified Kjeldahl method.*
- [3] ISO 11732: 1997, *Water quality - Determination of ammonium nitrogen by flow analysis (CFA and FIA) and spectrometric detection.*
- [4] EN 13651:2000, *Soil improvers and growing media - Extraction of calcium chloride/DTPA (CAT) soluble nutrients.*
- [5] EN 13652: 2000, *Soil improvers and growing media - Extraction of water soluble nutrients and elements.*
- [6] Hoffmann, G., 1997 Die Untersuchung von Böden, 4, Auflage, VDLUFA - Verlag, Darmstadt A2.2.1: Bestimmung von Gesamt-Stickstoff nach Kjeldhal (determination of total N according to Kjeldhal).
- [7] Page, A.L. et. al. Methods of soil analysis, Part 2(1982). American Society of Agronomy and Soil Science of America, Madison, WI.
- [8] Williams, P.C. J. Sci. Fd. Agric. 24 (1973), p, 343.

BSI — British Standards Institution

BSI is the independent national body responsible for preparing British Standards. It presents the UK view on standards in Europe and at the international level. It is incorporated by Royal Charter.

Revisions

British Standards are updated by amendment or revision. Users of British Standards should make sure that they possess the latest amendments or editions.

It is the constant aim of BSI to improve the quality of our products and services. We would be grateful if anyone finding an inaccuracy or ambiguity while using this British Standard would inform the Secretary of the technical committee responsible, the identity of which can be found on the inside front cover. Tel: 020 8996 9000. Fax: 020 8996 7400.

BSI offers members an individual updating service called PLUS which ensures that subscribers automatically receive the latest editions of standards.

Buying standards

Orders for all BSI, international and foreign standards publications should be addressed to Customer Services. Tel: 020 8996 9001. Fax: 020 8996 7001. Standards are also available from the BSI website at <http://www.bsi-global.com>.

In response to orders for international standards, it is BSI policy to supply the BSI implementation of those that have been published as British Standards, unless otherwise requested.

Information on standards

BSI provides a wide range of information on national, European and international standards through its Library and its Technical Help to Exporters Service. Various BSI electronic information services are also available which give details on all its products and services. Contact the Information Centre. Tel: 020 8996 7111. Fax: 020 8996 7048.

Subscribing members of BSI are kept up to date with standards developments and receive substantial discounts on the purchase price of standards. For details of these and other benefits contact Membership Administration. Tel: 020 8996 7002. Fax: 020 8996 7001. Further information about BSI is available on the BSI website at <http://www.bsi-global.com>.

Copyright

Copyright subsists in all BSI publications. BSI also holds the copyright, in the UK, of the publications of the international standardization bodies. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI.

This does not preclude the free use, in the course of implementing the standard, of necessary details such as symbols, and size, type or grade designations. If these details are to be used for any other purpose than implementation then the prior written permission of BSI must be obtained.

If permission is granted, the terms may include royalty payments or a licensing agreement. Details and advice can be obtained from the Copyright Manager. Tel: 020 8996 7070.