

BS ISO 7301:2011



BSI Standards Publication

Rice — Specification

bsi.

...making excellence a habit.™

National foreword

This British Standard is the UK implementation of ISO 7301:2011.

The UK participation in its preparation was entrusted to Technical Committee AW/4, Cereals and pulses.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

© BSI 2011

ISBN 978 0 580 62166 6

ICS 67.060

Compliance with a British Standard cannot confer immunity from legal obligations.

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 31 March 2011.

Amendments issued since publication

Date	Text affected
------	---------------

INTERNATIONAL STANDARD

BS ISO 7301:2011

ISO
7301

Third edition
2011-03-01

Rice — Specification

Riz — Spécifications



Reference number
ISO 7301:2011(E)

© ISO 2011

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.



COPYRIGHT PROTECTED DOCUMENT

© ISO 2011

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Contents		Page
Foreword		iv
1	Scope	1
2	Normative references	1
3	Terms and definitions	1
4	Specifications	4
4.1	General, sensory and health specifications	4
4.2	Physical and chemical specifications	5
4.3	Contract specifications	5
5	Test methods	5
6	Packaging	5
Annex A (normative) Methods of analysis for rice specifications		7
Annex B (normative) Determination of waxy rice in parboiled rice		17
Bibliography		19

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 7301 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 4, *Cereals and pulses*.

This third edition cancels and replaces the second edition (ISO 7301:2002), which has been technically revised.

Rice — Specification

1 Scope

This International Standard gives the minimum specifications for rice (*Oryza sativa* L.) which is subject to international trade. It is applicable to the following types: husked rice and milled rice, parboiled or not, intended for direct human consumption. It is neither applicable to other products derived from rice, nor to waxy rice (glutinous rice).

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 712, *Cereals and cereal products — Determination of moisture content — Reference method*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

paddy

paddy rice
rough rice
rice retaining its husk after threshing

3.2

husked rice

brown rice
cargo rice
paddy from which the husk only has been removed

NOTE The processes of husking and handling may result in some loss of bran.

3.3

milled rice

white rice
husked rice from which almost all of the bran and embryo have been removed by milling

3.3.1

undermilled rice

milled rice obtained by milling husked rice, but not to the degree necessary to meet the requirements of well-milled rice

3.3.2

well-milled rice

milled rice obtained by milling husked rice in such a way that most of the bran and part of the embryo have been removed

3.3.3

extra-well-milled rice

milled rice obtained by milling husked rice in such a way that almost all of the bran and the embryo have been removed

3.4

parboiled rice

husked or milled rice processed from paddy or husked rice that has been soaked in water and subjected to a heat treatment so that the starch is fully gelatinized, followed by a drying process

3.5

waxy rice

glutinous rice

varieties of rice whose kernels have a white and opaque appearance

NOTE The starch of waxy rice consists almost entirely of amylopectin. The kernels have a tendency to stick together after cooking.

3.6

whole kernel

husked or milled kernel without any broken part, or part of kernel with a length greater than or equal to nine-tenths of the **average length** (3.12) of the test sample kernels

NOTE See Figure 1.

3.7

head rice

whole kernel (3.6) or part of kernel with a length greater than or equal to three-quarters of the **average length** (3.12) of the test sample kernels

NOTE See Figure 1.

3.8

large broken kernel

part of kernel with a length less than three-quarters but greater than one-half of the **average length** (3.12) of the test sample kernels

NOTE See Figure 1.

3.9

medium broken kernel

part of kernel with a length less than or equal to one-half but greater than one-quarter of the **average length** (3.12) of the test sample kernels

NOTE See Figure 1.

3.10


small broken kernel

part of kernel with a length less than or equal to one-quarter of the **average length** (3.12) of the test sample kernels but which does not pass through test sieve with round apertures having diameter 1,4 mm

NOTE See Figure 1.

**3.11
chip**

part of kernel which passes through a test sieve complying with ISO 5223^[1], and with round apertures having diameter 1,4 mm



Key

- | | |
|------------------------------|--------------------------|
| 1 whole kernel (3.6) | 5 head rice (3.7) |
| 2 large broken kernel (3.8) | 6 broken kernel |
| 3 medium broken kernel (3.9) | 7 chips (3.11) |
| 4 small broken kernel (3.10) | \bar{L} average length |

^a Not passing through a test sieve with round apertures having diameter 1,4 mm.

Figure 1 — Size of kernels, broken kernels and chips

**3.12
average length**

\bar{L}

arithmetic mean of the length of the test sample kernels that are not immature or malformed and without any broken parts

NOTE 1 See definition of **immature kernel** or **malformed kernel** (3.16)

NOTE 2 Calculation of the average length is given in A.4.3.2.

**3.13
extraneous matter**

inorganic and organic components other than whole or broken kernels of rice

3.13.1

inorganic extraneous matter

inorganic components, such as stone, sand and dust

3.13.2

organic extraneous matter

extraneous matter including edible and non-edible

3.13.2.1

edible organic extraneous matter

extraneous matter, such as bran, non-toxic foreign seeds, flour lumps, and other food

3.13.2.2

non-edible organic extraneous matter

extraneous matter, including husks, pieces of straw, and impurities of animal origin, such as dead insects and their fragments

3.14

heat-damaged kernel

head rice or broken kernel that has changed its normal colour as a result of microbiological heating

NOTE This category includes kernel that is yellow to dark yellow in the case of non-parboiled rice and orange to dark orange in the case of parboiled rice, due to a microbiological alteration.

3.15

damaged kernel

head rice or broken kernel showing evident deterioration due to moisture, pests, disease or other causes, but excluding **heat-damaged kernels** (3.14)

3.16

immature kernel

malformed kernel

head rice or broken kernel which is unripe or badly developed

3.17

chalky kernel

head rice or broken kernel of non-parboiled rice, except **waxy rice** (3.5), whose whole surface has an opaque and floury appearance

3.18

red kernel

head rice or broken kernel having a red bran covering more than one-quarter of its surface

3.19

red-streaked kernel

head rice or broken kernel with red bran streaks of length greater than or equal to one-half of the **average length** (3.12), but where the surface covered by these red streaks is less than one-quarter of the total surface

3.20

partly gelatinized kernel

head rice or broken kernel of parboiled rice which is not fully gelatinized and shows a distinct white opaque area

3.21

peck

head rice or broken kernel of parboiled rice of which more than one-quarter of the surface is dark brown or black in colour due to the parboiling process

4 Specifications

4.1 General, sensory and health specifications

Kernels of rice, husked or milled, broken or not, shall be sound, clean and free from foreign odours or odour which indicates deterioration. They shall also be free from toxic or any harmful matter.

The level of additives and pesticides and other contaminants shall not exceed the maximum limits permitted by the national regulations of the country of destination or, in their absence, by CAC/MRL 1^[8] and associated database, CAC/MRL 2^[9], and CAC/MRL 3^[10] (developed by FAO/WHO Codex Alimentarius).

Living insects which are visible to the naked eye shall not be present.

4.2 Physical and chemical specifications

4.2.1 The mass fraction of moisture shall be not greater than 15,0 %.

NOTE Lower moisture contents can be required for certain destinations depending on the climate, duration of transport and storage. For further details, see ISO 6322-1^[2], ISO 6322-2^[3], and ISO 6322-3^[4].

4.2.2 The physical specifications shall be determined in accordance with the method specified in Annex A and shall not exceed the limits given in Table 1.

4.3 Contract specifications

All commercial contracts shall show clearly the:

- a) total percentage of broken kernels permitted, classified according to the agreed categories, and the relative proportion of each category;
- b) total percentage permitted, not exceeding the maximum values for the specifications detailed in Table 1, determined in accordance with the method described in Annex A.

If the contract deals with a specific kind of rice or a specific variety of rice, in order to evaluate the homogeneity of the lot, the contract can specify both the average length and its related coefficient of variation, determined according to A.4.3.2 and A.4.3.3, respectively.

Specifications shall be determined in accordance with the method described in Annex A.

5 Test methods

The moisture content shall be determined in accordance with ISO 712.

The other tests shall be carried out using the methods specified in Annexes A and B.

6 Packaging

The packaging material shall not transmit any smell or taste and shall not contain substances which may damage the product or constitute a health risk. If bags are used, they shall be clean, sufficiently strong and well stitched or sealed.

Table 1 — Contract specifications

Specification (definition reference)	Husked rice non-parboiled	Milled rice non-parboiled	Husked rice parboiled	Milled rice parboiled
Inorganic extraneous matter (3.13.1), % mass fraction	0,5	0,5	0,5	0,5
Organic extraneous matter (3.13.2), % mass fraction	1,0	0,5	1,0	0,5
Paddy (3.1), % mass fraction	2,5	0,3	2,5	0,3
Husked rice non-parboiled (3.2), % mass fraction	—	1,0	1,0	1,0
Milled rice non-parboiled (3.3), % mass fraction	1,0	—	1,0	1,0
Husked rice parboiled (3.2, 3.4), % mass fraction	1,0	1,0	—	1,0
Milled rice parboiled (3.3, 3.4), % mass fraction	1,0	1,0	1,0	—
Chip (3.11), % mass fraction	0,1	0,1	0,1	0,1
Heat-damaged kernel (3.14), % mass fraction	2,0 ^a	2,0	2,0 ^a	2,0
Damaged kernel (3.15), % mass fraction	4,0	3,0	4,0	3,0
Immature or malformed kernel (3.16), % mass fraction	8,0	2,0	8,0	2,0
Chalky kernel (3.17), % mass fraction	5,0 ^a	5,0	—	—
Red kernel and red-streaked kernel (3.18, 3.19), % mass fraction	12,0 ^b	12,0	12,0 ^b	12,0
Partly gelatinized kernel (3.20), % mass fraction	—	—	11,0 ^a	11,0
Peck (3.21), % mass fraction	—	—	4,0	2,0
Waxy rice (3.5), % mass fraction	1,0 ^a	1,0	1,0 ^a	1,0
— Not applicable.				
^a After milling.				
^b Only full red husked (cargo) rice is considered here.				

Annex A (normative)

Methods of analysis for rice specifications

A.1 Principle

Manual separation and weighing of the broken kernels and of the categories in Table 1.

A.2 Apparatus

A.2.1 Sample divider, conical sampler or multiple-slot sampler with a distribution system.

EXAMPLE Sample divider specified in ISO 24333^[6].

A.2.2 Test sieve, with round apertures of diameter 1,4 mm, ISO 5223^[1].

A.2.3 Tweezers, scalpel, and paintbrush.

A.2.4 Small bowls.

A.2.5 Balance, capable of being read to the nearest 0,01 g.

A.2.6 Tray, or other means, coloured in contrast with the colour of the rice to be evaluated.

A.2.7 Micrometer, or other measuring device not deforming the kernels, capable of being read to the nearest 0,01 mm.

A.3 Sampling

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 24333^[6].

It is important the laboratory receive a truly representative sample which has not been damaged or changed during transport or storage.

A.4 Procedure

A.4.1 General

Note if an odour, particular or foreign to rice, is detected, as well as the presence of all anomalies.

Verify the presence of living or dead insects by visual examination and report their number.

A.4.2 Preparation of the test sample

Weigh and carefully mix the laboratory sample to make it as homogeneous as possible. Then reduce it, if necessary, using a sample divider (A.2.1) to obtain a test sample of about 800 g.

Divide the obtained test sample into two equal test portions of about 400 g, using the sample divider (A.2.1).

A.4.3 Determination

A.4.3.1 General

When a kernel or a part of kernel can be classified in more than one category with reference to Table 1, it shall be classified in the category where the maximum permissible value is the lowest. All parts of kernels which get stuck in the apertures of a test sieve shall be considered as being retained by the test sieve.

A.4.3.2 Average length (3.12)

On one of the two test portions (A.4.2):

- a) separate two sets of 100 kernels without any broken part, by random sampling;
- b) measure the length of the kernels using the micrometer (A.2.7) and calculate the arithmetic means of the length for both sets of kernels (\bar{L}_1 and \bar{L}_2);
- c) calculate the average length \bar{L} (3.12) of the two sets of kernels $(\bar{L}_1 + \bar{L}_2)/2$; if the value of $100(\bar{L}_1 - \bar{L}_2)/\bar{L}$ is higher than 2, return all the kernels to the tray and repeat from step a);
- d) return all the kernels to the test portion.

A.4.3.3 Coefficient of variation of the length

From the measures of kernels in A.4.3.2 b), calculate the coefficient of variation as follows.

The values are:

$$L_1, L_2 \dots L_i \dots L_n$$

where L_i is the length of a single rice kernel.

The arithmetic mean is given by:

$$\bar{L} = \frac{\sum_{i=1}^n L_i}{n}$$

The standard deviation is given by:

$$s = \sqrt{\frac{\sum_{i=1}^n L_i^2 - \frac{\left(\sum_{i=1}^n L_i\right)^2}{n}}{n-1}}$$

The coefficient of variation, expressed as a percentage, is given by

$$C_V = \frac{s}{\bar{L}} \times 100 \%$$

Usually, the coefficient of variation of an homogeneous lot of rice does not exceed a value of 5 %.

A.4.3.4 Husked rice, non-parboiled (see Figure A.1)

Weigh one of the test portions (A.4.2) to the nearest 0,1 g, record the mass as m_w , and spread it on the tray (A.2.6). Separate the inorganic extraneous matter (3.13.1), the organic extraneous matter (3.13.2; 3.13.2.1 and 3.13.2.2), the paddy (3.1), the milled rice non-parboiled (3.3), the husked rice parboiled (3.2 and 3.4) and the milled rice parboiled (3.3 and 3.4), into small bowls (A.2.4), with the aid of tweezers, scalpel and paintbrush (A.2.3). Weigh the six fractions obtained to the nearest 0,01 g, and record the masses as m_1 , m_2 , m_3 , m_4 , m_5 , and m_6 .

Divide the second test portion with the divider (A.2.1) in order to obtain four different aliquot parts of about 100 g each.

Weigh the first aliquot part to the nearest 0,01 g and record the mass as m_x . Spread it out and separate the damaged kernels (3.15), the immature or malformed kernels (3.16) and the red kernels (3.18) into small bowls. Weigh the three fractions obtained to the nearest 0,01 g, and record the masses as m_7 , m_8 , and m_9 .

Weigh the second aliquot part to the nearest 0,01 g and record the mass as m_y . Separate the chips (3.11) by the test sieve (A.2.2), then spread out the remainder and separate the broken kernels, classifying them into large broken kernels (3.8), medium broken kernels (3.9), and small broken kernels (3.10). Put the fractions obtained into small bowls. Weigh the four fractions to the nearest 0,01 g and record the masses as m_{10} , m_{11} , m_{12} , and m_{13} .

Proceed with the laboratory milling of the third aliquot part. Weigh the obtained milled rice to the nearest 0,01 g and record the mass as m_z . Spread it out and separate the heat-damaged kernels (3.14), the chalky kernels (3.17) and waxy rice (3.5) into small bowls. Weigh the three fractions obtained to the nearest 0,01 g and record the masses as m_{14} , m_{15} , and m_{16} .

A.4.3.5 Milled rice, non-parboiled (see Figure A.2)

Weigh one of the test portions (A.4.2) to the nearest 0,1 g, record the mass as m_w , and spread it on the tray (A.2.6). Separate the inorganic extraneous matter (3.13.1), the organic extraneous matter (3.13.2; 3.13.2.1 and 3.13.2.2), the paddy (3.1), the husked rice non-parboiled (3.2), the husked rice parboiled (3.2 and 3.4) and the milled rice parboiled (3.3 and 3.4) into small bowls (A.2.4), with the aid of tweezers, scalpel and paintbrush (A.2.3). Weigh the six fractions obtained to the nearest 0,01 g and record the masses as m_1 , m_2 , m_3 , m_4 , m_5 , and m_6 .

Divide the second test portion with the divider (A.2.1) in order to obtain four different aliquot parts of about 100 g each.

Weigh the first aliquot part to the nearest 0,01 g and record the mass as m_x . Spread it out and separate the heat-damaged kernels (3.14), the damaged kernels (3.15), the immature or malformed kernels (3.16), the chalky kernels (3.17), the red kernels (3.18), together with the red-streaked kernels (3.19) and the waxy rice (3.5) into small bowls. Weigh the six fractions obtained to the nearest 0,01 g and record the masses as m_7 , m_8 , m_9 , m_{10} , m_{11} , and m_{16} .

Weigh the second aliquot part to the nearest 0,01 g and record the mass as m_y . Separate the chips (3.11) by the test sieve (A.2.2), then spread out the remainder and separate the broken kernels, classifying them into large broken kernels (3.8), medium broken kernels (3.9) and small broken kernels (3.10). Put the fractions obtained into small bowls. Weigh the four fractions to the nearest 0,01 g and record the masses as m_{12} , m_{13} , m_{14} , and m_{15} .

A.4.3.6 Husked rice, parboiled (see Figure A.3)

Weigh one of the test portions (A.4.2) to the nearest 0,1 g, record the mass as m_w , and spread it on the tray (A.2.6). Separate the inorganic extraneous matter (3.13.1), the organic extraneous matter (3.13.2; 3.13.2.1 and 3.13.2.2), the paddy (3.1), the husked rice non-parboiled (3.2), the milled rice non-parboiled (3.3) and the milled rice parboiled (3.3 and 3.4) into small bowls (A.2.4) with the aid of tweezers, scalpel and paintbrush

(A.2.3). Weigh the six fractions obtained to the nearest 0,01 g and record the masses as m_1 , m_2 , m_3 , m_4 , m_5 , and m_6 .

Divide the second test portion with the divider (A.2.1) in order to obtain four different aliquot parts of about 100 g each.

Weigh the first aliquot part to the nearest 0,01 g and record the mass as m_x . Spread it out and separate the damaged kernels (3.15), the immature or malformed kernels (3.16) and the red kernels (3.18) into small bowls. Weigh the three fractions obtained to the nearest 0,01 g and record the masses as m_7 , m_8 , and m_9 .

Weigh the second aliquot part to the nearest 0,01 g and record the mass as m_y . Separate the chips (3.11) by the test sieve (A.2.2), then spread out the remainder and separate the broken kernels, classifying them into large broken kernels (3.8), medium broken kernels (3.9) and small broken kernels (3.10). Put the four fractions obtained into small bowls. Weigh the fractions to the nearest 0,01 g and record the masses as m_{10} , m_{11} , m_{12} , and m_{13} .

Proceed with the laboratory milling of the third aliquot part. Weigh the milled rice to the nearest 0,01 g and record the mass as m_z . Spread it out and separate the heat-damaged kernels (3.14), the partly gelatinized kernels (3.20) and the pecks (3.21) into small bowls. Weigh the three fractions obtained to the nearest 0,01 g and record the masses as m_{14} , m_{15} , and m_{16} .

Proceed with the milling of the fourth aliquot part and determine the percentage of waxy rice (3.5) according to Annex B.

A.4.3.7 Milled rice, parboiled (see Figure A.4)

Weigh one of the test portions (A.4.2) so obtained to the nearest 0,1 g, record the mass as m_w , and spread it on the tray (A.2.6). Separate the inorganic extraneous matter (3.13.1), organic extraneous matter (3.13.2; 3.13.2.1 and 3.13.2.2), the paddy (3.1), the husked rice non-parboiled (3.2), the milled rice non-parboiled (3.3) and the husked rice parboiled (3.3 and 3.4) into small bowls (A.2.4), with the aid of tweezers, scalpel and paintbrush (A.2.3). Weigh the six fractions obtained to the nearest 0,01 g and record the masses as m_1 , m_2 , m_3 , m_4 , m_5 , and m_6 .

Divide the second test portion with the divider (A.2.1) in order to obtain four different aliquot parts of about 100 g each.

Weigh the first aliquot part to the nearest 0,01 g and record the mass as m_x . Spread it out and separate the heat-damaged kernels (3.14), the damaged kernels (3.15), the immature or malformed kernels (3.16), the red kernels (3.18), together with the red streaked kernels (3.19), the partly gelatinized kernels (3.20) and the pecks (3.21) into small bowls. Weigh the six fractions obtained to the nearest 0,01 g and record the masses as m_7 , m_8 , m_9 , m_{10} , m_{11} , and m_{12} .

Weigh the second aliquot part to the nearest 0,01 g and record the mass as m_y . Separate the chips (3.11) by the test sieve (A.2.2), then spread out the remainder and separate the broken kernels, classifying them into large broken kernels (3.8), medium broken kernels (3.9) and small broken kernels (3.10). Put the fractions obtained into small bowls. Weigh the four fractions to the nearest 0,01 g and record the masses as m_{13} , m_{14} , m_{15} , and m_{16} .

Weigh the third aliquot part to the nearest 0,01 g and determine the percentage of waxy rice (3.5) according to Annex B.

A.5 Expression of results

Report the result obtained for the categories given in Table A.1 as a mass fraction, expressed as a percentage, of the product as received.

Table A.1 — Expression of results

Categories	Husked rice non-parboiled ^a	Milled rice non-parboiled ^b	Husked rice parboiled ^c	Milled rice parboiled ^d
Inorganic extraneous matter (3.13.1)	$\frac{m_1 \times 100}{m_w}$	$\frac{m_1 \times 100}{m_w}$	$\frac{m_1 \times 100}{m_w}$	$\frac{m_1 \times 100}{m_w}$
Organic extraneous matter (3.13.2; 3.13.2.1; 3.13.2.2)	$\frac{m_2 \times 100}{m_w}$	$\frac{m_2 \times 100}{m_w}$	$\frac{m_2 \times 100}{m_w}$	$\frac{m_2 \times 100}{m_w}$
Paddy (3.1)	$\frac{m_3 \times 100}{m_w}$	$\frac{m_3 \times 100}{m_w}$	$\frac{m_3 \times 100}{m_w}$	$\frac{m_3 \times 100}{m_w}$
Husked rice non-parboiled (3.2)	—	$\frac{m_4 \times 100}{m_w}$	$\frac{m_4 \times 100}{m_w}$	$\frac{m_4 \times 100}{m_w}$
Milled rice non-parboiled (3.3)	$\frac{m_4 \times 100}{m_w}$	—	$\frac{m_5 \times 100}{m_w}$	$\frac{m_5 \times 100}{m_w}$
Husked rice parboiled (3.2 and 3.4)	$\frac{m_5 \times 100}{m_w}$	$\frac{m_5 \times 100}{m_w}$	—	$\frac{m_6 \times 100}{m_w}$
Milled rice parboiled (3.3 and 3.4)	$\frac{m_6 \times 100}{m_w}$	$\frac{m_6 \times 100}{m_w}$	$\frac{m_6 \times 100}{m_w}$	—
Heat-damaged kernels (3.14)	$\frac{m_{14} \times 100}{m_z}$	$\frac{m_7 \times 100}{m_x}$	$\frac{m_{14} \times 100}{m_z}$	$\frac{m_7 \times 100}{m_x}$
Damaged kernels (3.15)	$\frac{m_7 \times 100}{m_x}$	$\frac{m_8 \times 100}{m_x}$	$\frac{m_7 \times 100}{m_x}$	$\frac{m_8 \times 100}{m_x}$
Immature or malformed kernel (3.16)	$\frac{m_8 \times 100}{m_x}$	$\frac{m_9 \times 100}{m_x}$	$\frac{m_8 \times 100}{m_x}$	$\frac{m_9 \times 100}{m_x}$
Chalky kernel (3.17)	$\frac{m_{15} \times 100}{m_z}$	$\frac{m_{10} \times 100}{m_x}$	—	—
Partly gelatinized kernel (3.20)	—	—	$\frac{m_{15} \times 100}{m_z}$	$\frac{m_{11} \times 100}{m_x}$
Peck (3.21)	—	—	$\frac{m_{16} \times 100}{m_z}$	$\frac{m_{12} \times 100}{m_x}$
Red and red streaked kernel (3.18) and (3.19)	$\frac{m_9 \times 100}{m_x}$	$\frac{m_{11} \times 100}{m_x}$	$\frac{m_9 \times 100}{m_x}$	$\frac{m_{10} \times 100}{m_x}$
Waxy rice (3.5)	$\frac{m_{16} \times 100}{m_z}$	$\frac{m_{16} \times 100}{m_x}$	$\frac{m_{17} \times 100}{m_{17} + m_{18}}$	$\frac{m_{17} \times 100}{m_{17} + m_{18}}$
Large broken kernel (3.8)	$\frac{m_{10} \times 100}{m_y}$	$\frac{m_{12} \times 100}{m_y}$	$\frac{m_{10} \times 100}{m_y}$	$\frac{m_{13} \times 100}{m_y}$
Medium broken kernel (3.9)	$\frac{m_{11} \times 100}{m_y}$	$\frac{m_{13} \times 100}{m_y}$	$\frac{m_{11} \times 100}{m_y}$	$\frac{m_{14} \times 100}{m_y}$
Small broken kernel (3.10)	$\frac{m_{12} \times 100}{m_y}$	$\frac{m_{14} \times 100}{m_y}$	$\frac{m_{12} \times 100}{m_y}$	$\frac{m_{15} \times 100}{m_y}$
Chip (3.11)	$\frac{m_{13} \times 100}{m_y}$	$\frac{m_{15} \times 100}{m_y}$	$\frac{m_{13} \times 100}{m_y}$	$\frac{m_{16} \times 100}{m_y}$
Average length (A.4.3.2)	\bar{L}	\bar{L}	\bar{L}	\bar{L}
Coefficient of variation of the length (A.4.3.3)	C_V	C_V	C_V	C_V
<p>^a For the meaning of the symbols, refer to the scheme of procedure given in Figure A.1. ^b For the meaning of the symbols, refer to the scheme of procedure given in Figure A.2. ^c For the meaning of the symbols, refer to the scheme of procedure given in Figure A.3. ^d For the meaning of the symbols, refer to the scheme of procedure given in Figure A.4.</p>				

Report the result for each category to one decimal place by rounding it to the nearest integral multiple. If there are two successive integral multiples equally near to the given number, the even integral multiple should be selected as the rounded number. Carry out the rounding always in one step.

Although ISO 80000-1^[7] specifies two different rules for rounding if there are two successive integral multiples equally near to the given number, for the purposes of this International Standard the above-mentioned rule (i.e. ISO 80000-1^[7], rule A) should be used, as in this way the rounding errors are minimized.

EXAMPLES

Given number	Rounded number
18,23	18,2
18,26	18,3
18,37	18,4
18,25	18,2
18,35	18,4
18,347	18,3
18,251	18,3

A.6 Test report

The test report shall contain at least the following information:

- a) all information necessary for the complete identification of the sample;
- b) the sampling method used, if known;
- c) the sample mass;
- d) the test method used, with reference to this International Standard (ISO 7301:2011);
- e) the date of analysis;
- f) any operating details not specified in this annex, or regarded as optional, together with details of any incidents likely to have influenced the results;
- g) the test results obtained.




Figure A.1 — Scheme of procedure for husked rice non-parboiled




Figure A.2 — Scheme of procedure for milled rice non-parboiled




Figure A.3 — Scheme of procedure for husked rice parboiled




Figure A.4 — Scheme of procedure for milled rice parboiled

Annex B (normative)

Determination of waxy rice in parboiled rice

B.1 Principle

Waxy rice kernels have a reddish brown colour when stained in an iodine solution, while non-waxy rice kernels show a blue colour.

B.2 Reagents and materials

During the analysis, unless otherwise stated, use only reagents of recognized analytical grade and distilled or demineralized water or water of equivalent purity.

B.2.1 Iodine stock solution, containing 0,2 g iodine and 2,0 g potassium iodide in 100 ml water.

B.2.2 Iodine working solution, dilute the stock solution (B.2.1) two times (by volume) with water. Prepare daily.

B.2.3 Tissue paper.

B.3 Apparatus

B.3.1 Balance, capable of being read to the nearest 0,01 g.

B.3.2 Glass beaker, capacity 250 ml.

B.3.3 Small white-coloured bowls, or any white-coloured container.

B.3.4 Wire basket.

B.3.5 Stirrer rod.

B.3.6 Tweezers or forceps.

B.4 Sampling

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 24333^[6].

It is important the laboratory receive a truly representative sample which has not been damaged or changed during transport or storage.

B.5 Determination

B.5.1 Weigh a portion of about 100 g milled rice and put it into a glass beaker (B.3.2).

B.5.2 Add approximately 80 ml of iodine working solution (B.2.2) to soak the kernels and stir until all the kernels are submerged under the solution. Let the kernels soak in the solution for 30 s.

B.5.3 Pour the rice and solution into a wire basket (B.3.4) and shake the basket slightly in order to drain out the solution. Then place the basket on a piece of tissue paper (B.2.3) to absorb the excess liquid.

B.5.4 Pour the stained kernels into a bowl (B.3.3). Separate the reddish brown kernels of waxy rice from the dark blue kernels of non-waxy rice.

B.5.5 Weigh the waxy rice portion, m_{17} , and the non-waxy rice portion, m_{18} , to the nearest 0,1 g.

B.5.6 Calculate the waxy rice content, w , expressed as a percentage mass fraction, from the equation:

$$w = \frac{m_{17}}{m_{17} + m_{18}} \times 100 \%$$

Bibliography

- [1] ISO 5223, *Test sieves for cereals*
- [2] ISO 6322-1, *Storage of cereals and pulses — Part 1: General recommendations for the keeping of cereals*
- [3] ISO 6322-2, *Storage of cereals and pulses — Part 2: Practical recommendations*
- [4] ISO 6322-3, *Storage of cereals and pulses — Part 3: Control of attack by pests*
- [5] ISO 6646, *Rice — Determination of the potential milling yield from paddy and from husked rice*
- [6] ISO 24333, *Cereals and cereal products — Sampling*
- [7] ISO 80000-1, *Quantities and units — Part 1: General*
- [8] CAC/MRL 1, *Maximum residue limits (MRLs) for pesticides*. A database is available (viewed 2010-09-29) at: <http://www.codexalimentarius.net/pestres/data/pesticides/index.html>
- [9] CAC/MRL 2, *Maximum residue limits for veterinary drugs in food*
- [10] CAC/MRL 3, *Extraneous maximum residue limits (EMRLs)*

British Standards Institution (BSI)

BSI is the national body responsible for preparing British Standards and other standards-related publications, information and services.

BSI is incorporated by Royal Charter. British Standards and other standardization products are published by BSI Standards Limited.

About us

We bring together business, industry, government, consumers, innovators and others to shape their combined experience and expertise into standards-based solutions.

The knowledge embodied in our standards has been carefully assembled in a dependable format and refined through our open consultation process. Organizations of all sizes and across all sectors choose standards to help them achieve their goals.

Information on standards

We can provide you with the knowledge that your organization needs to succeed. Find out more about British Standards by visiting our website at bsigroup.com/standards or contacting our Customer Services team or Knowledge Centre.

Buying standards

You can buy and download PDF versions of BSI publications, including British and adopted European and international standards, through our website at bsigroup.com/shop, where hard copies can also be purchased.

If you need international and foreign standards from other Standards Development Organizations, hard copies can be ordered from our Customer Services team.

Subscriptions

Our range of subscription services are designed to make using standards easier for you. For further information on our subscription products go to bsigroup.com/subscriptions.

With **British Standards Online (BSOL)** you'll have instant access to over 55,000 British and adopted European and international standards from your desktop. It's available 24/7 and is refreshed daily so you'll always be up to date.

You can keep in touch with standards developments and receive substantial discounts on the purchase price of standards, both in single copy and subscription format, by becoming a **BSI Subscribing Member**.

PLUS is an updating service exclusive to BSI Subscribing Members. You will automatically receive the latest hard copy of your standards when they're revised or replaced.

To find out more about becoming a BSI Subscribing Member and the benefits of membership, please visit bsigroup.com/shop.

With a **Multi-User Network Licence (MUNL)** you are able to host standards publications on your intranet. Licences can cover as few or as many users as you wish. With updates supplied as soon as they're available, you can be sure your documentation is current. For further information, email bsmusales@bsigroup.com.

BSI Group Headquarters

389 Chiswick High Road London W4 4AL UK

Revisions

Our British Standards and other publications are updated by amendment or revision.

We continually improve the quality of our products and services to benefit your business. If you find an inaccuracy or ambiguity within a British Standard or other BSI publication please inform the Knowledge Centre.

Copyright

All the data, software and documentation set out in all British Standards and other BSI publications are the property of and copyrighted by BSI, or some person or entity that owns copyright in the information used (such as the international standardization bodies) and has formally licensed such information to BSI for commercial publication and use. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI. Details and advice can be obtained from the Copyright & Licensing Department.

Useful Contacts:

Customer Services

Tel: +44 845 086 9001

Email (orders): orders@bsigroup.com

Email (enquiries): cservices@bsigroup.com

Subscriptions

Tel: +44 845 086 9001

Email: subscriptions@bsigroup.com

Knowledge Centre

Tel: +44 20 8996 7004

Email: knowledgecentre@bsigroup.com

Copyright & Licensing

Tel: +44 20 8996 7070

Email: copyright@bsigroup.com



...making excellence a habit.™