

Soil quality — Sampling

**Part 6: Guidance on the collection,
handling and storage of soil under
aerobic conditions for the assessment
of microbiological processes, biomass
and diversity in the laboratory**

ICS 13.080.05

National foreword

This British Standard is the UK implementation of ISO 10381-6:2009. It supersedes BS 7755 SEC2.6:1994 which is withdrawn.

The UK participation in its preparation was entrusted to Technical Committee EH/4, Soil quality.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with a British Standard cannot confer immunity from legal obligations.

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 28 February 2010.

© BSI 2010

ISBN 978 0 580 58670 5

Amendments/corrigenda issued since publication

Date	Comments

INTERNATIONAL STANDARD

ISO
10381-6

Second edition
2009-03-15

Soil quality — Sampling —

Part 6:

Guidance on the collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory

Qualité du sol — Échantillonnage —

Partie 6: Lignes directrices pour la collecte, la manipulation et la conservation, dans des conditions aérobies, de sols destinés à l'évaluation en laboratoire des processus, de la biomasse et de la diversité microbiens



Reference number
ISO 10381-6:2009(E)

© ISO 2009

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.



COPYRIGHT PROTECTED DOCUMENT

© ISO 2009

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Contents

Page

Foreword	iv
Introduction	v
1 Scope	1
2 Terms and definitions	1
3 Procedure	1
3.1 Selection of sampling locations	1
3.2 Description of field site	2
3.3 Sampling conditions	2
3.4 Sampling methods	2
3.5 Sample marking	2
3.6 Transportation conditions	2
3.7 Soil processing	2
3.8 Storage conditions and storage periods	3
3.9 Pre-incubation	4
4 Sampling report	4
Bibliography	6

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 10381-6 was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

This second edition cancels and replaces the first edition (ISO 10381-6:1993), Subclauses 3.6, 3.7, 3.8 and Clause 4 of which have been technically revised. Table 1 has been added.

ISO 10381 consists of the following parts, under the general title *Soil quality — Sampling*:

- *Part 1: Guidance on the design of sampling programmes*
- *Part 2: Guidance on sampling techniques*
- *Part 3: Guidance on safety*
- *Part 4: Guidance on the procedure for investigation of natural, near-natural and cultivated sites*
- *Part 5: Guidance on the procedure for the investigation of urban and industrial sites with regard to soil contamination*
- *Part 6: Guidance on the collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory*
- *Part 7: Guidance on sampling of soil gas*
- *Part 8: Guidance on sampling of stockpiles*

Introduction

Soils are both complex and heterogeneous because they consist of both living and non-living components occurring in different combinations. Therefore, the condition of the soil, from collection to completion of an experiment, should be considered in relation to effects on the soil microflora. Temperature, water content, availability of oxygen and duration of storage are all known to affect the soil microflora, and thus the processes they mediate.

Soils can however be used effectively in laboratory systems to investigate microbially-mediated processes, provided that the dynamics of the living microflora are appreciated. This part of ISO 10381 provides guidance on the collection, handling and storage of soil for laboratory use where aerobic microbial activity is the main component of the study. It describes how to minimize the effects of differences in temperature, water content and availability of oxygen on aerobic processes to facilitate reproducible laboratory determinations [10], [11].

Soil quality — Sampling —

Part 6:

Guidance on the collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory

1 Scope

This part of ISO 10381 provides guidance on the collection, handling and storage of soil for subsequent testing under aerobic conditions in the laboratory. The recommendations in this document are not applicable to the handling of soil where anaerobic conditions are to be maintained throughout.

This part of ISO 10381 is mainly applicable to temperate soils. Soils collected from extreme climates (e.g. permafrost, tropical soils) may require special handling.

2 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

2.1

aerobic

descriptive of a condition in which molecular oxygen is freely available

2.2

anaerobic

descriptive of a condition in which molecular oxygen is not available

2.3

water content on a dry mass basis

mass of water evaporating from the soil when dried to constant mass at 105 °C divided by the dry mass of the soil and multiplied by 100

[ISO 11465:1993, 3.2]

3 Procedure

3.1 Selection of sampling locations

The locations of the sites from which samples are taken should be selected according to the purpose of the study.

These locations should be identified and recorded, e.g. on a map by reference to easily recognizable static objects or by using a detailed map reference or by GIS. If practicable, the locations should be marked so that they may be used for comparative tests or for obtaining replicate samples.

3.2 Description of field site

Selection of a soil sampling site depends on the purpose of a particular study, and knowledge of the field site history is always desirable. The site should be accurately described and its history given. Details of vegetation cover, the morphology of the sampling area (e.g. flat area, slopes, steepness), and of chemical and biological additions or accidental contamination, should be recorded and reported.

3.3 Sampling conditions

Soil required for studies conducted under laboratory conditions should, if practicable, be sampled in the field with a soil water content which facilitates sieving. Sampling should, unless the purpose of the study requires otherwise, be avoided during or immediately following long periods (e.g. 1 month) of drought, freezing or flooding. If laboratory tests are to be used for field monitoring, conditions existing in the field should be accepted. Soil samples may be frozen before investigations of, for example, ammonium oxidations.

3.4 Sampling methods

The sampling technique depends on the purpose of the study. If aerobic agricultural soil is required, sampling is usually conducted to the actual ploughing depth. Any surface vegetation cover, moss-covered litter layer, visible roots, large pieces of plant or woody plant litter and visible soil fauna should be removed to minimize the addition of fresh organic carbon to the soil. Organic constituents introduced from roots or other sources can cause unpredictable changes in the activities and composition of the soil microflora. If natural soils show distinct horizons, samples should be taken from these horizons.

3.5 Sample marking

Sample containers should be clearly and unambiguously marked and identified so that each sample can be related to the location from which it was taken. Use of containers which either absorb water from the soil or release materials, e.g. solvents or plasticizers, into the soil should be avoided.

3.6 Transportation conditions

Samples should be transported in a manner which minimizes changes in the soil water content, and should be kept in the dark with free access of air; a loosely-tied polyethylene bag is generally adequate for this purpose. Extreme environmental conditions should be avoided: the soil should be kept as cool as possible but it is essential that it is not allowed to dry out or become water-logged. Exposure to light for extended periods should be avoided as this encourages the growth of algae on the surface of the soil. Physical compaction should be avoided as far as is practicable.

Samples for DNA or RNA analysis shall be frozen quickly in the field using dry ice. During transportation to the laboratory, dry ice shall be used to maintain the temperature of those for RNA analysis. Samples for DNA analysis may be transported in a cooling box unless the circumstances are such that dry ice is needed for these as well.

3.7 Soil processing

The soil should be processed as soon as possible after sampling. Vegetation, larger soil fauna and stones should be removed prior to passing the soil through a 2 mm sieve. Sieving soil through a 2 mm sieve facilitates gaseous exchange between particles and is therefore recommended for maintaining the aerobic nature of the soil. It also removes small stones, fauna and plant debris. Some organic materials such as moor layers or peat do not pass easily through a 2 mm sieve and should be sieved in the moist condition through a 5 mm sieve. This necessitates manual operation and the quality of the material passing the sieve depends on the operator. If the soil is too wet to sieve, it should be spread out, in a gentle air stream where possible, to facilitate uniform drying. The soil should be finger crumbled and turned over frequently to avoid excessive surface drying. Usually, this should be performed at ambient temperature. If drying is required, the soil should not be dried more than necessary to facilitate sieving. Generally, drying of soils is not recommended although air-drying and rewetting is a common physiological stress for the microbial communities in surface soils. It has been shown that drying-rewetting events can induce significant changes in microbial C and N dynamics which

can last for more than a month after the last stress [12]. Rewetting after drying causes bursts of respiration and growth of distinct populations of bacteria [15]. If further storage is necessary following processing, consideration should be given to the parameters discussed in 3.8.

3.8 Storage conditions and storage periods

Samples should be stored in the dark at $(4 \pm 2)^\circ\text{C}$ with free access of air. A loosely tied plastic bag or equivalent is generally adequate for this purpose. Care should be taken to ensure that the soil is not stored in a quantity which allows anaerobic conditions to occur in the bottom of storage containers. The soil should be processed (see 3.7) before storage in order to ensure stable aerobic conditions. It is essential that the soil is not allowed to dry out or become water-logged during storage. Samples should not be stored on top of one another. It is preferable to use soils as soon as possible after sampling. Any delays due to transportation should be minimized. If storage is unavoidable, this should not exceed three months unless there is evidence showing continued microbial activity. Soil samples subject to DNA analyses should be frozen at -20°C if not processed immediately. For RNA analyses, samples should be frozen at -80°C .

If soil samples need to be stored for longer periods (> three months), freezing of samples at -20°C , -80°C or -180°C may be appropriate although not generally recommended. Examples for appropriate storage conditions for various test objectives are given in Table 1. It has been shown for a number of soils from temperate climates that storage at -20°C for up to 12 months does not inhibit microbial activity (e.g. ammonium oxidation). Moreover, soil samples for Phospholipid Fatty Acid (PLFA) and DNA analyses can be stored at -20°C for one to two years. Samples for rRNA analyses can be stored at -80°C for the same period. Shock freezing with liquid nitrogen is recommended for freezing samples subjected to DNA, RNA and PLFA/PLEL analyses.

Longer storage periods are mainly needed if the influence of added pollutants on soil microbes and microbial processes is tested with the same soil material or if the community structure (PLFA, DNA, RNA) of soils is evaluated at a distinct point of time during the year. In these cases, the time needed for analyses can easily exceed three months (chemical, pollutant testing). For structural analyses of the microflora, storage at 4°C is not suitable.

Table 1 — Storage conditions for the assessment of aerobic microbial processes when analysis cannot be performed immediately

Test objective	International Standard	Wet 4 °C months	Wet –20 °C ^a years	Wet –80 °C or liquid nitrogen (–180 °C) ^a years
PLFA, PLEL		—	2	10
DNA		—	2	10
RNA		—	—	10
Biomass				
– Substrate-induced respiration method	ISO 14240-1	3	1	10
– Fumigation-extraction method	ISO 14240-2			
Potential ammonium oxidation	ISO 15685	3	1	10
Nitrogen mineralization	ISO 14238	3	1	10
Microbial soil respiration	ISO 16072	3	1	10
Soil respiration curves	ISO 17155	3	1	10
Dehydrogenase activity	ISO 23753-1 ISO 23753-2	3	1	—

symbols: — = no storage possible.

^a The soils shall be divided into sub-samples for further investigation before storage. Alternatively, PLFA should be extracted from field-moist soils (<2 mm) immediately after sampling. The extracts can be stored at –20 °C for several months prior to further separation steps and analysis with GC/MS. Shock freezing in liquid nitrogen is recommended before storage at –20 °C or –80 °C.

3.9 Pre-incubation

Before the processed soil is used for a specific laboratory test, it should be pre-incubated to allow germination and removal of seeds, and to re-establish an equilibrium of microbial metabolism following the change from sampling or storage conditions to incubation conditions. Pre-incubation conditions depend on the purpose of the study but should approach test conditions as far as is practicable. The period of pre-incubation depends on the purpose of the study, the soil composition and the storage/pre-incubation conditions. A period between 2 d and 28 d is generally adequate.

If samples were frozen, special attention shall be given to the thawing of samples. For the analyses of microbial activity (e.g. soil respiration), a thawing period of one week at 4 °C and another three days at 20 °C are recommended. A thawing period of one day at 20 °C might also be suitable. For DNA, RNA and PLFA/PLEL analyses, the thawing period shall be as short as possible to avoid degradation processes.

Freezing the samples can change the water-holding capacity; therefore, for such samples the water-holding capacity should be determined after thawing.

4 Sampling report

The detailed sampling report depends on the sampling objectives but, in general, the following data should be reported:

- a reference to this part of ISO 10381;
- location of the site (sufficiently precise for another person to find it without further guidance);
- a comprehensive description of the relevant details and features of the site;

- d) history of the site, including previous use and any known accidental or intentional chemical or biological additions;
- e) the date and time of sample collection;
- f) the weather conditions at the time or immediately prior to sampling, including air temperature, rainfall, sunshine, cloud, etc.;
- g) the precise location from which the sample was taken;
- h) the type of device used to take the sample;
- i) whether or not the sample needed drying before sieving;
- j) the number of samples taken, the area of sampling plot or sampling area;
- k) the depth of sampling;
- l) whether the individual soil samples were kept separate or pooled into composite samples;
- m) the time elapsed between sampling, transport and post-sampling handling;
- n) any other factor that might influence the results of subsequent testing.

Bibliography

- [1] ISO 11465:1993, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method*
- [2] ISO 14238, *Soil quality — Biological methods — Determination of nitrogen mineralization and nitrification in soils and the influence of chemicals on these processes*
- [3] ISO 14240-1, *Soil quality — Determination of soil microbial biomass — Part 1: Substrate-induced respiration method*
- [4] ISO 14240-2, *Soil quality — Determination of soil microbial biomass — Part 2: Fumigation-extraction method*
- [5] ISO 15685, *Soil quality — Determination of potential nitrification and inhibition of nitrification — Rapid test by ammonium oxidation*
- [6] ISO 16072, *Soil quality — Laboratory methods for determination of microbial soil respiration*
- [7] ISO 17155, *Soil quality — Determination of abundance and activity of soil microflora using respiration curves*
- [8] ISO 23753-1, *Soil quality — Determination of dehydrogenase activity in soils — Part 1: Method using triphenyltetrazolium chloride (TTC)*
- [9] ISO 23753-2, *Soil quality — Determination of dehydrogenase activity in soils — Part 2: Method using iodotetrazolium chloride (INT)*
- [10] ANDERSON, J.P.E. Handling and Storage of Soils and method for measuring the persistence and biological effects of pesticides in soil. *Soil Science*, **100**, 1965, pp. 68-70
- [11] BARTHA, R. and PRAMER, D., *Features of a flask for Pesticide Experiments, Pesticide Effects and Soil Microflora* (eds. Sommerville, L. and Greaves, M.P.), 1987, pp. 45-60, Taylor and Francis
- [12] FIERER, N. and SCHIMEL, J.P. Effects of drying-rewetting frequency on soil carbon and nitrogen transformations. *Soil Biology & Biochemistry*, **34**, 2002, pp. 777-787
- [13] GOBERNA, M., INSAM, H., PASCUAL, J.A. and SANCHEZ, J. Storage effects on the community level physiological profiles of Mediterranean forest soils. *Soil Biology and Biochemistry*, **37**, 2005, pp. 173-178
- [14] HAYNES, R.J. and BEARE, M.H. Aggregation and organic matter storage in meso-thermal, humid soils. In: *Soil Structure and Organic Matter Storage* (eds. Carter, M.R., Steward, B.A.), CRC/Lewis Publishers, Boca Raton, 1996, pp. 213-262
- [15] LUND, V. and GOKSØR, J. Effects of Water Fluctuations on Microbial Mass and Activity in Soil. *Microbial Ecol.*, **6**, 1980, pp. 115-123
- [16] SHISHIDO, M. and CHANWAY, C.P. Storage effect on indigenous microbial communities and PGPR efficacy. *Soil Biology and Biochemistry*, **30**, 1998, pp. 939-947
- [17] STENBERG, B., JOHANSSON, M., PELL, M., SJODAHL-SVENSSON, K., STENSTROM, J. and TORSTENSSON, L. Microbial biomass and activities in soil as affected by frozen and cold storage. *Soil Biology and Biochemistry*, **30**, 1998, pp. 293-402
- [18] VERCHOT, L.V. Cold storage of a tropical soil decrease nitrification potential. *Soil Science Society of America Journal*, **63**, 1999, pp. 1942-1944

ICS 13.080.05

Price based on 6 pages

This page has been intentionally left blank

BSI - British Standards Institution

BSI is the independent national body responsible for preparing British Standards. It presents the UK view on standards in Europe and at the international level. It is incorporated by Royal Charter.

Revisions

British Standards are updated by amendment or revision. Users of British Standards should make sure that they possess the latest amendments or editions.

It is the constant aim of BSI to improve the quality of our products and services. We would be grateful if anyone finding an inaccuracy or ambiguity while using this British Standard would inform the Secretary of the technical committee responsible, the identity of which can be found on the inside front cover. Tel: +44 (0)20 8996 9000. Fax: +44 (0)20 8996 7400.

BSI offers members an individual updating service called PLUS which ensures that subscribers automatically receive the latest editions of standards.

Buying standards

Orders for all BSI, international and foreign standards publications should be addressed to Customer Services. Tel: +44 (0)20 8996 9001. Fax: +44 (0)20 8996 7001 Email: orders@bsigroup.com You may also buy directly using a debit/credit card from the BSI Shop on the Website <http://www.bsigroup.com/shop>

In response to orders for international standards, it is BSI policy to supply the BSI implementation of those that have been published as British Standards, unless otherwise requested.

Information on standards

BSI provides a wide range of information on national, European and international standards through its Library and its Technical Help to Exporters Service. Various BSI electronic information services are also available which give details on all its products and services. Contact Information Centre. Tel: +44 (0)20 8996 7111 Fax: +44 (0)20 8996 7048 Email: info@bsigroup.com

Subscribing members of BSI are kept up to date with standards developments and receive substantial discounts on the purchase price of standards. For details of these and other benefits contact Membership Administration. Tel: +44 (0)20 8996 7002 Fax: +44 (0)20 8996 7001 Email: membership@bsigroup.com

Information regarding online access to British Standards via British Standards Online can be found at <http://www.bsigroup.com/BSOL>

Further information about BSI is available on the BSI website at <http://www.bsigroup.com>

Copyright

Copyright subsists in all BSI publications. BSI also holds the copyright, in the UK, of the publications of the international standardization bodies. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI.

This does not preclude the free use, in the course of implementing the standard, of necessary details such as symbols, and size, type or grade designations. If these details are to be used for any other purpose than implementation then the prior written permission of BSI must be obtained.

Details and advice can be obtained from the Copyright and Licensing Manager. Tel: +44 (0)20 8996 7070 Email: copyright@bsigroup.com