

**Soil quality —
Determination of the
effects of pollutants on
soil flora — Screening
test for emergence of
lettuce seedlings
(*Lactuca sativa* L.)**

ICS 13.080.30

National foreword

This British Standard reproduces verbatim ISO 17126:2005 and implements it as the UK national standard.

The UK participation in its preparation was entrusted to Technical Committee EH/4, Soil quality, which has the responsibility to:

- aid enquirers to understand the text;
- present to the responsible international/European committee any enquiries on the interpretation, or proposals for change, and keep the UK interests informed;
- monitor related international and European developments and promulgate them in the UK.

A list of organizations represented on this committee can be obtained on request to its secretary.

Cross-references

The British Standards which implement international publications referred to in this document may be found in the *BSI Catalogue* under the section entitled “International Standards Correspondence Index”, or by using the “Search” facility of the *BSI Electronic Catalogue* or of British Standards Online.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with a British Standard does not of itself confer immunity from legal obligations.

Summary of pages

This document comprises a front cover, an inside front cover, the ISO title page, pages ii to v, a blank page, pages 1 to 7 and a back cover.

The BSI copyright notice displayed in this document indicates when the document was last issued.

Amendments issued since publication

Amd. No.	Date	Comments

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 4 March 2005

© BSI 4 March 2005

ISBN 0 580 45559 9

INTERNATIONAL
STANDARD

ISO
17126

First edition
2005-02-01

**Soil quality — Determination of the
effects of pollutants on soil flora —
Screening test for emergence of lettuce
seedlings (*Lactuca sativa* L.)**

*Qualité du sol — Détermination des effets des polluants sur la flore du
sol — Essai de détection de l'émergence des plantules de laitue
(*Lactuca sativa* L.)*



Reference number
ISO 17126:2005(E)

Contents

Page

Foreword	iv
Introduction	v
1 Scope.....	1
2 Normative references	1
3 Terms and definitions.....	1
4 Principle	2
5 Materials.....	2
6 Apparatus	2
7 Procedure	3
7.1 Testing of samples of soil and other test materials	3
7.2 Testing of chemicals.....	3
7.3 Temperature and light regime	4
7.4 Water content	4
7.5 Reference substance	4
7.6 Preparation and start of test dishes.....	4
7.7 Test duration	5
7.8 Measurements	5
7.9 Recordings	5
8 Expression of results.....	5
9 Validity of the test	6
10 Test report	6
Bibliography	7

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 17126 was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

Introduction

For the assessment of the suitability of soil to sustain living organisms, there is a need for simple, rapid, inexpensive biological test methods as a complement to chemical analysis. The method described in this International Standard has been developed for the testing of contaminated soil as well as other contaminated samples. It is cost effective and can be conducted within a short period of time. Furthermore, the test organism is easily available, it does not require advanced equipment for measurements or for growing plants, and it can be conducted by any skilled laboratory technician without special training.

This International Standard is based on US EPA 600/3-88-029 (1989)^[1].

Soil quality — Determination of the effects of pollutants on soil flora — Screening test for emergence of lettuce seedlings (*Lactuca sativa* L.)

1 Scope

This International Standard specifies test procedures for the determination of effects of contaminated soils or other contaminated samples on the emergence of lettuce seeds.

This International Standard is applicable to contaminated soils, soil materials, compost, sludge and chemical testing. It is applicable to the measurement of effects of substances deliberately added to the soil and to the comparison of soils of known and unknown quality.

This International Standard is not applicable to volatile contaminants.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 10381-6, *Soil quality — Sampling — Part 6: Guidance on the collection, handling and storage of soil for the assessment of aerobic microbial processes in the laboratory*

ISO 10390, *Soil quality — Determination of pH*

ISO 11265, *Soil quality — Determination of the specific electrical conductivity*

ISO 11267:1999, *Soil quality — Inhibition of reproduction of *Collembola* (*Folsomia candida*) by soil pollutants*

ISO 11274, *Soil quality — Determination of the water-retention characteristic — Laboratory methods*

ISO 11465, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

seedling emergence

appearance of the seedling (i.e. visible seedling) above the surface of the cover material

3.2

EC_x

concentration of test material (or test substance) estimated to reduce the seedling emergence by *x* % as compared to the control

3.3

test material

material to be tested

EXAMPLES Soils, soil materials, compost, sludge.

3.4

test mixture

mixture of test material (or test substance) and growth medium

4 Principle

Lettuce seeds are exposed to the test material under investigation in a geometric dilution series with test material and growth medium. Incubation takes place under controlled conditions of light and temperature, and lasts usually 5 days (120 h). It is also possible to use this International Standard for chemical testing. In this case, seeds are planted in control pots and in pots containing soil to which the test chemical has been added. At the end of the test, the number of seedlings visible above the sand are counted and recorded. The effect on seedling emergence is expressed as EC50 (possibly EC20), calculated from numbers of emerged seedlings in the control pots (pure growth medium) and in pots containing the test material (or test substance).

5 Materials

5.1 Biological material, in this case lettuce seeds, *Lactuca sativa* L.

Seeds coated with insecticides and/or fungicides ("dressed" seeds) should be avoided.

After purchase, examine the seeds and remove any trash, empty hulls and damaged seeds. A uniform emergence is dependent on uniform seed size. To reduce variability of emergence, the seed batch may be sized before use by means of four sieves with oblong holes (see 6.4) placed on top of each other. In this case, select for testing the fraction with the largest number of seeds.

Pack the seeds in small portions in air-tight containers. The storage time of the seeds should not exceed the expiration date given by the supplier.

Storage at 4 °C is recommended but good emergence may also be accomplished by storage in the dark at 18 °C.

The seeds should not be soaked in water before testing.

5.2 Growth medium, in this case washed, fine quartz sand, e.g. with grain size 0,4 mm to 0,8 mm.

5.3 Cover material, in this case washed, coarse quartz sand, e.g. with grain size 0,7 mm to 1,2 mm (possibly 0,8 mm to 1,4 mm).

The coarse quality of the cover material ensures air exchange between the growth medium and the surroundings.

6 Apparatus

Standard laboratory equipment (pH-meter, thermometer, pipettes, etc.) including the following.

6.1 Balance, with an accuracy of 0,1 g.

6.2 Lower parts of plastic Petri dishes (diameter 15 cm), or other containers with similar surface area, for use as test containers.

- 6.3 Re-sealable polyethylene bags** that fit the test containers (20 cm × 25 cm for a 15 cm Petri dish).
- 6.4 Sieves for seeds**, with oblong mesh dimensions of 0,75 mm × 10 mm, 0,8 mm × 10 mm, 0,85 mm × 10 mm and 0,9 mm × 10 mm.
- 6.5 Sieve** for contaminated soil, stainless steel, with mesh size 2 mm.
- 6.6 Controlled environment chamber:**
- 6.7 Magnifier**

7 Procedure

7.1 Testing of samples of soil and other test materials

Usually the test material is not dried before the test. If necessary, upon reception air-dry it at room temperature to a water content that enables sieving. Immediately thereafter, sieve the test material through a stainless steel sieve (6.5) and store in the dark at $4\text{ °C} \pm 2\text{ °C}$ until testing (in accordance with ISO 10381-6). Preferentially, storage should not exceed three months but if prolonged storage is necessary, it shall be stored at -18 °C . For sieving, a mesh size of 2 mm is preferable but if this is not possible, coarser sieves (e.g. 5 mm) may be used.

Determine the following properties of the sieved test material and register them prior to the test:

- water content (in accordance with ISO 11465);
- water-holding capacity (in accordance with ISO 11274, or alternatively Annex C of ISO 11267:1999);
- conductivity (in accordance with ISO 11265);
- pH (in accordance with ISO 10390).

The water content and water-holding capacity of the test material and the water-holding capacity of the growth medium are used for calculation of the amounts of water to be used for the test. Before the test, weigh and mix the moist test material and the dry growth medium.

Use a geometric dilution series between test material and growth medium (minimum of 5 concentrations), for which the dilution factor should not exceed two. The range of concentrations should include those at which 0 (or minimum) and 100 % emergence are expected, e.g. based on a preliminary test (7.6.1).

The calculations are based on dry mass, and the concentrations are calculated and expressed as grams dry mass test material per gram dry mass test mixture (i.e. test material and growth medium).

NOTE A commonly used dilution factor is $\sqrt[4]{10}$ which is approximately 1,8, resulting in concentrations of e.g. 10, 18, 32, 56, 100.

7.2 Testing of chemicals

7.2.1 General

For testing of chemicals, the test is basically conducted with growth medium only and the chemical is added to the growth medium. The concentration of the chemical is calculated based on the dry mass of the growth medium using the data reported in Clause 8.

7.2.2 Water-soluble substances

Dissolve the chemical in water and add to the test dishes with the moisturizing water (see 7.4).

7.2.3 Substances insoluble in water

Dissolve the test substance in a volume as small as possible of a suitable organic solvent (e.g. acetone or ethanol). Then mix this solution (maximum 1 ml of solvent) with 10 g of the test growth medium per treatment and replicate. Allow the solvent to evaporate, add 90 g of growth medium and mix carefully in order to achieve a homogeneous distribution.

A set of control dishes should be prepared using the same amount of solvent.

7.3 Temperature and light regime

Incubate in a controlled environmental chamber at the optimum temperature that allows germination of the lettuce seeds. This may depend on the strain used (e.g. some strains germinate at 24 °C while other seeds do not tolerate temperatures exceeding 20 °C). The temperature should be kept constant within ± 2 °C of the selected temperature.

During the first 48 h, store the test units in complete darkness. Thereafter, a diurnal cycle (16 h light, 8 h dark) should be maintained with fluorescent light at $4\ 300\ \text{lx} \pm 430\ \text{lx}$ ($30\ \mu\text{E}/\text{m}^2/\text{s} \pm 3\ \mu\text{E}/\text{m}^2/\text{s}$) for the remaining test period.

7.4 Water content

Moisten the test mixture with deionized water to approximately 85 % of the water-holding capacity. Retain the moisture in the experimental units during the test by using polyethylene bags. Measurement of water content during or after the test is thus not normally necessary.

7.5 Reference substance

It is recommended that a reference substance be tested to demonstrate the uniformity of the laboratory test conditions. 2-chloroacetamide or boric acid is suggested as reference substance. A reference test should be carried out regularly and after any major changes in operating procedures are introduced, for example change in phytotron/growth room/greenhouse, change in soil or change in watering regime, etc.

NOTE EC50 values for 2-chloroacetamide and boric acid are found to be 10,4 mg/kg and 406 mg/kg artificial soil, respectively. The value for 2-chloroacetamide is based on artificial soil consisting of silica sand, whereas the value for boric acid is based on artificial soil consisting of 70 % silica sand, 20 % kaolinite clay and 10 % sphagnum peat.

7.6 Preparation and start of test dishes

7.6.1 Preliminary test

In order to establish the range of concentrations of test material or chemical within which the effect is between 0 and 100 %, a preliminary test can be conducted.

For this, the procedures described for the final test (7.6.2) apply, with the exception that Petri dishes with a diameter of 9 cm with 15 seeds in each shall be used and only one replicate is necessary.

7.6.2 Final test

Weigh sufficient test material (moist) and growth medium (dry) to mix an amount equivalent to 300 g to 400 g dry mass of each dilution/concentration. Carefully mix test material and growth medium. Place an amount equivalent to 100 g dry mass of each test mixture in each of three replicate Petri dishes and smooth the surface.

For testing of chemicals, place 100 g dry mass of growth medium in each Petri dish.

As a control, prepare three dishes with growth medium only, following the same procedure as for the test dishes.

Place 40 lettuce seeds on top of the test mixture/growth medium. Distribute the seeds evenly over the area but not closer than 1 cm from the edge of the test container. Press the seeds gently into the medium, e.g. using the bottom of a clean beaker.

Moisten the contents of the dishes with the amount of water calculated to obtain 85 % of the water-holding capacity. Spread the water evenly over the surface.

For chemical testing of water-soluble substances, dissolve the substance in the water before the growth medium is moistened. If soil characteristics cause floating of the seeds upon addition of water, the seeds should be placed in the test container after addition of water.

Cover the contents of each Petri dish evenly with 90 g of dry cover material.

Between operations, cover the dishes with the lid in order to reduce evaporation. Immediately before the dishes are placed in polyethylene bags, remove the lid. Elevate the polyethylene bag to allow room for air in each bag before sealing the bags.

Randomly place the Petri dishes in bags inside the controlled environment chamber for incubation. The specified light regime (7.3) should be ensured during incubation.

7.7 Test duration

Continue incubation until emergence of seedlings has been completed in control dishes, usually 120 h. Depending on the temperature used for the emergence (see 7.3), this period may have to be adjusted. However, the test duration shall not exceed 7 days.

7.8 Measurements

At the beginning of the test period and at the end of the test period (usually 120 h), measure and record the pH and conductivity of samples from moistened sand (control) and from the test mixture least diluted with growth medium. Make both measurements on the overall medium, i.e. test mixture plus cover material.

Temperature shall be recorded daily in the chamber and inside one of the bags, chosen at random, in order to ensure that the temperature is not elevated inside the bags.

7.9 Recordings

At the end of the test, determine the number of emerged seedlings by counting each seedling that has emerged (i.e. is visible) above the surface of the cover material. Additional information may be obtained by careful inspection of the seedlings. Any observed effects should be recorded.

8 Expression of results

The results, i.e. the number of seedlings emerged, are expressed by means of probit analysis or other applicable statistical methods, as the EC50, which corresponds to 50 % of the mean seedling emergence in the controls, or the EC20, which corresponds to 80 % of the mean seedling emergence in the controls. EC50 and/or EC20 shall be expressed as grams dry mass test material per gram dry mass test mixture.

The following data shall be reported:

- a) EC50 and /or EC20, as required;
- b) the 95 % confidence limits of the above value(s);
- c) the mean seedling emergence in the controls;
- d) the method of estimation of the confidence limits.

The 95 % confidence interval can be estimated by a Fieller approximation [4]. If the dose-response relationship proceeds in a non-linear fashion, it is recommended to seek statistical expertise. If no dose-response relationship is obtained, repeat the test.

The data should be presented in tabular form, presenting the result of each test container and indicating the mean number and standard deviation of emerged seeds for each concentration. The statistical method for estimating EC_x values should be specified.

9 Validity of the test

The results are considered to be valid if the following condition is met in the control :

- mean seedling emergence is at least 80 %.

10 Test report

The test report shall provide the following information:

- a) a reference to this International Standard (ISO 17126:2005);
- b) a full description of the experimental design and procedures;
- c) all data required for the identification of the test substance, preparation or test substrate;
- d) pretreatment, if any, of the waste or soil submitted to testing (storage time and storage conditions, drying);
- e) complete description of the biological material, i.e. seed variety, batch number, seed size, etc;
- f) description of the test substrate (water content, water-holding capacity, conductivity and pH of the sieved test material);
- g) description of preparation of test mixtures;
- h) environmental conditions of the test;
- i) a table indicating the emergence obtained in each container, for each test mixture and for controls;
- j) statistically calculated values (EC_x) including 95 % confidence limits, the method used to calculate EC_x, parameter estimates, and a plot of the dose-response relationship;
- k) results obtained if a positive control was performed;
- l) all operating details not specified in this International Standard and conditions that are likely to influence the results;
- m) discussion of the results.

Bibliography

- [1] US EPA 600/3-88-029 (1989): Protocols for short term toxicity screening of hazardous waste sites. A.8.6. Lettuce seed germination (*Lactuca sativa*)
- [2] KEDDY C., J.C. GREENE & M.A. BONNELL (1994): A review of whole organism bioassays for assessing the quality of soil, freshwater sediment and freshwater in Canada. *Environment Canada*, Scientific Series No. 198, Ottawa, Ontario, Canada. 183 pp.
- [3] KEDDY C., J.C. GREENE & M.A. BONNELL (1995): Review of whole-organism bioassays: Soil, freshwater sediment and freshwater assessment in Canada. *Ecotoxicology and Environmental Safety*, **30**, 221-251
- [4] FINNEY, D. J. (1978) *Statistical Methods in Biological Assay*. Charles Griffin and Co. Ltd, London

BSI — British Standards Institution

BSI is the independent national body responsible for preparing British Standards. It presents the UK view on standards in Europe and at the international level. It is incorporated by Royal Charter.

Revisions

British Standards are updated by amendment or revision. Users of British Standards should make sure that they possess the latest amendments or editions.

It is the constant aim of BSI to improve the quality of our products and services. We would be grateful if anyone finding an inaccuracy or ambiguity while using this British Standard would inform the Secretary of the technical committee responsible, the identity of which can be found on the inside front cover.
Tel: +44 (0)20 8996 9000. Fax: +44 (0)20 8996 7400.

BSI offers members an individual updating service called PLUS which ensures that subscribers automatically receive the latest editions of standards.

Buying standards

Orders for all BSI, international and foreign standards publications should be addressed to Customer Services. Tel: +44 (0)20 8996 9001.
Fax: +44 (0)20 8996 7001. Email: orders@bsi-global.com. Standards are also available from the BSI website at <http://www.bsi-global.com>.

In response to orders for international standards, it is BSI policy to supply the BSI implementation of those that have been published as British Standards, unless otherwise requested.

Information on standards

BSI provides a wide range of information on national, European and international standards through its Library and its Technical Help to Exporters Service. Various BSI electronic information services are also available which give details on all its products and services. Contact the Information Centre.
Tel: +44 (0)20 8996 7111. Fax: +44 (0)20 8996 7048. Email: info@bsi-global.com.

Subscribing members of BSI are kept up to date with standards developments and receive substantial discounts on the purchase price of standards. For details of these and other benefits contact Membership Administration.
Tel: +44 (0)20 8996 7002. Fax: +44 (0)20 8996 7001.
Email: membership@bsi-global.com.

Information regarding online access to British Standards via British Standards Online can be found at <http://www.bsi-global.com/bsonline>.

Further information about BSI is available on the BSI website at <http://www.bsi-global.com>.

Copyright

Copyright subsists in all BSI publications. BSI also holds the copyright, in the UK, of the publications of the international standardization bodies. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI.

This does not preclude the free use, in the course of implementing the standard, of necessary details such as symbols, and size, type or grade designations. If these details are to be used for any other purpose than implementation then the prior written permission of BSI must be obtained.

Details and advice can be obtained from the Copyright & Licensing Manager.
Tel: +44 (0)20 8996 7070. Fax: +44 (0)20 8996 7553.
Email: copyright@bsi-global.com.