

Erythema reference action spectrum and standard erythema dose

ICS 11.040.01: 17.080.20

National foreword

This British Standard reproduces verbatim ISO 17166:1999 and implements it as the UK national standard.

The UK participation in its preparation was entrusted to Technical Committee CPL/34, Lamps and related equipment, which has the responsibility to:

- aid enquirers to understand the text;
- present to the responsible international/European committee any enquiries on the interpretation, or proposals for change, and keep the UK interests informed;
- monitor related international and European developments and promulgate them in the UK.

A list of organizations represented on this committee can be obtained on request to its secretary.

Cross-references

The British Standards which implement international publications referred to in this document may be found in the BSI Standards Catalogue under the section entitled “International Standards Correspondence Index”, or by using the “Find” facility of the BSI Standards Electronic Catalogue.

A British Standard does not purport to include all the necessary provisions of a contract. Users of British Standards are responsible for their correct application.

Compliance with a British Standard does not of itself confer immunity from legal obligations.

Summary of pages

This document comprises a front cover, an inside front cover, the ISO title page, pages ii to vi, pages 1 to 4, an inside back cover and a back cover.

The BSI copyright notice displayed in this document indicates when the document was last issued.

This British Standard, having been prepared under the direction of the Electrotechnical Sector Committee, was published under the authority of the Standards Committee and comes into effect on 15 March 2000

© BSI 03-2000

Amendments issued since publication

| Amd. No. | Date | Comments |
|----------|------|----------|
| | | |
| | | |
| | | |
| | | |
| | | |

INTERNATIONAL
STANDARD

ISO
17166
CIE S 007/E

First edition
1999-12-15

**Erythema reference action spectrum and
standard erythema dose**

Spectre d'action érythémale de référence et dose érythémale normalisée



Reference number
ISO 17166:1999(E)
CIE S 007/E-1998

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standard ISO 17166 was prepared as Standard CIE S 007/E by the International Commission on Illumination, which has been recognized by the ISO Council as an international standardizing body. It was adopted by ISO under a special procedure which requires approval by at least 75 % of the member bodies casting a vote, and is published as a joint ISO/CIE edition.

The International Commission on Illumination (abbreviated as CIE from its French title) is an organization devoted to international cooperation and exchange of information among its member countries on all matters relating to the science and art of lighting.

Annex A of this International Standard is for information only.



ISO 17166:1999(E)
CIE S 007/E-1998

Standard

Erythema Reference Action Spectrum and Standard Erythema Dose

Spectre d'action erithémale de référence et dose erithémale normalisée

Erythemale Referenzwirkungsfunktion und standardisierte Erythemdosis

CIE Standards are copyrighted and shall not be reproduced in any form, entirely or partly, without the explicit agreement of the CIE.

CIE Central Bureau,
Kegelgasse 27, A-1030 Vienna, Austria

S 007/E

UDC: 612.014.481
614.898

Descriptor: Optical radiation effects on humans
Protection of the skin

Contents

| | |
|---|---|
| Foreword | 1 |
| 1. Introduction | 1 |
| 2. Scope | 1 |
| 3. Normative references | 1 |
| 4. Definitions | 2 |
| 5. The concept of erythema irradiance and dose | 2 |
| <i>5.1 Erythema effective irradiance</i> | 2 |
| <i>5.2 The erythema action spectrum</i> | 3 |
| <i>5.3 The standard erythema dose (erythema quantity and radiometric equivalence)</i> | 3 |
| Annex A: Bibliography (Non-normative annex) | 4 |

Foreword

Standards produced by the Commission Internationale de l'Éclairage (CIE) are a concise documentation of data defining aspects of light and lighting, for which international harmony requires such unique definition. CIE Standards are therefore a primary source of internationally accepted and agreed data, which can be taken, essentially unaltered, into universal standard systems.

The CIE undertook a major review of its official recommendations on photobiological effects, their dose relationships and measurement. Based on these investigations the present standard describes present day knowledge of the subject.

This Standard has been approved by the National Committees of the CIE and supersedes the recommendations made in Publication *CIE 106/4* - 1993 (reprint from *CIE-Journal 6/1* 17-22 1987) *A reference action spectrum for ultraviolet induced erythema in human skin*.

This CIE Standard reflects present day knowledge of UV radiation effect on humans, specially erythema, but does not absolve those carrying out experiments with humans from their responsibility for the safety and well being of the subjects involved.

1. Introduction

The problem of dosimetry in skin photobiology lies in the fact that the ability of ultraviolet (UV) radiation to elicit erythema in human skin depends strongly on wavelength, encompassing a range of four orders of magnitude between 250 nm and 400 nm. Thus a statement that a subject received an exposure dose of $1 \text{ J}\cdot\text{cm}^{-2}$ ($10^4 \text{ J}\cdot\text{m}^{-2}$) of UV radiation conveys nothing about the consequences of that exposure in terms of erythema. If the radiation source was a UVA fluorescent lamp, no erythema response would be seen apart from in people exhibiting severe, abnormal pathological photosensitivity. The same dose delivered from an unfiltered mercury arc lamp or fluorescent sun-lamp would result in marked violaceous erythema in most white skinned individuals. Consequently, photobiologists have long recognised the need to express the exposure as an erythemally-weighted quantity¹.

Recently the term *minimal erythema dose (MED)* has been used widely as a 'measure' of erythemal radiation. This is unreasonable because the *MED* is not a standard measure of anything but, on the contrary, encompasses the variable nature of individual sensitivity to ultraviolet radiation. Variables which affect the *MED* include optical and radiometric characteristics of the source; determinants of the exposure such as dose increment and field size; nature of the skin such as pigmentation, previous light exposure, and anatomical site; and observational factors such as definition of the end point, time of reading after exposure, and ambient illumination.

To avoid further confusing misuse of the term *MED*, we propose that this term be reserved solely for observational studies in humans and other animals, and that a new term, the *standard erythema dose (SED)* be used as a standardized measure of erythemogenic UV radiation.

2. Scope

This Standard specifies the erythema reference action spectrum ($s_{\text{er}}(\lambda)$), and the Standard Erythema Dose (*SED*).

3. Normative references

The following standards contain provisions which, through reference in this text, constitute provisions of this Standard. At the time of publication, the editions indicated were valid. All standards are subject to revision, and parties to agreements based on this Standard are encouraged to investigate the possibility of applying most recent editions of the standards indicated below. Members of CIE, the International Electrotechnical Commission (IEC) and the International Organization for Standardization (ISO) maintain registers of currently valid international standards.

- CIE 17.4-1987: International Lighting Vocabulary - equivalent to IEC 50(845)
- CIE 90-1991: Sunscreen testing (UVB).
- CIE 98-1992: Personal dosimetry of UV radiation.

ISO 17166:1999(E)

CIE S 007/E-1998

CIE 103/3-1993: Reference action spectra for ultraviolet induced erythema and pigmentation of different human skin (CIE Collection in Photobiology and Photochemistry).

CIE 125-1997: Standard Erythema Dose, a Review.

4. Definitions

erythema action spectrum ($s_{er}(\lambda)$)

Spectral dependence of the ability of UV radiation to produce erythema in human skin.

Note: It is usual to tabulate and plot the $s_{er}(\lambda)$ normalized to its maximum.

erythemal effective irradiance (E_{er})

Quantity defined by the equation:

$$E_{er} = \int E_{\lambda} \cdot s_{er}(\lambda) d\lambda$$

where E_{λ} is the spectral irradiance in $W \cdot m^{-2} \cdot nm^{-1}$ and $s_{er}(\lambda)$ is the erythema action spectrum normalized to its maximum.

erythemal effective radiant exposure (H_{er}), also called the effective dose or erythemal dose:

The time integral of erythemal effective irradiance defined by the equation

$$H_{er} = \iint E_{\lambda} \cdot s_{er}(\lambda) d\lambda dt$$

where E_{λ} is the spectral irradiance in $W \cdot m^{-2} \cdot nm^{-1}$ and $s_{er}(\lambda)$ is the erythema action spectrum.

minimal erythema dose (MED):

A subjective measure based on the reddening of the skin; it depends on many variables, e.g. individual sensitivity to UVR, radiometric characteristics of the source, skin pigmentation, anatomic site, elapsed time between irradiation and observing the reddening (typical value: 24 hours), etc. It should be reserved solely for observational studies in humans and other animals.

standard erythema dose (SED):

Standardised measure of erythemogenic UV radiation, 1 SED is equivalent to an erythemal effective radiant exposure of $100 J \cdot m^{-2}$.

ultraviolet radiation (UVR):

Optical radiation in the 100 nm to 400 nm wavelength range.

5. The concept of erythemal irradiance and dose

5.1 Erythemal effective irradiance

The erythemal effective irradiance (E_{er}) from a source of ultraviolet radiation is obtained by weighting the spectral irradiance of the radiation at wavelength λ in nm by the effectiveness of radiation of this wavelength to cause a minimal erythema and summing over all wavelengths present in the source spectrum. This can be expressed mathematically as:

$$E_{er} = \int E_{\lambda} \cdot s_{er}(\lambda) d\lambda \quad (1)$$

or

$$E_{er} = \sum E_{\lambda} \cdot s_{er}(\lambda) \cdot \Delta\lambda$$

E_{λ} is the spectral irradiance in $W \cdot m^{-2} \cdot nm^{-1}$ at wavelength λ in nm and $\Delta\lambda$ is the wavelength interval used in the summation. $s_{er}(\lambda)$ is a measure of the effectiveness of radiation of wavelength λ in nm relative to some reference wavelength in producing a minimal erythema. Integration has to be carried out in the wavelength range where neither E_{λ} or $s_{er}(\lambda)$ equal zero. As it is a ratio, $s(\lambda)$ is of dimension unity. The effective irradiance is equivalent to a hypothetical irradiance of monochromatic radiation having a

wavelength at which $s_{er}(\lambda)$ is equal to unity. The time integral of effective irradiance is the erythema effective radiant exposure (also called the *effective dose* or *erythema dose*).

The dose (expressed as an *erythema quantity*) received after an exposure period of t seconds is

$$H_{er} = E_{er} \cdot t / \Phi \quad (2)$$

where Φ is the numerical value in $\text{J}\cdot\text{m}^{-2}$ equivalent to one *erythema quantity*.

5.2 The erythema action spectrum

The ability of UV radiation to produce erythema in human skin is highly dependent upon the radiation wavelength, and is expressed by the erythema action spectrum. Erythema action spectra have been the subject of experimental and theoretical interest for over 70 years. The CIE first considered the adoption of a so-called standard erythema curve in 1935². The erythema action spectrum defined in this standard was first introduced by the CIE in 1987³ (See also ⁴⁻⁶). It is represented by relatively simple functions over three clearly defined spectral regions:

$$s_{er}(\lambda) = 1,0 \quad \text{for } 250 \leq \lambda \leq 298 \text{ nm} \quad (3)$$

$$s_{er}(\lambda) = 10^{0,094(298-\lambda)} \quad \text{for } 298 < \lambda \leq 328 \text{ nm} \quad (4)$$

$$s_{er}(\lambda) = 10^{0,015(140-\lambda)} \quad \text{for } 328 < \lambda \leq 400 \text{ nm} \quad (5)$$

5.3 The standard erythema dose (erythema quantity and radiometric equivalence)

There are many biological endpoints in photobiology in which the effectiveness varies with wavelength. For this reason the new quantity shall contain the word *erythema* to make it explicit that we are referring here only to that specific biological response. The term *standard erythema dose*⁷, acronym *SED*, shall be used to express this quantity.

1 SED is equivalent to an erythema effective radiant exposure of $100 \text{ J}\cdot\text{m}^{-2}$.

Note: When deciding upon the numerical value of the radiometric equivalence, the choice is arbitrary - there is no 'right' value. However in order to avoid confusion between *MED* and the *SED* we do not want the *SED* to be interpreted as the *MED* in some particular skin type. Consequently⁸ 1 SED is equivalent to an erythema effective radiant exposure of $100 \text{ J}\cdot\text{m}^{-2}$. The *MEDs* in subjects with skin types I to IV would be expected to lie between erythema effective radiant exposures of $150 \text{ J}\cdot\text{m}^{-2}$ to $600 \text{ J}\cdot\text{m}^{-2}$, equivalent to 1,5 SED to 6 SED.

Annex A: Bibliography (Non-normative annex)

1. Diffey BL. Whatever happened to the erythema unit? *Photodermatology* 1984; **1**: 103-105.
2. International Commission on Illumination. Berlin. *Comptes Rendus de la Commission Internationale de l'Éclairage* 1935; **9**: 596-625.
3. McKinlay AF, Diffey BL, A reference action spectrum for ultraviolet induced erythema in human skin. *CIE J* 1987; **6**: 17-22.
4. Urbach F, Man and ultraviolet radiation. In *Human Exposure to Ultraviolet Radiation: Risks and Regulations* (Passchier WF and Bosnjakovic BFM, eds), Excerpta Medica, Amsterdam, 1987, pp. 3-17.
5. Diffey BL, Observed and predicted minimal erythema doses: a comparative study. *Photochem Photobiol.* 1994; **60**: 380-382.
6. International Commission on Illumination, Standard Erythema Dose, Publication *CIE 125* - 1997.
7. Wulf HC, The need for a standard erythema dose: proposed definition. In: Proceedings of the 12th International Congress on Photobiology, Organizzazione Editoriale Medico Farmaceutica, Milan, in press.
8. Lock-Andersen J, Wulf HC, Mortensen NM, Erythemally weighted radiometric dose and standard erythema dose (SED). In: Proceedings of the 12th International Congress on Photobiology, Organizzazione Editoriale Medico Farmaceutica, Milan, in press.

Copies of CIE Publications are available from the National Committees of most CIE member countries. Purchasers in other countries may obtain them from the CIE Central Bureau, Kegelgasse 27, A-1030 Vienna, Austria.

CIE Central Bureau
Kegelgasse 27
A-1030 Vienna
Austria

Tel.: +43 1 714 3187 0
Fax: +43 1 713 0838 18
e-mail: ciecb@ping.at

WWW: <http://www.cie.co.at/cie/home.html>

BSI — British Standards Institution

BSI is the independent national body responsible for preparing British Standards. It presents the UK view on standards in Europe and at the international level. It is incorporated by Royal Charter.

Revisions

British Standards are updated by amendment or revision. Users of British Standards should make sure that they possess the latest amendments or editions.

It is the constant aim of BSI to improve the quality of our products and services. We would be grateful if anyone finding an inaccuracy or ambiguity while using this British Standard would inform the Secretary of the technical committee responsible, the identity of which can be found on the inside front cover. Tel: 020 8996 9000. Fax: 020 8996 7400.

BSI offers members an individual updating service called PLUS which ensures that subscribers automatically receive the latest editions of standards.

Buying standards

Orders for all BSI, international and foreign standards publications should be addressed to Customer Services. Tel: 020 8996 9001. Fax: 020 8996 7001.

In response to orders for international standards, it is BSI policy to supply the BSI implementation of those that have been published as British Standards, unless otherwise requested.

Information on standards

BSI provides a wide range of information on national, European and international standards through its Library and its Technical Help to Exporters Service. Various BSI electronic information services are also available which give details on all its products and services. Contact the Information Centre. Tel: 020 8996 7111. Fax: 020 8996 7048.

Subscribing members of BSI are kept up to date with standards developments and receive substantial discounts on the purchase price of standards. For details of these and other benefits contact Membership Administration. Tel: 020 8996 7002. Fax: 020 8996 7001.

Copyright

Copyright subsists in all BSI publications. BSI also holds the copyright, in the UK, of the publications of the international standardization bodies. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI.

This does not preclude the free use, in the course of implementing the standard, of necessary details such as symbols, and size, type or grade designations. If these details are to be used for any other purpose than implementation then the prior written permission of BSI must be obtained.

If permission is granted, the terms may include royalty payments or a licensing agreement. Details and advice can be obtained from the Copyright Manager. Tel: 020 8996 7070.