

DD ISO/TS 23647:2010



BSI Standards Publication

Vegetable fats and oils — Determination of wax content by gas chromatography

NO COPYING WITHOUT BSI PERMISSION EXCEPT AS PERMITTED BY COPYRIGHT LAW

raising standards worldwide™



National foreword

This Draft for Development is the UK implementation of ISO/TS 23647:2010.

This publication is not to be regarded as a British Standard.

It is being issued in the Draft for Development series of publications and is of a provisional nature. It should be applied on this provisional basis, so that information and experience of its practical application can be obtained.

Comments arising from the use of this Draft for Development are requested so that UK experience can be reported to the international organization responsible for its conversion to an international standard. A review of this publication will be initiated not later than 3 years after its publication by the international organization so that a decision can be taken on its status. Notification of the start of the review period will be made in an announcement in the appropriate issue of *Update Standards*.

According to the replies received by the end of the review period, the responsible BSI Committee will decide whether to support the conversion into an international Standard, to extend the life of the Technical Specification or to withdraw it. Comments should be sent to the Secretary of the responsible BSI Technical Committee at British Standards House, 389 Chiswick High Road, London W4 4AL.

The UK participation in its preparation was entrusted to Technical Committee AW/307, Oil seeds, animal and vegetable fats and oils and their by products.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

© BSI 2010

ISBN 978 0 580 66083 2

ICS 67.200.10

Compliance with a British Standard cannot confer immunity from legal obligations.

This Draft for Development was published under the authority of the Standards Policy and Strategy Committee on 31 October 2010.

Amendments issued since publication

Date	Text affected
------	---------------

TECHNICAL SPECIFICATION

DD ISO/TS 23647:2010
**ISO/TS
23647**

First edition
2010-10-01

Vegetable fats and oils — Determination of wax content by gas chromatography

Corps gras d'origine végétale — Détermination de la teneur en cires par chromatographie en phase gazeuse



Reference number
ISO/TS 23647:2010(E)

© ISO 2010

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.



COPYRIGHT PROTECTED DOCUMENT

© ISO 2010

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Contents

Page

Foreword	iv
Introduction.....	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle.....	1
5 Reagents and materials	1
6 Apparatus	2
7 Sampling.....	3
8 Preparation of the test sample	3
9 Procedure	3
9.1 Preparation of silver nitrate-impregnated silica gel 60	3
9.2 Column packing.....	3
9.3 Isolation of waxes.....	3
9.4 Gas chromatographic determination of waxes	4
10 Precision.....	6
10.1 Results of interlaboratory test	6
10.2 Repeatability	6
10.3 Reproducibility	6
11 Test report.....	7
Annex A (informative) Chromatograms	8
Annex B (informative) Results of interlaboratory tests	12
Annex C (informative) Preparation of crystalline sunflower wax standard	13
Bibliography.....	16

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

In other circumstances, particularly when there is an urgent market requirement for such documents, a technical committee may decide to publish other types of document:

- an ISO Publicly Available Specification (ISO/PAS) represents an agreement between technical experts in an ISO working group and is accepted for publication if it is approved by more than 50 % of the members of the parent committee casting a vote;
- an ISO Technical Specification (ISO/TS) represents an agreement between the members of a technical committee and is accepted for publication if it is approved by 2/3 of the members of the committee casting a vote.

An ISO/PAS or ISO/TS is reviewed after three years in order to decide whether it will be confirmed for a further three years, revised to become an International Standard, or withdrawn. If the ISO/PAS or ISO/TS is confirmed, it is reviewed again after a further three years, at which time it must either be transformed into an International Standard or be withdrawn.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO/TS 23647 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 11, *Animal and vegetable fats and oils*.

Introduction

Vegetable fats and oils consist predominantly of triacylglycerols, but they also contain small amounts of non-glyceridic substances often called minor components. The composition of these minor constituents (e.g. sterols, steryl esters, triterpene dialcohols, and waxes) provides highly characteristic information about the identity of the oils. Since the physical and chemical steps in the processing of vegetable fats and oils can cause alteration in their composition and content, analysis of these minor components can be conveniently applied to characterize the processing steps through which an oil was obtained.

Waxes are natural compounds occurring in various vegetable oils and can easily crystallize at low temperatures resulting in cloudiness. Wax removal is very often a part of the vegetable oil-refining process (e.g. sunflower, rice bran, corn oil) and its efficiency requires measurement.

This Technical Specification does not cover the analysis of waxes of olive oil.

With the exception of olive oil, there is currently no reliable official method to measure the wax content and composition of vegetable fats and oils. Cold tests give neither qualitative nor quantitative results; methods developed for olive oils are not applicable to seed oils and can cause a lot of problems in the interpretation of results. There is an industrial and commercial need for an International Standard which is applicable to crude, degummed, pre-dewaxed, winterized, and fully refined vegetable fats and oils.

Waxes from different vegetable oils and fats are separated from triacylglycerols and from other non-glyceridic compounds containing double bonds by column chromatography using a mixed column packing consisting of silica gel and silica gel impregnated with AgNO_3 .

The wax fraction is further analysed by capillary gas chromatography. The method also gives information about the total content of waxes and their composition. The information given by the method can be easily used to check oil quality and to track the efficiency of the dewaxing process.

Vegetable fats and oils — Determination of wax content by gas chromatography

1 Scope

This Technical Specification specifies a gas chromatographic method for determining the wax content of crude, degummed, neutralized, winterized, and fully refined vegetable oils, such as sunflower, soybean, rapeseed, corn, and rice bran oils. It is not applicable to olive oils or olive pomace oils.

Waxes are esters of long chain fatty acids and fatty alcohols (having C₂₀ or longer saturated carbon chain).

The wax content is expressed in milligrams per kilogram of oil.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 661, *Animal and vegetable fats and oils — Preparation of test sample*

3 Terms and definitions

For the purpose of this document, the following terms and definitions apply.

3.1

wax content

mass fraction of those substances in the sample, which are determined under the conditions specified in this Technical Specification

NOTE The wax content is expressed in milligrams per kilogram of oil.

4 Principle

The waxes are separated by column chromatography using a mixed column packing consisting of silica gel and silica gel impregnated with AgNO₃. Determination of the waxes is carried out using capillary gas chromatography (GC), applying an internal standard, previously added to the oil.

5 Reagents and materials

WARNING — Attention is drawn to the regulations which specify the handling of dangerous matter. Technical, organizational and personal safety measures shall be followed.

During the analysis, unless otherwise stated, use only reagents of recognized analytical grade suitable for HPLC column chromatography and distilled or demineralized water or water of equivalent purity.

- 5.1 *n*-Hexane.
- 5.2 Dichloromethane.
- 5.3 *n*-Heptane.
- 5.4 Chloroform.
- 5.5 Silica gel 60, of particle size 0,063 mm to 0,200 mm (70 mesh to 230 mesh), such as Merck No. 107734¹).
- 5.6 Silver nitrate (AgNO₃).
- 5.7 Hexatriacontane²) (C₃₆ paraffin).
- 5.7.1 Internal standard solution, mass concentration 0,1 mg/ml in *n*-heptane.
- 5.7.2 Standard for determination of response factor, mass concentration 1 mg/ml in *n*-heptane.
- 5.8 Standards for peak identification in the gas chromatogram, pure waxes³), e.g. C₄₀, C₄₄.
- 5.8.1 Standard for determination of response factor, stearyl stearate or stearic acid stearyl ester (C₃₆ wax), mass concentration 1 mg/ml in chloroform.
- 5.9 Crystallized sunflower wax, prepared from winter cake or from crude sunflower oil using crystallization. For the preparation, see Annex C.
- 5.10 Cotton wool, surgical quality, non-absorbent.

6 Apparatus

- 6.1 Chromatographic column, made of glass, of internal diameter 30 mm and length 450 mm, equipped with a polytetrafluoroethylene (PTFE) tap.
- 6.2 Glass rod, about 600 mm in length.
- 6.3 Pasteur pipette, ISO 7712^[4].
- 6.4 Vacuum rotary evaporator.
- 6.5 Round-bottomed flask, capacity 250 ml (to collect wax fraction).
- 6.6 Pear-shaped flask, capacity 25 ml (to collect the concentrated wax fraction).
- 6.7 Gas chromatograph, equipped with flame ionization detector, split/splitless injector and integrator or data acquisition system.

1) Merck No. 107734 is an example of a suitable product available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of this product. Equivalent products may be used if they can be shown to lead to the same results.

2) Hexatriacontane is available from Sigma-Aldrich with a purity of min. 99 % mass fraction. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of this supplier.

3) Pure waxes e.g. behenic acid stearyl ester (C₄₀), stearic acid stearyl ester (C₃₆) and behenic acid behenyl ester (C₄₄) are available from Sigma-Aldrich, purity approximately 99 % mass fraction. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of this supplier.

6.8 Fused silica capillary column, length 25 m, internal diameter 0,32 mm or preferably 0,20 mm, coated with dimethylpolysiloxane (e.g. HP-1 or OV-1 or equivalent), film thickness 0,11 μm .

7 Sampling

Sampling is not part of the method specified in this Technical Specification. A recommended sampling method is given in ISO 5555^[1].

It is important the laboratory receive a truly representative sample which has not been damaged or changed during transport or storage.

8 Preparation of the test sample

Prepare the test sample in accordance with ISO 661.

Heat the laboratory sample up to 130 °C. Mix it so as to melt completely the crystallized waxes and remove any moisture (a magnetic stirrer or microwave oven may be used).

9 Procedure

9.1 Preparation of silver nitrate-impregnated silica gel 60

Place 100 g of silica gel 60 (5.5) into a ceramic bowl. Pour a silver nitrate solution, containing 5 g AgNO_3 (5.6) dissolved in 240 ml water on to the silica gel and, after thorough mixing, make its surface flat. Put this suspension into a cold electric oven, heat to 170 °C and activate the gel overnight. Allow to cool slowly to 50 °C in the electric oven (in the dark). Keep this column packing in the dark in a well-sealed bottle. If necessary, remove the dark particles from the impregnated silica gel surface.

The gel contains 5 % mass fraction AgNO_3 .

9.2 Column packing

Using the glass rod (6.2) place a wad of the cotton wool (5.10) in the lower part of the column (6.1) and press it down. Pour about 30 ml of *n*-hexane (5.1) into the column and remove air by pressing the cotton wool down with the rod. Allow the solvent layer to rise approximately 2 mm to 3 mm above the cotton wool before putting the silica gel into the column.

Introduce 3 g of impregnated silica gel (9.1) into the glass column (6.1), remove any entrapped air, and flatten the surface. Add 12 g of silica gel 60 (5.5) on the top of the previous packing, remove any air, and protect the packing against light. Wash the column packing with approximately 90 ml to 100 ml of *n*-hexane, removing any air bubbles by slight tapping if necessary. The tap should be open when adding the solvent for washing. Allow approximately 5 mm of solvent layer above the column packing before adding the sample solution.

9.3 Isolation of waxes

9.3.1 Sample preparation

Solvent mixture A: *n*-hexane (5.1) and dichloromethane (5.2), volume fraction of hexane, $\varphi_{\text{C}_6\text{H}_{14}} = 95 \text{ ml}/100 \text{ ml}$ and of dichloromethane, $\varphi_{\text{CH}_2\text{Cl}_2} = 5 \text{ ml}/100 \text{ ml}$.

Weigh the sample according to the expected wax content (sunflower oil: 3 g from crude oil and neutralized oil, and 4 g to 4,5 g from winterized and fully refined oil, other oils: 4 g to 4,5 g). Add 3 ml of internal standard solution (5.7.1) and add 7 ml of solvent mixture A to the sample. Mix thoroughly and transfer approximately 2 ml of this solution by means of a Pasteur pipette (6.3) into the column.

Adjust the flow rate to 1,5 ml/min to 2 ml/min and wash the inner surface of the glass column three times with approximately 3 ml of the solvent mixture A. Do not let the column run dry, allow approximately 5 mm of solvent layer above the column packing before adding the next portion of the solvent.

9.3.2 Column chromatography

Solvent mixture B: *n*-hexane (5.1) and dichloromethane (5.2), volume fraction of hexane, $\varphi_{\text{C}_6\text{H}_{14}} = 80$ ml/100 ml and of dichloromethane, $\varphi_{\text{CH}_2\text{Cl}_2} = 20$ ml/100 ml.

Elute the waxes with 190 ml of Solvent B. Adjust the flow rate to approximately 3 ml/min.

Collect all material eluting from the column in a round-bottomed flask (6.5), and evaporate the solvent by means of a vacuum rotary evaporator (6.4). Dissolve the residue in a small quantity of chloroform (5.4) transfer the solution into a pear-shaped flask (6.6), evaporate the solvent again and re-dissolve the waxes in approximately 1 ml of chloroform. The GC analysis is carried out using this solution.

9.4 Gas chromatographic determination of waxes

9.4.1 Recommended gas chromatographic conditions

Column:	HP-1 (25 m × 0,20 mm, 0,11 μm film)
Oven:	170 °C (0,1 min); 170 °C to 350 °C (6 °C/min); 350 °C isotherm (20 min)
Carrier gas:	H ₂
Flow rate:	1,4 ml/min
Injector:	355 °C, split ratio: 1:30
Detector:	360 °C (FID)

These conditions may be adjusted in accordance with the characteristics of the gas chromatograph apparatus and the column to give good separation for the wax compounds.

9.4.2 Determination of response factor

Mix equal volumes of the standard solutions (5.7.2 and 5.8.1), and inject 1 μl of the mixture into the GC to determine the response factor, F_r

$$F_r = \frac{A_{\text{IS}} \rho}{A \rho_{\text{IS}}}$$

where

A_{IS} is the area of the hexatriacontane (5.7.2) peak;

A is the area of the stearyl stearate peak;

ρ is the mass concentration of stearyl stearate (5.8.1);

ρ_{IS} is the mass concentration of hexatriacontane (5.7.2).

The value of F_r should be between 1,1 and 1,2. Keep this solution in a sealed vial in the dark and repeat the determination as required, normally once a week.

9.4.3 Calculation of wax content

Inject 1,5 µl to 2 µl of solution of waxes (isolated according to 9.3) into the GC. The elution order is the following: solvent, hydrocarbons, internal standard (hexatriacontane), waxes.

If the isolation (9.3) is done properly, the wax fraction (especially the sunflower wax) should show a typical pattern: continuously decreasing peak area after that for C₄₈ (see Figure A.1 and Table C.1). However, overlapping peaks or big peaks (e.g. sterol esters and triglycerides) at the end of the wax region can make data analysis impossible. In this case, repeat the column chromatography (see Figure A.4).

9.4.3.1 Sunflower oil

The main components of sunflower wax are: C₄₄, C₄₆, C₄₈ waxes. The sunflower wax contains mainly waxes with an even number of carbon atoms e.g. C₄₄, C₄₆, C₄₈, C₅₀, C₅₂, C₅₄. Among these major peaks, minor peaks with an odd carbon number can also be found. All the peaks (minor and major peaks) are used in the calculation.

The calculation is based on a series of previous measurements and the literature (References [5][6][7]). The wax content, w_t , expressed as a mass fraction in milligrams per kilogram oil, is given by:

$$w_t = \frac{F_r \left[A_{>C_{44}} + (1+k) A_{C_{44}} \right] m_{IS}}{A_{IS} m}$$

where

F_r is the response factor (9.4.2);

$A_{>C_{44}}$ is the total area of peaks due to all waxes having more than C₄₄;

$A_{C_{44}}$ is the area of the C₄₄ wax peak;

A_{IS} is the area of the hexatriacontane peak;

k is an empirical factor ($k = 1,0$ for refined oils and winterized oils; $k = 0,5$ for non-refined oils, see Reference [5]);

m_{IS} is the mass, in micrograms, of internal standard added to the oil;

m is the mass, in grams, of the oil.

9.4.3.2 Other oils

The main wax components of vegetable oils (soybean, rapeseed, and corn) are: C₄₀, C₄₁, C₄₂, C₄₄, C₄₆ waxes. Their content is generally below 20 mg/kg in a well-refined deodorized oil. The exception is crude rice bran oil, which can have 1 500 mg/kg to 2 000 mg/kg wax content with a similar composition to sunflower.

9.4.3.2.1 Rice bran oil

The wax content, w_t , expressed as a mass fraction in milligrams per kilogram oil, is given by:

$$w_t = \frac{F_r (A_{\geq C_{42}} m_{IS})}{A_{IS} m}$$

where

- F_r is the response factor (9.4.2);
- $A_{\geq C_{42}}$ is the total area of peaks due to all waxes with C_{42} and above;
- A_{IS} is the area of the hexatriacontane peak;
- m_{IS} is the mass, in micrograms, of internal standard added to the oil;
- m is the mass, in grams, of the oil.

9.4.3.2.2 Other oils (soybean, rapeseed, and corn)

The wax content, w_t , expressed as a mass fraction in milligrams per kilogram oil, is given by

$$w_t = \frac{F_r \left[A_{>C_{44}} + (1+k) A_{C_{44}} \right] m_{IS}}{A_{IS} m}$$

where

- F_r is the response factor (9.4.2);
- $A_{>C_{44}}$ is the total area of peaks due to all waxes having more than C_{44} ;
- $A_{C_{44}}$ is the area of the C_{44} wax peak;
- A_{IS} is the area of hexatriacontane peak;
- k is an empirical factor, $k = 0,5$ (see Reference [5]);
- m_{IS} is the mass, in micrograms, of internal standard added to the oil;
- m is the mass, in grams, of the oil.

NOTE For soybean oil, the sum of all wax peaks can result in a very high wax content and cause inconvenience.

10 Precision

10.1 Results of interlaboratory test

Details of an inter-laboratory test on the precision of the method are summarized in Annex B. The values derived from this interlaboratory test may not be applicable to ranges and matrices other than those given.

10.2 Repeatability

The absolute difference between two independent single test results, obtained using the same method on identical test material in the same laboratory by the same operator using the same equipment within a short interval of time, will in not more than 5 % of cases exceed the values in Table B.1.

10.3 Reproducibility

The absolute difference between two single test results, obtained using the same method on identical test material in two different laboratories with different operators using different equipment, will in not more than 5 % of cases exceed the values in Table B.1.


11 Test report

The test report shall contain at least the following information:


- a) all information necessary for the complete identification of the sample;
- b) the sampling method used, if known;
- c) the test method, with reference to this Technical Specification (ISO/TS 23647:2010);
- d) all operating details not specified in this Technical Specification, or regarded as optional, together with details of any incidents which may have influenced the results;
- e) the test results obtained;
- f) if the repeatability has been checked, the final quoted result obtained.

Annex A (informative)

Chromatograms



a) Chromatogram A — Full gas chromatographic profile




b) Chromatogram B — Enlargement

Key

- I response
- t time
- 1 hexatriacontane internal standard solution, 0,1 mg/ml in *n*-heptane

Figure A.1 — Example of a typical chromatograms of crude sunflower oil wax composition



Key

I response


t time

1 hexatriacontane internal standard solution, 1 mg/ml in *n*-heptane

2 stearyl stearate 1,032 mg/ml in chloroform

NOTE In this case, the response factor, $F_r = 1,09$.

Figure A.2 — Typical chromatogram showing response factor determination




Key

I response

t time


1 hexatriacontane internal standard solution, 1 mg/ml in *n*-heptane

Figure A.3 — Typical chromatogram of winterized, fully refined sunflower oil wax composition




- Key**
- I* response
 - t* time
 - 1 hexatriacontane internal standard solution, 1 mg/ml in *n*-heptane
 - 2 triglycerides

Figure A.4 — Typical chromatogram of a crude sunflower wax co-eluted with triglycerides




- Key**
- I* response
 - t* time
 - 1 hexatriacontane internal standard solution, 1 mg/ml in *n*-heptane

Figure A.5 — Typical chromatogram of soybean oil wax composition



Key
y response
t time
 1 hexatriacontane internal standard solution, 1 mg/ml in *n*-heptane

Figure A.6 — Typical chromatogram of rice bran oil wax composition



Key
I response
t time
 1 hexatriacontane internal standard solution, 1 mg/ml in *n*-heptane
 2 triglycerides

Figure A.7 — Typical chromatogram of corn oil wax composition

Annex B (informative)

Results of interlaboratory tests

An international collaborative test involving seven laboratories in four countries was carried out on eight samples (four duplicate samples).

The results obtained were subjected to statistical analysis in accordance with ISO 5725-1^[2] and ISO 5725-2^[3] to give the precision data shown in Table B.1.

Table B.1 — Statistical evaluation of the collaborative trial

Parameter	Samples of different sunflower oils							
	Crude oil, not winterized		Refined and 10 % crude oil, winterized		Bleached and crude oil, not winterized		Refined oil winterized	
Number of laboratories participating	7	7	7	7	7	7	7	7
Number of laboratories after eliminating outliers	7	7	7	7	7	7	7	7
Number of test results from remaining laboratories	2	2	2	2	2	2	2	2
Mean value, \bar{w}_t , mg/kg	406,9	379,6	107,8	100,4	241,0	241,6	39,6	37,0
Repeatability standard deviation, s_r , mg/kg	21,4	9,9	17,2	6,8	9,09	9,1	1,9	3,2
Coefficient of variation of repeatability, $C_{V,r}$, %	5	3	16	7	4	4	5	9
Repeatability limit, r , (2,8 s_r) mg/kg	59,8	27,6	48,3	19,1	24,4	25,6	5,4	8,9
Reproducibility standard deviation, s_R , mg/kg	66,6	61,0	22,6	26,7	49,9	30,6	12,1	10,9
Coefficient of variation of reproducibility, $C_{V,R}$, %	16	16	21	27	21	13	31	29
Reproducibility limit, R , (2,8 s_R) mg/kg	186,5	170,8	63,2	74,7	139,6	85,8	33,8	30,4

Annex C (informative)

Preparation of crystalline sunflower wax standard

C.1 Reagents and materials

During the analysis, unless otherwise stated, use only reagents of recognized analytical grade suitable for HPLC column chromatography.

C.1.1 *n*-Hexane.

C.1.2 Dichloromethane.

C.1.3 Acetone.

C.1.4 Filter paper, quantitative, 80 g/m².

C.1.5 Sunflower winter cake, filter aid together with oil and wax, oil mass fraction is 50 % to 70 %.

C.1.6 Dried, degummed sunflower oil, with at least 500 mg/kg of wax.

C.1.7 Filter aid, Clarcel⁴⁾, Perfil⁴⁾ or any other similar type.

C.1.8 Bleaching clay, acid activated.

C.2 Apparatus

C.2.1 Magnetic stirrer.

C.2.2 Glass funnel, diameter 120 mm.

C.2.3 Round-bottomed flask, capacity 500 ml.

C.2.4 Round-bottomed flask, capacity 2 000 ml.

C.2.5 Beaker, capacity 1 000 ml.

C.2.6 Erlenmeyer flask, capacity 500 ml.

C.2.7 Vacuum rotary evaporator.

C.2.8 Büchner funnel, internal diameter 80 mm.

4) Product available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of this product.

C.3 Procedure

C.3.1 Method to prepare sunflower wax from winter cake

C.3.1.1 Sample preparation

Weigh approximately 100 g of winter cake (filtration residue of sunflower oil winterization) (C.1.5), add 350 ml of *n*-hexane (C.1.1), and stir it on a magnetic stirrer (C.2.1) for 1 h. After separation of the phases, decant the upper hexane layer which contains the oil. Let the residue stand in a ventilated fume cupboard to allow the hexane to evaporate.

C.3.1.2 Extraction of the wax

Add 100 ml of dichloromethane (C.1.2) to the residue of the winter cake (C.3.1.1) and stir it for 10 min. After separation of the layers, filter it through the filter paper (C.1.4) into a round-bottomed flask (C.2.3) and wash the residue with an additional 300 ml of dichloromethane. Evaporate the solvent on the vacuum rotary evaporator (C.2.7). The residue contains the wax.

C.3.1.3 Purification of the wax

Add 60 ml of acetone (C.1.3) to the isolated wax fraction (C.3.1.2), pour it into a beaker (C.2.5), stir it for 5 min on a magnetic stirrer (C.2.1) and let the phases separate. Filter the upper layer through the filter paper (C.1.4) and add an additional 60 ml of acetone to the residue and repeat the whole procedure at least four or five times. After the last washing step, pour the entire residue on to the filter paper, wash it with approximately 50 ml of acetone and remove the solvent residue in a ventilated fume cupboard.

To remove any traces of filter aid particles, dissolve the isolated wax in dichloromethane, filter it through the filter paper (C.1.4), wash the filter paper with dichloromethane, evaporate the solvent on a vacuum rotary evaporator (C.2.7) and dry the wax in a drying oven at 105 °C.

The purity of wax can be checked by gas chromatography.

NOTE The typical composition of sunflower wax standard prepared according to this method (see Table C.1) depends on the process conditions the winter cake originated from.

C.3.2 Method to prepare sunflower wax from degummed sunflower oil

C.3.2.1 Sample preparation

Weigh approximately 1 000 g of dried degummed sunflower oil (C.1.6) in a round-bottomed flask (C.2.4) equipped with a magnetic stirrer (C.2.1), heat under reduced pressure (10 kPa) to 70 °C, disconnect the vacuum after reaching the desired temperature, add approximately 2 % of bleaching clay (C.1.8), reconnect the vacuum, heat to 100 °C under reduced pressure using mechanical stirring and maintain it under these conditions for 30 min. Cool under reduced pressure, disconnect the vacuum, and filter the contents of the flask through a Büchner funnel (C.2.8), using filter paper (C.1.4) with a 20 mm thick layer of filter aid (C.1.7).

C.3.2.2 Extraction of the wax

Pour the bleached oil into a screw-capped glass container, close, and store in a refrigerator at 0 °C ± 2 °C for at least 2 days. The crystallized wax forms a sediment in the container. Decant the upper layer (pure oil), warm the residue to 130 °C, homogenize and weigh a 20 g aliquot into a glass flask. Add 50 ml of acetone (C.1.3), stir it for 5 min on a magnetic stirrer, allow the phases to separate, decant the upper phase, and pass it through filter paper (C.1.4).

Add another 50 ml of acetone to the lower phase and repeat the procedure in the previous paragraph five or six times.

After the last washing step, place the entire wax residue on a filter paper and wash it with approximately 30 ml of acetone, remove the solvent residue in a ventilated fume cupboard. Check the purity and composition by gas chromatography.

NOTE An example of a typical sunflower wax composition prepared from degummed sunflower oil is given in Table C.2.

Table C.1 — Two examples for a typical sunflower wax composition prepared from winter cake

Wax compound	Without pre-dewaxing step % mass fraction	With pre-dewaxing step % mass fraction
C ₄₂	5,3	5,9
C ₄₃	0,7	0,9
C ₄₄	19,7	23,3
C ₄₅	1,9	2,0
C ₄₆	22,0	23,3
C ₄₇	1,8	1,5
C ₄₈	15,1	14,9
C ₄₉	1,2	1,0
C ₅₀	9,7	8,9
C ₅₁	1,0	0,8
C ₅₂	8,8	7,2
C ₅₃	0,9	0,7
C ₅₄	5,3	4,3
C ₅₅	0,6	0,4
C ₅₆	3,1	2,4
C ₅₇	—	0,3
C ₅₈	1,8	1,4
C ₅₉	—	—
C ₆₀	1,1	0,8

Table C.2 — Example of a typical sunflower wax composition prepared from degummed sunflower oil

Wax compound	Content % mass fraction
C ₄₂	2,3
C ₄₃	0,5
C ₄₄	15,1
C ₄₅	1,6
C ₄₆	21,1
C ₄₇	1,5
C ₄₈	16,6
C ₄₉	1,2
C ₅₀	11,2
C ₅₁	1,1
C ₅₂	10,1
C ₅₃	1,1
C ₅₄	6,6
C ₅₅	0,7
C ₅₆	4,0
C ₅₇	0,5
C ₅₈	2,5
C ₅₉	0,3
C ₆₀	1,6

Bibliography

- [1] ISO 5555, *Animal and vegetable fats and oils — Sampling*
- [2] ISO 5725-1:1994, *Accuracy (trueness and precision) of measurement methods and results — Part 1: General principles and definitions*
- [3] ISO 5725-2:1994, *Accuracy (trueness and precision) of measurement methods and results — Part 2: Basic method for the determination of repeatability and reproducibility of standard measurement method*
- [4] ISO 7712, *Laboratory glassware — Disposable Pasteur pipettes*
- [5] HENON, G. Chromatographic determination of waxes in sunflower oil. *Rev. Fr. Corps Gras* 1986, **33**, pp. 475-482
- [6] RECSEG, K. KEMÉNY, Z. KÖVÁRI, K., DENISE, J. Investigation of non-glyceridic esters of vegetable oils. In: KOSOGLU, S.S., RHEE, K.C., WILSON, R.F., editors. *Proceedings of the World Conference on Oilseed and Edible Oils Processing*. Champaign, IL: AOCS Press, 1998, Vol. II, pp. 342-345
- [7] HENON, G RECSEG, K. KÖVÁRI, K. Wax analysis of vegetable oils using liquid chromatography on a double-adsorbent layer of silica gel and silver nitrate-impregnated silica gel. *J. Am. Oil Chem. Soc.* 2001, **78**, pp. 401-410

ICS 67.200.10

Price based on 16 pages

British Standards Institution (BSI)

BSI is the independent national body responsible for preparing British Standards and other standards-related publications, information and services.

It presents the UK view on standards in Europe and at the international level.

It is incorporated by Royal Charter.

Revisions

British Standards are updated by amendment or revision. Users of British Standards should make sure that they possess the latest amendments or editions.

It is the constant aim of BSI to improve the quality of our products and services. We would be grateful if anyone finding an inaccuracy or ambiguity while using this British Standard would inform the Secretary of the technical committee responsible, the identity of which can be found on the inside front cover.

Tel: +44 (0)20 8996 9001 Fax: +44 (0)20 8996 7001

BSI offers Members an individual updating service called PLUS which ensures that subscribers automatically receive the latest editions of standards.

Tel: +44 (0)20 8996 7669 Fax: +44 (0)20 8996 7001

Email: plus@bsigroup.com

Buying standards

You may buy PDF and hard copy versions of standards directly using a credit card from the BSI Shop on the website www.bsigroup.com/shop. In addition all orders for BSI, international and foreign standards publications can be addressed to BSI Customer Services.

Tel: +44 (0)20 8996 9001 Fax: +44 (0)20 8996 7001

Email: orders@bsigroup.com

In response to orders for international standards, it is BSI policy to supply the BSI implementation of those that have been published as British Standards, unless otherwise requested.

Information on standards

BSI provides a wide range of information on national, European and international standards through its Knowledge Centre.

Tel: +44 (0)20 8996 7004 Fax: +44 (0)20 8996 7005

Email: knowledgecentre@bsigroup.com

Various BSI electronic information services are also available which give details on all its products and services.

Tel: +44 (0)20 8996 7111 Fax: +44 (0)20 8996 7048

Email: info@bsigroup.com

BSI Subscribing Members are kept up to date with standards developments and receive substantial discounts on the purchase price of standards. For details of these and other benefits contact Membership Administration.

Tel: +44 (0)20 8996 7002 Fax: +44 (0)20 8996 7001

Email: membership@bsigroup.com

Information regarding online access to British Standards via British Standards Online can be found at www.bsigroup.com/BSOL

Further information about BSI is available on the BSI website at www.bsigroup.com/standards

Copyright

Copyright subsists in all BSI publications. BSI also holds the copyright, in the UK, of the publications of the international standardization bodies. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI. This does not preclude the free use, in the course of implementing the standard of necessary details such as symbols, and size, type or grade designations. If these details are to be used for any other purpose than implementation then the prior written permission of BSI must be obtained. Details and advice can be obtained from the Copyright & Licensing Manager.

Tel: +44 (0)20 8996 7070

Email: copyright@bsigroup.com

BSI Group Headquarters

389 Chiswick High Road London W4 4AL UK

Tel +44 (0)20 8996 9001

Fax +44 (0)20 8996 7001

www.bsigroup.com/standards

raising standards worldwide™

