

Food analysis — Determination of pesticide residues by LC-MS/MS — Tandem mass spectrometric parameters

ICS 67.050

National foreword

This Published Document is the UK implementation of CEN/TR 15641:2007. The UK participation in its preparation was entrusted to Technical Committee AW/-3, Food analysis — Horizontal methods.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

This Published Document was published under the authority of the Standards Policy and Strategy Committee on 28 September 2007

© BSI 2007

ISBN 978 0 580 58353 7

Amendments issued since publication

Amd. No.	Date	Comments

ICS 67.050

English Version

Food analysis - Determination of pesticide residues by LC-MS/MS - Tandem mass spectrometric parameters

Analyse des produits alimentaires - Détermination des résidus de pesticides par CL-SM/SM - Paramètres de spectrométrie de masse en tandem

Lebensmitteluntersuchung - Bestimmung von Pestizidrückständen mit LC-MS/MS - Parameter für die Tandem-Massenspektrometrie

This Technical Report was approved by CEN on 25 July 2007. It has been drawn up by the Technical Committee CEN/TC 275.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: rue de Stassart, 36 B-1050 Brussels

Contents

Page

Foreword.....	3
Introduction	4
1 Scope	5
2 Analyte specific parameters for selective reaction monitoring of pesticides.....	5
2.1 General.....	5
2.2 LC Parameters.....	6
2.3 General MS/MS parameters	6
2.4 Analyte specific MS/MS Parameters	7
Bibliography.....	23

Foreword

This document (CEN/TR 15641:2007) has been prepared by Technical Committee CEN/TC 275 “Food analysis - Horizontal methods”, the secretariat of which is held by DIN.

Introduction

Pesticide residue analysis employs multi methods involving extraction of residues from foods and clean up of the extract to obtain as many pesticide residues as possible in the purified extracts. Afterwards the extracts can be analysed by different kinds of instruments.

The hyphenation of liquid chromatography (LC) and tandem mass spectrometry (MS/MS) has become one of the most universal, but selective and sensitive analysis techniques for identification and quantification of pesticide residues in extracts of foods.

For the ionization of the analytes (M) in LC-MS/MS, electro spray ionization (ESI) or atmospheric pressure chemical ionization (APCI) interfaces are most commonly used. Depending on the selected polarity of the ionization, protonated or deprotonated molecular ions like $[M+H]^+$ or $[M-H]^-$ are generated. Using ESI, relatively stable adducts (clusters) of the analytes (M) and components of the mobile phase like ions (e.g. ammonium, sodium or potassium ions) or solvent molecules (e.g. methanol) can be additionally formed. To obtain a high yield of quasi molecular ions and therefore to increase the sensitivity of the measurement these clusters have to be broken. When using ESI the formed adducts are accelerated by a potential (e.g. declustering potential or cone voltage) in the ion source at atmospheric pressure. Due to collision with neutral gas molecules the clusters (adducts) are broken in the ion source. Under certain conditions some adduct-ions formed are stable (e.g. with ammonium cation $[M+NH_4]^+$). It should be mentioned, however, that ammonium adducts are usually not generated in the APCI-mode and that their formation when using ESI strongly depends on the concentration of ammonium ions in the LC mobile phase. Adduct ions like $[M+NH_4]^+$ can also be used alternatively for quantification if they were shown to provide reproducible signals. Sodium adducts are usually not suitable for quantitative analysis as their formation and decomposition tends to be highly irreproducible.

The selective determination of each target compound is performed by acquisition of characteristic product ions of the precursor ion (quasi molecular ion or adduct) in the 'selected reaction monitoring' (SRM). Decomposition of the precursor ions in the collision cell is induced by collision with molecules of the collision gas (nitrogen or argon). The collision energy (CE) necessary for fragmentation is a very substantial parameter of the MS/MS optimization for maximum sensitivity.

If LC-MS/MS measurement should be used not only for quantification but also for confirmation of positive findings, at least two SRM transitions have to be recorded.

1 Scope

This Technical Report lists mass spectrometric parameters which are useful for the application of European Standards for the determination of pesticide residues in foods of plant origin that use LC-MS/MS, such as the standards in preparation:

prEN 15637 "Foods of plant origin — Determination of pesticide residues using LC-MS/MS following methanol extraction and clean up using diatomaceous earth"

prEN 15662 "Foods of plant origin — Determination of pesticide residues using GC-MS and/or LC-MS/MS following acetonitrile extraction/partitioning by dispersive SPE — QuEChERS-method"

To facilitate the determination of active substances and/or metabolites using LC-MS/MS, Table 1 specifies the precursor ions and product ions suitable for quantification, which can be used independently of the type of triple quadrupole mass spectrometer. However, using an ion trap mass spectrometer other product ions can be generated or at least the relative intensities of the ions are different to triple quadrupole instruments. Furthermore, the additional parameters declustering potential (DP), collision energy (CE), relative retention times and an approximate classification of detection sensitivity are presented in Table 1. These were derived using the API 2000¹⁾ and should be applicable at least for other instruments of the API type (Applied Biosystems).

2 Analyte specific parameters for selective reaction monitoring of pesticides

2.1 General

All values indicated in Table 2 were acquired using the above mentioned LC-MS/MS system under the experimental conditions as outlined in 2.2. Comparative investigations showed that these parameters can be transferred simply on instruments of other types of the same manufacturer or after adjustment also on devices of other manufacturers (see in 2.3)

¹⁾ Instruments of the API type are products supplied by Applied Biosystems (Foster City, CA, USA). This information is given for the convenience of users of this Technical Report and does not constitute an endorsement by CEN of the product named. Equivalent products may be used if they can be shown to lead to the same results.

2.2 LC Parameters

The following LC operating conditions have been proven to be satisfactory. This is an example for appropriate experimental conditions. Equivalent conditions may be used if they can be shown to lead to the same results.

HPLC pump	HP1100 ^{®2)} Binary Pump (G1312A)	
Autosampler	HP1100 [®] (G1313A)	
Injection volume	20 µl	
Column	Phenomenex ^{® 3)} Aqua 5 µ C18 125 Å, 50 mm × 2 mm	
Mobile phase A	Methanol/water 2+8 (V/V) with 5 mmol/l ammonium formate	
Mobile phase B	Methanol/water 9+1 (V/V) with 5 mmol/l ammonium formate	
Flow rate	0,2 ml/min	
Column temperature	20 °C	
Gradient	Linear:	
	0 min	0 % B
	11 min	100 % B
	23 min	100 % B
	25 min	0 % B
	36 min	0 % B (equilibration time)

As slight fluctuations in the measurement conditions influence the retention time, usually relative retention times (RRT), related to a standard substance, are compared. The standard substance for the calculation of the RRT values in Table 1 was Imazalil (RRT = 1,000).

It could be shown that the use of other mobile phase systems does not shift the order of elution substantially, except for those compounds which are sensitive to the pH of eluent. Often retention data can be transferred to HPLC columns of other manufacturers, if a typical reversed phase is used.

2.3 General MS/MS parameters

The following general MS/MS operating conditions have been proven to be satisfactory. This is an example for appropriate experimental conditions. Equivalent conditions may be used if they can be shown to lead to the same results

MS/MS instrument	Applied Biosystems API 2000 [®]
Ion source	Turbo Ion Spray [®] (ESI)

²⁾ Instruments of the HP type are products supplied by Agilent Technologies Inc. (Palo Alto, CA, USA). This information is given for the convenience of users of this Technical Report and does not constitute an endorsement by CEN of the product named. Equivalent products may be used if they can be shown to lead to the same results.

³⁾ HPLC columns of the Aqua type are products supplied by Phenomenex (Torrance, CA, USA). This information is given for the convenience of users of this Technical Report and does not constitute an endorsement by CEN of the product named. Equivalent products may be used if they can be shown to lead to the same results.

Table 1 — Ion source and general parameters

Ion polarity	positive	Gas 2 temperature	400 °C
Curtain gas	nitrogen, 35 psi (241 kPa)	Resolution MS 1	unit
Collision gas	nitrogen, 2 arbitrary units	Resolution MS 2	unit
Ion spray voltage	5500 V	Dwell time	25 ms
Gas 1	nitrogen, 60 psi (414 kPa)	Focusing potential	360 V
Gas 2	nitrogen, 60 psi (414 kPa)		

2.4 Analyte specific MS/MS Parameters

The analyte specific parameters for selective reaction monitoring of pesticides are listed in Table 1 [1]. The names of the individual analytes are supplemented by the CAS number (Chemical Abstracts Service), which is useful for the search in databases. It is usually taken from [2], but there can be several numbers in individual cases, e.g. for isomers and racemates.

The values for the declustering potential (DP), indicated in Table 1 for the API 2000[®], have to be increased by 20 V for tandem mass spectrometers of the type API 3000[®] or API 4000[®]. It is to be considered that DP breaks not only the clusters but can already induce fragmentation of the precursor ions (at too high values) in the ESI source before entering into the first MS.

NOTE It is not necessary to change the collision energy for API 3000[®] or API 4000[®] instruments as the differences for the CE are less than 5 V.

When using tandem mass spectrometers of other manufacturers the correct value for the collision energy should be determined in tuning experiments for analytes with particularly low and high collision energy since it has relevant influence on the intensity of the SRM transition. Based on these data, in a first approximation the values for the collision energy of further pesticides can be derived proportionally from the data in the table and the observed difference (difference of CE at instrument X to the API 2000[®]). The values of the declustering potential (other name "cone voltage") for other instruments have to be determined individually. On the intensity of the SRM transitions this parameter has a smaller influence than the collision energy.

Table 2 — MS/MS Parameters of 497 analytes

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
2,4-D	94-75-7	ESI - [M-H]-		219,0	-21	160,9	-14	124,9	-34	0,69	***
2,4-DB	94-82-6	ESI - [M-H]-		247,0	-66	160,8	-12	124,9	-34	0,86	***
2-Naphthoxyacetic acid	120-23-0	ESI + [M+NH ₄] ⁺		220,1	36	157,1	19	127,1	43	0,66	n.a.
2-Naphthoxyacetic acid	120-23-0	ESI - [M-H]-		201,1	-71	143,0	-18	114,9	-50	0,66	***
3,4,5-Trimethacarb	2686-99-9	ESI + [M+H] ⁺		194,1	61	137,1	15	122,0	35	0,86	****
4-CPA	122-88-3	ESI - [M-H]-		185,0	-71	126,8	-18	140,7	-12	0,47	**
Acephate	30560-19-1	ESI + [M+H] ⁺		184,1	6	124,9	25	142,9	13	0,11	****
Acetamiprid	135410-20-7	ESI + [M+H] ⁺		223,0	36	126,0	27	90,1	45	0,58	****

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Acibenzolar-S-methyl	135158-54-2	ESI + [M+H] ⁺		210,9	26	136,1	39	140,0	31	0,92	**
Aclonifen	74070-46-5	ESI + [M+H] ⁺		265,0	56	182,1	39	218,0	33	0,99	**
Acrinathrin	101007-06-1	ESI + [M+NH ₄] ⁺		559,1	26	208,1	23	181,1	43	1,20	*
Alachlor	15972-60-8	ESI + [M+H] ⁺		270,1	31	238,1	15	162,2	25	0,97	****
Aldicarb	116-06-3	ESI + [M+NH ₄] ⁺		208,1	1	89,1	21	116,0	13	0,66	****
Aldicarb-sulfoxide	1646-87-3	ESI + [M+H] ⁺		207,1	36	89,1	17	131,9	11	0,15	****
Aldoxycarb	1646-88-4	ESI + [M+NH ₄] ⁺		240,1	11	148,0	19	86,1	27	0,19	****
Alloxydim	55634-91-8	ESI + [M+H] ⁺		324,2	11	178,3	27	234,2	19	0,77	****
Ametryn	834-12-8	ESI + [M+H] ⁺		228,1	36	186,2	25	96,1	35	0,90	****
Amidosulfuron	120923-37-7	ESI + [M+H] ⁺		370,0	21	217,9	31	260,9	19	0,46	****
Aminocarb	2032-59-9	ESI + [M+H] ⁺		209,1	16	152,1	19	137,2	31	0,74	****
Amitraz	33089-61-1	ESI + [M+H] ⁺		294,2	16	163,1	21	122,1	41	1,19	****
Amitrole	61-82-5	ESI + [M+H] ⁺		85,0	51	58,2	29	57,0	23	0,07	**
Aramit	140-57-8	ESI + [M+NH ₄] ⁺		352,1	41	191,2	19	105,0	57	1,09	****
Atrazine	1912-24-9	ESI + [M+H] ⁺		216,1	21	174,0	25	103,9	27	0,83	****
Atrazine, 2-hydroxy-	2163-68-0	ESI + [M+H] ⁺		198,1	66	69,0	47	156,2	25	0,65	****
Atrazine, desethyl-	6190-65-4	ESI + [M+H] ⁺		188,1	56	104,0	33	146,0	25	0,59	***
Atrazine, desethyl-2-hydroxy-	6190-65-4	ESI + [M+H] ⁺		170,1	66	128,1	23	86,0	31	0,14	****
Atrazine, desisopropyl-	1007-28-9	ESI + [M+H] ⁺		174,1	56	104,2	31	96,0	27	0,39	***
Avermectin B1a	65195-55-3	ESI + [M+NH ₄] ⁺		890,5	41	305,1	35	145,2	43	1,33	***
Avermectin B1b	65195-56-4	ESI + [M+NH ₄] ⁺		876,5	41	291,1	35	145,2	43	1,26	***
Azaconazole	60207-31-0	ESI + [M+H] ⁺		300,0	56	231,0	23	159,0	37	0,86	****
Azamethiphos	35575-96-3	ESI + [M+H] ⁺		325,0	16	183,0	21	139,2	33	0,74	****
Azimsulfuron	120162-55-2	ESI + [M+H] ⁺		425,1	31	182,1	23	156,1	43	0,55	***
Azinphos-ethyl	2642-71-9	ESI + [M+H] ⁺		346,0	26	132,2	21	160,2	15	0,96	****
Azinphos-methyl	86-50-0	ESI + [M+H] ⁺		318,0	16	132,2	21	160,2	13	0,89	****
Azocyclotin	41083-11-8	ESI + [M-OH] ⁺		369,2	76	204,8	23	287,0	17	n.a.	n.a.
Azoxystrobin	131860-33-8	ESI + [M+H] ⁺		404,1	36	371,9	19	343,9	29	0,90	****
Beflubutamid	113614-08-7	ESI + [M+NH ₄] ⁺		373,1	26	91,2	47	162,1	39	1,00	***
Benalaxyl	71626-11-4	ESI + [M+H] ⁺		326,2	26	148,2	27	208,2	21	1,01	****
Bendiocarb	22781-23-3	ESI + [M+H] ⁺		224,1	6	167,2	13	108,9	21	0,76	****
Benfuracarb	82560-54-1	ESI + [M+H] ⁺		411,2	1	195,1	31	252,0	19	1,05	****
Benomyl	17804-35-2	ESI + [M+H] ⁺		291,1	16	160,1	35	192,2	17	1,03	**
Bensulfuron-methyl	83055-99-6	ESI + [M+H] ⁺		411,1	51	148,9	27	119,0	51	0,85	***

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Bentazone	25057-89-0	ESI - [M-H]-		239,1	-51	132,0	-32	197,0	-24	0,33	***
Benzoximate	29104-30-1	ESI + [M+H]+		364,1	1	199,1	17	105,1	35	1,02	****
Bifenox	42576-02-3	ESI + [M+NH4]+		358,9	6	309,9	17	189,1	35	1,04	**
Bifenthrin	82657-04-3	ESI + [M+NH4]+		440,1	36	181,2	21	166,2	55	1,33	****
Binapacryl	485-31-4	ESI + [M+NH4]+		340,1	26	83,2	21	54,9	63	1,23	*
Bioresmethrin	28434-01-7	ESI + [M+NH4]+		356,2	21	171,2	21	128,1	53	1,23	n.a.
Bitertanol	55179-31-2	ESI + [M+H]+		338,2	1	70,0	25	269,2	15	1,02	***
Boscalid	188425-85-6	ESI + [M+H]+		343,0	71	307,0	27	139,9	27	0,92	****
Bromacil	314-40-9	ESI + [M+H]+		261,0	21	205,0	19	187,9	37	0,75	***
Bromophos-ethyl	4824-78-6	ESI + [M+H]+		394,9	51	338,7	23	366,9	17	1,20	**
Bromoxynil	1689-84-5	ESI - [M-H]-		273,9	-46	79,0	-36	80,9	-40	0,56	***
Bromuconazole	116255-48-2	ESI + [M+H]+		378,0	46	159,0	37	69,9	35	0,95	***
Bupirimate	41483-43-6	ESI + [M+H]+		317,1	31	166,1	33	108,1	35	0,98	****
Buprofezin	69327-76-0	ESI + [M+H]+		306,2	6	201,2	17	116,2	21	1,09	****
Butafenacil	134605-64-4	ESI + [M+NH4]+		492,1	36	180,1	59	331,0	29	0,95	**
Butocarboxim	34681-10-2	ESI + [M+NH4]+		208,1	1	116,1	11	75,0	15	0,66	****
Butocarboxim-sulfoxide	34681-24-8	ESI + [M+H]+		207,1	6	131,9	11	75,0	19	0,13	****
Butoxycarboxim	34681-23-7	ESI + [M+NH4]+		240,1	6	106,0	19	166,0	13	0,18	**
Buturon	3766-60-7	ESI + [M+H]+		237,1	41	84,1	21	126,1	37	0,85	n.a.
Butylate	2008-41-5	ESI + [M+H]+		218,2	66	57,1	29	156,2	17	1,07	***
Cadusafos	95465-99-9	ESI + [M+H]+		271,1	66	159,0	19	97,0	47	1,05	***
Carbaryl	63-25-2	ESI + [M+H]+		202,1	11	144,9	15	127,0	35	0,79	****
Carbendazim	10605-21-7	ESI + [M+H]+		192,1	41	160,0	25	132,0	41	0,64	****
Carbetamide	16118-49-3	ESI + [M+H]+		237,1	21	118,1	19	192,0	13	0,72	****
Carbofuran	1563-66-2	ESI + [M+H]+		222,1	16	165,1	17	123,0	29	0,75	****
Carbofuran, 3-hydroxy-	16655-82-6	ESI + [M+H]+		238,1	21	181,1	15	163,1	19	0,56	***
Carbosulfan	55285-14-8	ESI + [M+H]+		381,2	36	118,1	25	160,2	21	1,31	***
Carboxin	5234-68-4	ESI + [M+H]+		236,1	26	142,9	21	86,9	33	0,78	****
Carfentrazone-ethyl	128639-02-1	ESI + [M+H]+		412,0	66	365,9	25	345,9	31	0,99	***
Cartap hydrochloride	15263-52-2	ESI + [M+H]+		238,1	26	73,0	37	150,1	19	0,14	**
Chinomethionat	2439-01-2	ESI + [M+H]+		234,9	41	207,1	21	163,0	39	0,72	*
Chlorbromuron	13360-45-7	ESI + [M+H]+		292,9	51	182,1	23	204,0	21	0,92	***
Chlorbufam	1967-16-4	ESI + [M+NH4]+		241,0	6	172,1	17	154,1	27	0,91	***
Chlorfenvinphos	470-90-6	ESI + [M+H]+		358,9	36	155,0	19	99,2	43	1,00	****

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Chlorfluazuron	71422-67-8	ESI + [M+H] ⁺		539,9	71	158,0	27	383,0	27	1,19	***
Chlorfluazuron	71422-67-8	ESI - [M-H] ⁻		538,0	-36	517,9	-18	354,9	-28	1,18	n.a.
Chloridazon	1698-60-8	ESI + [M+H] ⁺		222,0	56	92,2	35	104,2	31	0,58	****
Chlorimuron-ethyl	90982-32-4	ESI + [M+H] ⁺		415,0	46	121,1	53	186,1	27	0,76	***
Chlormephos	24934-91-6	ESI + [M+H] ⁺		234,9	31	97,0	33	143,0	19	0,98	*
Chlorobenzilate	510-15-6	ESI + [M+NH ₄] ⁺		342,0	21	307,1	17	139,1	47	1,05	*
Chlorotoluron	15545-48-9	ESI + [M+H] ⁺		213,1	36	72,0	33	140,0	33	0,83	****
Chloroxuron	1982-47-4	ESI + [M+H] ⁺		291,1	51	72,0	41	218,1	33	0,95	****
Chlorpyrifos	2921-88-2	ESI + [M+H] ⁺		349,9	21	96,9	41	198,0	25	1,13	***
Chlorpyrifos-methyl	5598-13-0	ESI + [M+H] ⁺		321,9	26	125,1	27	289,8	25	1,04	**
Chlorsulfuron	64902-72-3	ESI + [M+H] ⁺		358,0	51	141,0	23	167,0	25	0,59	****
Chlorthiamid	1918-13-4	ESI + [M+H] ⁺		205,9	36	118,9	55	154,0	45	0,57	**
Chlorthiophos	60238-56-4	ESI + [M+H] ⁺		360,9	41	305,0	21	333,0	17	1,16	****
Cinidon-ethyl	142891-20-1	ESI + [M+NH ₄] ⁺		411,1	41	348,0	29	107,2	45	1,11	***
Cinosulfuron	94593-91-6	ESI + [M+H] ⁺		414,1	36	182,9	23	214,9	21	0,62	****
Clethodim	99129-21-2	ESI + [M+H] ⁺		360,1	46	268,2	17	164,0	25	0,92	**
Clethodim	99129-21-2	ESI - [M-H] ⁻		358,1	-66	238,1	-30	no		0,85	n.a.
Clethodim-imin-sulfon	not available	ESI + [M+H] ⁺		302,1	71	98,0	41	208,1	27	0,58	n.a.
Clethodim-imin-sulfoxid	not available	ESI + [M+H] ⁺		286,1	26	208,1	21	166,0	31	0,57	n.a.
Clethodim-sulfon	not available	ESI + [M+H] ⁺		392,1	1	164,2	33	208,1	27	0,67	n.a.
Clethodim-sulfon, 5-hydroxy-	not available	ESI - [M-H] ⁻		406,1	-61	92,8	-30	64,1	-80	0,18	n.a.
Clethodim-sulfon, 5-Hydroxy-	not available	ESI + [M+H] ⁺		408,1	16	204,0	27	176,1	35	0,55	n.a.
Clethodim-sulfoxid	not available	ESI + [M+H] ⁺		376,1	1	206,0	19	164,1	29	0,67	n.a.
Clodinafop-propargyl	105512-06-9	ESI + [M+H] ⁺		350,0	41	266,1	21	91,2	41	0,98	****
Clofentezine	74115-24-5	ESI + [M+H] ⁺		303,1	56	102,1	47	138,1	21	1,02	***
Clomazone	81777-89-1	ESI + [M+H] ⁺		240,1	26	125,0	27	89,1	65	0,88	****
Clopyralid	1702-17-6	ESI + [M+H] ⁺		191,9	36	146,0	29	110,1	47	0,10	**
Cloquintocet-mexyl	99607-70-2	ESI + [M+H] ⁺		336,1	16	192,2	37	238,1	21	1,11	****
Clothianidin	210880-92-5	ESI + [M+H] ⁺		250,0	76	169,1	19	132,1	19	0,51	***
Coumaphos	56-72-4	ESI + [M+H] ⁺		363,0	66	227,0	35	307,0	23	1,01	****
Crotoxyphos	7700-17-6	ESI + [M+NH ₄] ⁺		332,1	1	127,1	33	211,0	15	0,92	****
Cyanazine	21725-46-2	ESI + [M+H] ⁺		241,1	41	214,1	23	104,1	41	0,73	***
Cyanofenphos	13067-93-1	ESI + [M+H] ⁺		304,0	46	276,0	17	157,1	27	1,00	**
Cyanophos	2636-26-2	ESI + [M+NH ₄] ⁺		261,0	6	125,0	27	212,0	27	0,84	*

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Cyazofamid	120116-88-3	ESI + [M+H] ⁺		325,0	36	108,0	19	261,0	15	0,97	****
Cyclanilide	113136--77-9	ESI - [M-H] ⁻		272,0	-56	159,9	-28	228,0	-14	0,82	****
Cycloate	1134-23-2	ESI + [M+H] ⁺		216,1	21	154,3	17	134,2	19	1,05	**
Cycloxydim	101205-02-1	ESI + [M+H] ⁺		326,2	51	280,0	17	180,0	25	0,93	**
Cycloxydim	101205-02-1	ESI - [M-H] ⁻		324,2	-61	235,9	-68	133,8	-68	0,84	n.a.
Cyfluthrin	68359-37-5	ESI + [M+NH ₄] ⁺		451,1	26	191,0	21	127,0	41	1,16	**
Cyhalofop-butyl	122008-85-9	ESI + [M+NH ₄] ⁺		375,1	31	256,1	21	120,0	41	1,05	*
Cyhexatin	13121-70-5	ESI + [M-OH] ⁺		369,2	76	204,8	23	287,0	17	n.a.	n.a.
Cymoxanil	57966-95-7	ESI + [M+H] ⁺		199,1	46	128,0	13	110,9	25	0,61	****
Cypermethrin	52315-07-8	ESI + [M+NH ₄] ⁺		433,1	1	191,0	21	127,0	39	1,16	**
Cyphenothrin[(1R)-trans-isomers]	39515-40-7	ESI + [M+NH ₄] ⁺		393,2	31	151,2	21	123,1	33	1,21	**
Cyproconazole	113096-99-4	ESI + [M+H] ⁺		292,1	16	70,2	35	125,1	35	0,93	***
Cyprodinil	121552-61-2	ESI + [M+H] ⁺		226,1	61	76,9	63	92,9	45	1,01	***
Cyromazine	66215-27-8	ESI + [M+H] ⁺		167,1	46	125,0	25	108,1	29	0,15	***
Daminozide	1596-84-5	ESI + [M+H] ⁺		161,1	46	142,9	15	61,0	19	0,06	**
Deltamethrin	52918-63-5	ESI + [M+NH ₄] ⁺		522,9	16	280,7	23	181,3	51	1,20	***
Demeton-S-methyl	919-86-8	ESI + [M+NH ₄] ⁺		248,0	6	89,1	17	61,0	47	0,77	****
Demeton-S-methyl sulfon	17040-19-6	ESI + [M+H] ⁺		263,0	71	108,9	37	169,0	21	0,30	****
Desmedipham	13684-56-5	ESI + [M+NH ₄] ⁺		318,1	31	182,2	19	136,0	33	0,87	****
Dialifos	10311-84-9	ESI + [M+H] ⁺		394,0	36	208,1	23	186,9	17	1,04	n.a.
Di-allate	2303-16-4	ESI + [M+H] ⁺		270,0	41	86,1	23	108,8	37	1,05	****
Diazinon	333-41-5	ESI + [M+H] ⁺		305,1	21	169,1	29	96,6	41	1,01	****
Dicamba	1918-00-9	ESI - [M-H] ⁻		219,0	-21	175,0	-6	144,8	-14	0,27	*
Dichlofenthion	97-17-6	ESI + [M+H] ⁺		314,9	26	258,9	21	286,9	15	1,00	**
Dichlofluanid	1085-98-9	ESI + [M+NH ₄] ⁺		350,0	21	123,0	41	223,9	21	0,93	***
Dichlorprop-P	15165-67-0	ESI - [M-H] ⁻		233,0	-21	161,0	-14	125,1	-36	0,78	***
Dichlorvos	62-73-7	ESI + [M+H] ⁺		220,9	26	127,1	27	108,9	25	0,74	***
Diclobutrazol	75736-33-3	ESI + [M+H] ⁺		328,1	21	70,1	39	159,0	45	0,99	****
Diclofop-methyl	51338-27-3	ESI + [M+NH ₄] ⁺		358,0	26	281,0	21	120,0	39	1,09	****
Dicloran	99-30-9	ESI - [M-H] ⁻		205,0	-66	175,0	-20	168,9	-24	0,88	*
Dicrotophos	3735-78-3	ESI + [M+H] ⁺		238,1	16	127,1	23	112,1	17	0,45	****
Diethofencarb	87130-20-9	ESI + [M+H] ⁺		268,1	31	226,1	15	180,1	23	0,90	****
Difenoconazole	119446-68-3	ESI + [M+H] ⁺		406,1	41	250,9	37	337,0	23	1,04	****
Difenoxuron	14214-32-5	ESI + [M+H] ⁺		287,1	36	123,1	25	72,1	39	0,87	n.a.

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Diflubenzuron	35367-38-5	ESI - [M-H]-		309,0	-36	155,9	-12	288,9	-8	1,00	***
Diflufenican	83164-33-4	ESI + [M+H]+		395,1	56	265,8	33	246,1	45	1,04	***
Dimefuron	34205-21-5	ESI + [M+H]+		339,1	61	167,0	29	256,0	21	0,88	***
Dimethachlor	50563-36-5	ESI + [M+H]+		256,1	26	224,2	19	148,1	33	0,86	****
Dimethenamide	87674-68-8	ESI + [M+H]+		276,1	11	244,1	19	168,1	33	0,90	****
Dimethoate	60-51-5	ESI + [M+H]+		230,0	11	125,0	29	198,8	13	0,55	****
Dimethomorph	110488-70-5	ESI + [M+H]+		388,1	46	301,1	27	165,0	43	0,92	****
Dimetilan	644-64-4	ESI + [M+H]+		241,1	66	72,1	25	196,0	15	0,59	n.a.
Diniconazole	83657-24-3	ESI + [M+H]+		326,0	56	70,0	45	159,0	39	1,04	***
Dinoseb	88-85-7	ESI - [M-H]-		239,1	-36	134,0	-52	193,0	-30	0,84	***
Dinoterb	1420-07-1	ESI - [M-H]-		239,1	-21	207,0	-32	176,0	-48	0,86	****
Dioxathion	78-34-2	ESI + [M+NH4]+		474,0	31	271,0	19	97,1	61	1,09	****
Diphenylamine	122-39-4	ESI + [M+H]+		170,1	21	93,1	37	151,9	37	0,93	**
Disulfoton	298-04-4	ESI + [M+H]+		275,0	6	89,2	17	61,1	43	1,04	***
Dithianon	3347-22-6	ESI - [M-H]-		296,0	-49	264,0	-24	238,0	-24	0,95	n.a.
Diuron	330-54-1	ESI + [M+H]+		233,0	66	72,0	31	159,9	33	0,86	****
DNOC	534-52-1	ESI - [M-H]-		197,0	-26	137,0	-24	108,8	-28	0,56	***
Dodemorph	1593-77-7	ESI + [M+H]+		282,3	51	116,1	29	98,2	39	1,21	****
Edifenphos	17109-49-8	ESI + [M+NH4]+		328,0	1	109,0	45	283,0	23	1,00	***
Endosulfansulfat	1031-07-8	ESI - [M-H]-		420,8	-66	96,8	-38	79,9	-130	1,02	***
EPN	2104-64-5	ESI + [M+H]+		324,0	46	156,9	29	296,1	19	1,04	**
Epoxiconazole	106325-08-0	ESI + [M+H]+		330,1	36	121,0	27	101,2	63	0,97	****
EPTC	759-94-4	ESI + [M+H]+		190,1	61	128,1	15	86,1	19	0,98	***
Esfenvalerate	66230-04-4	ESI + [M+NH4]+		437,1	41	167,2	23	125,0	51	1,19	*
Ethametsulfuron-methyl	97780-06-8	ESI + [M+H]+		411,1	31	196,1	23	168,1	39	0,68	****
Ethidimuron	30043-49-3	ESI + [M+H]+		265,0	81	208,2	19	114,1	25	0,53	n.a.
Ethiofencarb	29973-13-5	ESI + [M+H]+		226,1	16	107,2	21	164,0	13	0,80	***
Ethiofencarb-sulfone	53380-23-7	ESI + [M+NH4]+		275,3	11	106,9	25	201,0	15	0,45	****
Ethiofencarb-sulfoxide	53380-22-6	ESI + [M+H]+		242,1	41	107,1	23	185,0	13	0,47	****
Ethion	563-12-2	ESI + [M+H]+		385,0	16	199,1	17	171,0	23	1,11	****
Ethirimol	23947-60-6	ESI + [M+H]+		210,2	41	98,1	37	140,0	31	0,85	****
Ethofumesate	26225-79-6	ESI + [M+NH4]+		304,1	36	121,1	27	161,2	31	0,90	***
Ethoprophos	13194-48-4	ESI + [M+H]+		243,0	21	131,0	29	97,0	41	0,96	****
Ethoxyquin	91-53-2	ESI + [M+H]+		218,2	81	160,2	43	174,2	37	0,96	**

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Ethylthiourea (ETU)	96-45-7	ESI + [M+H] ⁺	103,0	31	60,0	47	44,2	27	0,08	**	
Etofenprox	80844-07-1	ESI + [M+NH ₄] ⁺	394,2	16	177,3	21	107,1	53	1,34	****	
Etoxazole	153233-91-1	ESI + [M+H] ⁺	360,2	66	141,0	37	113,0	79	1,15	****	
Etrinfos	38260-54-7	ESI + [M+H] ⁺	293,1	26	125,0	33	265,1	21	1,00	****	
Famoxadone	131807-57-3	ESI + [M+NH ₄] ⁺	392,2	16	238,0	23	330,9	15	1,01	****	
Fenamidone	161326-34-7	ESI + [M+H] ⁺	312,1	41	92,2	33	236,1	19	0,90	****	
Fenamiphos	22224-92-6	ESI + [M+H] ⁺	304,1	41	217,1	31	202,0	45	0,98	****	
Fenarimol	60168-88-9	ESI + [M+H] ⁺	331,0	46	81,0	47	268,0	31	0,96	***	
Fenazaquin	120928-09-8	ESI + [M+H] ⁺	307,2	51	161,2	31	146,9	25	1,28	***	
Fenbuconazole	114369-43-6	ESI + [M+H] ⁺	337,1	41	125,1	37	70,0	33	0,97	****	
Fenbutatin oxide	13356-08-6	ESI + [M-O/2] ⁺	519,2	101	91,1	65	196,9	67	n.a.	n.a.	
Fenfuram	24691-80-3	ESI + [M+H] ⁺	202,1	41	109,0	27	120,1	21	0,80	****	
Fenhexamid	126833-17-8	ESI + [M+H] ⁺	302,1	91	97,2	33	55,1	57	0,95	***	
Fenitrothion	122-14-5	ESI + [M+H] ⁺	278,1	41	125,0	29	108,9	25	0,93	*	
Fenothiocarb	62850-32-2	ESI + [M+H] ⁺	254,1	61	72,1	29	160,1	15	0,99	***	
Fenoxaprop-P-ethyl	71283-80-2	ESI + [M+H] ⁺	362,1	46	288,1	23	121,0	37	1,08	****	
Fenoxycarb	79127-80-3	ESI + [M+H] ⁺	302,1	66	88,0	29	116,0	17	0,98	****	
Fenpiclonil	74738-17-3	ESI + [M+NH ₄] ⁺	253,9	1	202,1	35	140,0	57	0,89	***	
Fenpropathrin	39515-41-8	ESI + [M+H] ⁺	350,2	41	125,1	19	97,2	43	1,17	****	
Fenpropidin	67306-00-7	ESI + [M+H] ⁺	274,2	51	147,1	37	117,1	65	0,91	****	
Fenpropimorph	67306-03-0	ESI + [M+H] ⁺	304,3	46	147,1	39	116,9	71	1,27	****	
Fenpyroximate	111812-58-9	ESI + [M+H] ⁺	422,2	26	366,3	23	135,2	41	1,21	****	
Fenthion	55-38-9	ESI + [M+H] ⁺	279,1	21	169,1	23	247,1	17	1,00	****	
Fentin	668-34-8	ESI + [M] ⁺	351,0	101	196,9	39	119,8	41	1,00	n.a.	
Fenuron	101-42-8	ESI + [M+H] ⁺	165,0	21	72,1	27	120,0	23	0,50	n.a.	
Fenvalerate	51630-58-1	ESI + [M+NH ₄] ⁺	437,1	36	167,0	23	125,0	57	1,19	*	
Fipronil	120068-37-3	ESI + [M+NH ₄] ⁺	453,9	21	368,1	29	255,1	51	0,96	***	
Fipronil	120068-37-3	ESI - [M-H] ⁻	434,9	-31	329,7	-18	249,9	-36	0,98	n.a.	
Fipronil-desulfinyl	205650-65-3	ESI - [M-H] ⁻	387,0	-31	350,8	-18	281,8	-38	0,96	n.a.	
Fipronil-sulfid	120067-83-6	ESI - [M-H] ⁻	418,9	-56	261,9	-34	382,8	-16	0,98	n.a.	
Fipronil-sulfon	120068-36-2	ESI - [M-H] ⁻	450,9	-11	281,9	-34	414,7	-20	1,00	n.a.	
Flamprop-M-isopropyl	63782-90-1	ESI + [M+H] ⁺	364,1	31	77,1	71	105,2	23	0,99	****	
Flamprop-M-methyl	63729-98-6	ESI + [M+H] ⁺	336,0	36	105,1	21	77,1	67	0,93	****	
Flazasulfuron	104040-78-0	ESI + [M+H] ⁺	408,1	41	182,1	25	226,9	25	0,66	****	

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Florasulam	145701-23-1	ESI + [M+H] ⁺		360,0	71	129,1	29	191,9	23	0,55	***
Fluazifop (free acid)	83066-88-0	ESI + [M+H] ⁺		328,1	31	282,0	23	254,1	35	0,80	n.a.
Fluazifop-P-butyl	69806-50-4	ESI + [M+H] ⁺		384,1	51	282,1	27	328,1	23	1,08	****
Fluazifop-P	83066-88-0	ESI - [M-H] ⁻		326,1	-31	253,8	-18	108,1	-50	0,81	***
Fluazinam	79622-59-6	ESI - [M-H] ⁻		462,9	-11	415,8	-24	397,9	-20	1,08	****
Flucythrinate	70124-77-5	ESI + [M+NH ₄] ⁺		469,2	36	181,0	45	199,1	25	1,12	**
Fludioxonil	131341-86-1	ESI - [M-H] ⁻		247,0	-56	125,9	-42	169,0	-42	0,93	***
Flufenacet	142459-58-3	ESI + [M+H] ⁺		364,1	11	194,2	17	152,1	27	0,95	****
Flufenoxuron	101463-69-8	ESI + [M+H] ⁺		489,0	86	158,1	27	141,0	57	1,16	****
Flumioxazin	103361-09-7	ESI + [M+NH ₄] ⁺		372,1	26	327,1	27	76,9	95	0,88	**
Fluometuron	2164-17-2	ESI + [M+H] ⁺		233,1	36	72,0	37	160,2	37	0,82	****
Fluoroglycofene-ethyl	77501-90-7	ESI + [M+NH ₄] ⁺		465,0	16	344,0	21	223,0	43	1,05	***
Flupyr-sulfuron-methyl-sodium	144740-54-5	ESI + [M+H] ⁺		466,1	61	139,1	63	182,2	29	0,70	****
Fluquinconazole	136426-54-5	ESI + [M+H] ⁺		376,0	56	307,1	33	349,0	25	0,95	***
Flurenol	467-69-6	ESI + [M+NH ₄] ⁺		244,1	6	181,2	29	209,2	15	0,27	*
Fluridone	59756-60-4	ESI + [M+H] ⁺		330,1	76	310,2	37	259,1	59	0,89	****
Flurochloridone	61213-25-0	ESI + [M+H] ⁺		312,0	61	291,9	29	145,0	63	0,92	**
Fluroxypyr	69377-81-7	ESI + [M+H] ⁺		255,0	71	209,1	21	181,1	31	0,40	*
Fluroxypyr-meptyl	81406-37-3	ESI + [M+H] ⁺		367,1	31	255,0	17	209,0	29	1,15	****
Flurtamone	96525-23-4	ESI + [M+H] ⁺		334,1	51	247,1	35	178,1	57	0,90	n.a.
Flusilazole	85509-19-9	ESI + [M+H] ⁺		316,1	36	247,1	25	165,0	35	0,98	****
Flutolanil	66332-96-5	ESI + [M+H] ⁺		324,1	86	262,1	25	242,1	35	0,92	****
Flutriafol	76674-21-0	ESI + [M+H] ⁺		302,1	41	122,9	39	109,0	43	0,84	***
Folpet	133-07-3	ESI + [M+NH ₄] ⁺		314,9	1	130,1	39	163,0	19	0,79	*
Fomesafen	72178-02-0	ESI + [M+NH ₄] ⁺		456,0	51	344,1	21	222,9	45	0,88	n.a.
Fonofos	944-22-9	ESI + [M+H] ⁺		247,0	11	109,1	25	137,0	15	1,01	***
Foramsulfuron	173159-57-4	ESI + [M+H] ⁺		453,1	51	182,2	27	139,2	63	0,64	***
Formetanate	22259-30-9	ESI + [M+H] ⁺		222,1	21	165,1	19	120,1	35	0,73	**
Fosthiazate	98886-44-3	ESI + [M+H] ⁺		284,0	61	104,1	27	227,8	15	0,81	****
Fuberidazole	3878-19-1	ESI + [M+H] ⁺		185,1	26	157,1	31	65,1	59	0,76	****
Furathiocarb	65907-30-4	ESI + [M+H] ⁺		383,2	51	195,0	23	251,9	19	1,09	****
Glufosinate	77182-82-2	ESI + [M+H] ⁺		182,1	31	136,1	19	119,0	25	0,06	n.a.
Halfenprox	111872-58-3	ESI + [M+NH ₄] ⁺		494,1	46	183,1	27	168,0	75	1,41	**
Halofenozide	112226-61-6	ESI + [M+NH ₄] ⁺		348,1	1	105,0	27	275,2	17	0,91	**

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Halosulfuron-methyl	100784-20-1	ESI + [M+H] ⁺		435,0	56	182,1	27	83,1	73	0,70	***
Haloxypop-etotyl	87237-48-7	ESI + [M+H] ⁺		434,1	81	315,9	25	287,9	35	1,07	****
Haloxypop-P	95977-29-0	ESI + [M+NH ₄] ⁺		379,0	36	316,0	29	91,1	47	0,90	**
Haloxypop-P	95977-29-0	ESI - [M-H] ⁻		360,0	-81	287,9	-14	no		0,91	n.a.
Haloxypop-P-methyl	72619-32-0	ESI + [M+H] ⁺		376,1	91	315,9	23	287,8	33	1,04	****
Heptenophos	23560-59-0	ESI + [M+H] ⁺		251,0	36	127,0	19	109,0	37	0,86	****
Hexaconazole	79983-71-4	ESI + [M+H] ⁺		314,1	36	70,1	39	159,0	37	1,02	****
Hexaflumuron	86479-06-3	ESI - [M-H] ⁻		459,0	-6	438,8	-14	275,9	-22	1,07	***
Hexazinone	51235-04-2	ESI + [M+H] ⁺		253,2	16	171,1	21	71,1	47	0,76	****
Hexythiazox	78587-05-0	ESI + [M+H] ⁺		353,1	66	227,9	21	168,1	33	1,13	****
Imazalil	35554-44-0	ESI + [M+H] ⁺		297,0	26	158,9	31	200,9	23	1,00	****
Imazapyr	81334-34-1	ESI + [M+H] ⁺		262,1	56	217,1	27	149,1	35	0,11	n.a.
Imazaquin	81335-37-7	ESI + [M+H] ⁺		312,1	46	199,1	37	128,1	69	0,52	n.a.
Imazethapyr	81335-77-5	ESI + [M+H] ⁺		290,1	51	177,2	37	245,3	27	0,31	n.a.
Imibenconazole	86598-92-7	ESI + [M+H] ⁺		411,0	71	125,1	39	171,1	27	1,11	***
Imidacloprid	138261-41-3	ESI + [M+H] ⁺		256,1	46	175,0	25	208,9	21	0,51	***
Imidacloprid, 5-hydroxide	not available	ESI + [M+H] ⁺		272,0	56	191,1	23	224,8	23	0,43	**
Imidacloprid, olefine	not available	ESI + [M+H] ⁺		254,0	56	204,9	21	125,8	39	0,38	**
Indoxacarb	173584-44-6	ESI + [M+H] ⁺		528,1	76	203,0	51	56,0	55	1,04	***
Iodosulfuron-methyl-sodium	185119-76-0	ESI + [M+H] ⁺		507,9	36	167,2	27	141,0	35	0,72	***
Ioxynil	1689-83-4	ESI - [M-H] ⁻		369,8	-46	126,8	-36	242,8	-26	0,72	****
Iprobenfos	26087-47-8	ESI + [M+NH ₄] ⁺		306,1	1	91,1	35	205,1	19	0,99	****
Iprodione	36734-19-7	ESI + [M+H] ⁺		330,0	56	101,0	33	143,2	21	1,05	n.a.
Iprodione	36734-19-7	ESI - [M-H] ⁻		328,0	-1	141,1	-14	no		1,07	***
Iprovalicarb	140923-17-7	ESI + [M+H] ⁺		321,2	46	119,0	23	202,9	13	0,95	****
Isazofos	42509-80-8	ESI + [M+H] ⁺		314,0	41	120,0	35	162,2	21	0,94	n.a.
Isofenphos	25311-71-1	ESI + [M+H] ⁺		346,1	56	217,0	33	245,0	17	1,02	****
Isofenphos-oxon	31120-85-1	ESI + [M+H] ⁺		330,1	1	229,1	17	201,0	29	0,95	****
Isoprocab	2631-40-5	ESI + [M+NH ₄] ⁺		211,3	31	95,0	25	137,2	17	0,83	***
Isoprothiolane	50512-35-1	ESI + [M+H] ⁺		291,1	11	231,1	17	189,0	27	0,93	***
Isoproturon	34123-59-6	ESI + [M+H] ⁺		207,1	46	165,2	19	72,0	33	0,85	***
Isoxadifen-ethyl	163520-33-0	ESI + [M+NH ₄] ⁺		313,2	11	232,1	27	204,2	39	0,99	***
Isoxaflutole	141112-29-0	ESI + [M+NH ₄] ⁺		377,0	26	251,0	25	no		0,85	***
Isoxathion	18854-01-8	ESI + [M+H] ⁺		314,1	46	105,0	21	170,0	19	1,02	****

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Kresoxim-methyl	143390-89-0	ESI + [M+H] ⁺		314,1	16	115,9	21	206,1	13	0,99	***
lambda-Cyhalothrin	91465-08-6	ESI + [M+NH ₄] ⁺		467,1	16	225,0	23	141,2	57	1,15	**
Lenacil	2164-08-1	ESI + [M+H] ⁺		235,1	56	153,1	21	136,1	43	0,84	****
Linuron	330-55-2	ESI + [M+H] ⁺		249,0	66	159,9	23	181,9	21	0,91	****
Lufenuron	103055-07-8	ESI + [M+H] ⁺		510,9	61	158,2	27	141,2	67	1,10	***
Malaoxon	1634-78-2	ESI + [M+H] ⁺		315,0	31	127,1	17	99,2	31	0,77	****
Malathion	121-75-5	ESI + [M+H] ⁺		331,0	26	127,0	17	99,0	29	0,93	****
Maleic hydrazide	123-33-1	ESI - [M-H] ⁻		111,0	-76	81,9	-20	55,0	-20	n.a.	n.a.
Maleic hydrazide	123-33-1	ESI + [M+H] ⁺		113,0	76	85,2	23	66,9	27	0,06	n.a.
MCPA	94-74-6	ESI - [M-H] ⁻		199,0	-46	140,8	-18	no		0,70	***
MCPA, 2-Ethylhexyl-	29450-45-1	ESI + [M+NH ₄] ⁺		330,2	11	155,0	23	201,0	15	1,31	**
MCPA, Butotyl-	19480-43-4	ESI + [M+NH ₄] ⁺		318,1	6	227,0	17	101,2	19	1,07	***
MCPB	94-81-5	ESI - [M-H] ⁻		227,0	-26	140,9	-10	104,8	-36	0,86	***
Mecarbam	2595-54-2	ESI + [M+H] ⁺		330,0	26	227,0	15	97,1	45	0,96	***
Mecoprop-P	16484-77-8	ESI - [M-H] ⁻		213,0	-51	140,7	-14	no		0,77	***
Mefenpyr-diethyl	135590-91-9	ESI + [M+NH ₄] ⁺		390,1	1	327,1	23	160,1	49	1,02	****
Mepanipyrim	110235-47-7	ESI + [M+H] ⁺		224,1	31	77,0	49	106,0	35	0,96	****
Mepiquat	24307-26-4	ESI + [M] ⁺		114,1	6	58,1	37	70,1	45	0,63	n.a.
Mepronil	55814-41-0	ESI + [M+H] ⁺		270,1	76	119,0	31	91,2	55	0,93	****
Mesosulfuron-methyl	208465-21-8	ESI + [M+H] ⁺		504,1	61	182,1	33	83,0	75	0,71	****
Mesotrione	104206-82-8	ESI + [M+NH ₄] ⁺		357,1	21	227,8	29	104,1	49	0,17	***
Metalaxyl-M	70630-17-0	ESI + [M+H] ⁺		280,1	46	220,0	19	159,9	31	0,85	****
Metamitron	41394-05-2	ESI + [M+H] ⁺		203,1	46	175,0	29	104,1	29	0,54	***
Metazachlor	67129-08-2	ESI + [M+H] ⁺		278,1	1	210,1	15	134,2	29	0,84	****
Metconazole	125116-23-6	ESI + [M+H] ⁺		320,1	31	70,1	45	125,0	49	1,02	****
Methabenzthiazuron	18691-97-9	ESI + [M+H] ⁺		222,1	11	165,1	23	150,0	43	0,84	****
Methacrifos	30864-28-9	ESI + [M+H] ⁺		241,0	31	209,1	13	125,1	25	0,86	**
Methamidophos	10265-92-6	ESI + [M+H] ⁺		142,0	26	124,9	19	93,9	19	0,09	****
Methfuroxam	28730-17-8	ESI + [M+H] ⁺		230,0	26	137,1	27	111,1	21	0,88	n.a.
Methidathion	950-37-8	ESI + [M+H] ⁺		302,9	16	145,1	15	85,1	27	0,88	****
Methiocarb	2032-65-7	ESI + [M+H] ⁺		226,1	61	121,0	25	169,0	17	0,91	****
Methiocarb-sulfone	2179-25-1	ESI + [M+NH ₄] ⁺		275,1	16	122,0	29	201,1	17	1,21	**
Methomyl	16752-77-5	ESI + [M+H] ⁺		163,0	46	106,0	13	88,0	13	0,27	***
Methomyl-oxime	13749-94-5	ESI + [M+H] ⁺		106,0	41	57,9	19	73,9	13	0,17	n.a.

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Methoxyfenozide	161050-58-4	ESI + [M+H] ⁺		369,2	36	149,1	23	133,0	31	0,94	****
Methoxyfenozide	161050-58-4	ESI - [M-H] ⁻		367,2	-66	148,8	-24	no		0,94	n.a.
Metobromuron	3060-89-7	ESI + [M+H] ⁺		259,0	56	170,0	25	147,9	21	0,83	****
Metolcarb	1129-41-5	ESI + [M+H] ⁺		166,1	61	109,0	17	94,1	43	0,72	***
Metosulam	139528-85-1	ESI + [M+H] ⁺		418,0	26	175,1	27	140,1	69	0,76	****
Metoxuron	19937-59-8	ESI + [M+H] ⁺		229,0	26	156,1	31	72,1	35	0,69	****
Metrafenone	220899-03-6	ESI + [M+H] ⁺		409,1	41	209,1	21	226,9	25	1,04	****
Metribuzin	21087-64-9	ESI + [M+H] ⁺		215,1	31	187,2	25	84,0	29	0,74	***
Metsulfuron-methyl	74223-64-6	ESI + [M+H] ⁺		382,1	31	198,9	27	167,0	21	0,52	****
Mevinphos	7786-34-7	ESI + [M+H] ⁺		225,0	31	193,1	13	127,0	21	0,65	****
Molinate	2212-67-1	ESI + [M+H] ⁺		188,1	21	83,2	25	126,1	19	0,94	***
Monocrotophos	6923-22-4	ESI + [M+H] ⁺		224,1	46	127,0	21	98,0	17	0,39	***
Monolinuron	1746-81-2	ESI + [M+H] ⁺		215,1	61	125,9	25	148,0	19	0,80	****
Monuron	150-68-5	ESI + [M+H] ⁺		199,1	31	72,0	29	126,0	35	0,74	****
Myclobutanil	88671-89-0	ESI + [M+H] ⁺		289,1	36	70,1	33	125,1	41	0,94	***
Naled	300-76-5	ESI + [M+NH ₄] ⁺		397,7	1	127,1	25	108,9	53	0,86	***
Napropamide	15299-99-7	ESI + [M+H] ⁺		272,1	31	129,3	21	171,1	23	0,96	****
Neburon	555-37-3	ESI + [M+H] ⁺		275,1	41	88,1	23	114,1	21	0,99	****
Nicosulfuron	111991-09-4	ESI + [M+H] ⁺		411,1	61	182,1	25	213,0	23	0,44	***
Nicotine	54-11-5	ESI + [M+H] ⁺		163,1	16	132,0	21	84,1	25	0,33	**
Nitenpyram	120738-89-8	ESI + [M+H] ⁺		271,1	51	126,1	39	237,2	25	0,17	***
Nitrothal-isopropyl	10552-74-6	ESI + [M+NH ₄] ⁺		313,1	11	211,8	29	253,9	17	1,07	n.a.
Norflurazon	27314-13-2	ESI + [M+H] ⁺		304,0	81	284,1	31	160,1	41	0,86	****
Norflurazon-desmethyl	23576-24-1	ESI + [M+H] ⁺		290,0	96	270,0	29	160,1	41	0,82	****
Nuarimol	63284-71-9	ESI + [M+H] ⁺		315,0	41	81,1	43	252,1	31	0,91	***
Ofurace	58810-48-3	ESI + [M+H] ⁺		282,1	41	254,0	17	160,1	29	0,76	****
Omethoate	1113-02-6	ESI + [M+H] ⁺		214,1	51	109,0	35	125,0	29	0,13	****
Oxadiazon	19666-30-9	ESI + [M+NH ₄] ⁺		362,1	41	220,0	31	177,1	45	1,09	****
Oxadixyl	77732-09-3	ESI + [M+H] ⁺		279,1	46	219,2	17	133,3	29	0,71	***
Oxamyl	23135-22-0	ESI + [M+NH ₄] ⁺		237,1	1	72,0	21	90,0	13	0,19	****
Oxamyl-oxime	30558-43-1	ESI + [M+H] ⁺		163,0	46	71,9	19	89,9	23	0,13	n.a.
Oxasulfuron	144651-06-9	ESI + [M+H] ⁺		407,1	56	150,1	25	107,1	63	0,58	****
Oxycarboxin	5259-88-1	ESI + [M+H] ⁺		268,0	36	175,1	19	147,1	29	0,61	****
Oxydemeton-methyl	301-12-2	ESI + [M+H] ⁺		247,0	21	169,0	19	109,0	35	0,24	****

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Oxyfluorfen	42874-03-3	ESI + [M+NH4] ⁺		379,0	21	316,0	23	237,1	39	1,08	*
Paclobutrazol	76738-62-0	ESI + [M+H] ⁺		294,1	36	70,1	39	125,2	49	0,92	****
Paraoxon	311-45-5	ESI + [M+H] ⁺		276,1	76	220,0	19	94,2	45	0,83	****
Paraoxon-methyl	950-35-6	ESI + [M+H] ⁺		248,0	51	109,1	35	202,1	25	0,71	***
Parathion	56-38-2	ESI + [M+H] ⁺		292,0	51	236,1	21	97,0	39	0,97	**
Parathion-methyl	298-00-0	ESI + [M+H] ⁺		264,0	56	125,0	25	232,1	23	0,89	*
Penconazole	66246-88-6	ESI + [M+H] ⁺		284,1	41	158,9	39	70,1	29	1,00	***
Pencycuron	66063-05-6	ESI + [M+H] ⁺		329,1	41	125,1	33	99,1	83	1,03	****
Pendimethalin	40487-42-1	ESI + [M+H] ⁺		282,1	6	212,2	15	194,0	23	1,14	***
Permethrin	52645-53-1	ESI + [M+NH4] ⁺		408,1	31	183,1	25	153,1	63	1,28	***
Phenmedipham	13684-63-4	ESI + [M+H] ⁺		301,1	56	136,1	25	168,0	15	0,88	****
Phenthoate	2597-03-7	ESI + [M+H] ⁺		321,0	36	163,1	17	79,1	51	0,99	***
Phorate	298-02-2	ESI + [M+NH4] ⁺		278,1	1	74,9	23	170,8	21	1,01	**
Phorat-sulfone	2588-04-7	ESI + [M+H] ⁺		293,0	36	97,0	41	171,1	17	0,82	****
Phorat-sulfoxide	2588-03-6	ESI + [M+H] ⁺		277,0	51	199,1	15	143,0	25	0,82	****
Phosalone	2310-17-0	ESI + [M+H] ⁺		367,9	51	182,0	21	110,9	51	1,02	****
Phosmet	732-11-6	ESI + [M+H] ⁺		317,9	31	133,1	49	160,1	19	0,89	***
Phosphamidon	13171-21-6	ESI + [M+H] ⁺		300,0	36	127,1	27	174,1	19	0,71	****
Phoxim	14816-18-3	ESI + [M+H] ⁺		299,0	26	129,1	17	77,1	41	1,01	****
Picloram	1918-02-1	ESI - [M-H] ⁻		240,9	-66	196,9	-14	122,6	-30	0,10	n.a.
Picolinafen	137641-05-5	ESI + [M+H] ⁺		377,1	56	145,0	69	238,1	35	1,10	****
Picoxystrobin	117428-22-5	ESI + [M+H] ⁺		368,1	36	145,0	27	205,2	15	0,98	****
Piperonyl butoxide	51-03-6	ESI + [M+NH4] ⁺		356,2	11	177,1	19	119,1	47	1,10	****
Pirimicarb	23103-98-2	ESI + [M+H] ⁺		239,1	16	72,1	31	181,9	21	0,82	****
Pirimicarb, desmethyl-	30614-22-3	ESI + [M+H] ⁺		225,1	16	72,0	27	168,1	19	0,69	****
Pirimicarb, desmethyl-formamido-	59333-83-4	ESI + [M+H] ⁺		253,1	11	72,0	25	225,1	15	0,76	****
Pirimiphos-ethyl	23505-41-1	ESI + [M+H] ⁺		334,1	16	198,1	29	182,2	27	1,10	****
Pirimiphos-methyl	29232-93-7	ESI + [M+H] ⁺		306,1	26	164,1	29	108,1	39	1,02	****
Primisulfuron-methyl	86209-51-0	ESI + [M+H] ⁺		469,0	71	253,8	25	199,0	25	0,82	***
Prochloraz	67747-09-5	ESI + [M+H] ⁺		376,0	16	308,0	17	265,9	23	1,02	****
Procymidone	32809-16-8	ESI + [M+NH4] ⁺		301,0	6	256,0	29	no		0,74	*
Profenofos	41198-08-7	ESI + [M+H] ⁺		372,9	56	302,9	25	97,0	43	1,07	****
Prohexadione	88805-35-0	ESI - [M-H] ⁻		211,1	-21	123,1	-16	166,9	-16	0,10	n.a.
Promecarb	2631-37-0	ESI + [M+H] ⁺		208,1	11	108,9	21	150,9	13	0,92	****

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Prometon	1610-18-0	ESI + [M+H] ⁺		226,1	21	142,0	29	184,3	23	0,89	****
Prometryn	7287-19-6	ESI + [M+H] ⁺		242,1	31	158,1	31	200,2	25	0,95	****
Propachlor	1918-16-7	ESI + [M+H] ⁺		212,1	36	170,0	21	94,1	37	0,84	****
Propamocarb hydrochloride	24579-73-5	ESI + [M+H] ⁺		189,2	16	102,0	23	144,0	17	0,19	***
Propaquizafop	111479-05-1	ESI + [M+H] ⁺		444,1	76	299,1	31	100,0	27	1,09	****
Propargite	2312-35-8	ESI + [M+NH ₄] ⁺		368,1	1	175,1	21	231,2	17	1,15	****
Propazin, 2-hydroxy-	not available	ESI + [M+H] ⁺		212,1	46	128,1	31	170,2	25	0,76	****
Propazine	139-40-2	ESI + [M+H] ⁺		230,1	31	146,0	29	188,0	23	0,89	****
Propetamphos	31218-83-4	ESI + [M+H] ⁺		282,1	31	156,2	17	138,0	21	0,93	****
Propham	122-42-9	ESI + [M+H] ⁺		180,1	16	138,1	13	120,1	25	0,83	***
Propiconazole	60207-90-1	ESI + [M+H] ⁺		342,1	46	69,1	33	159,0	37	1,01	***
Propoxur	114-26-1	ESI + [M+H] ⁺		210,1	11	111,0	19	168,0	11	0,74	****
Propoxycarbazone-sodium	181274-15-7	ESI + [M+NH ₄] ⁺		416,1	11	116,0	41	199,1	23	0,61	***
Propyzamide	23950-58-5	ESI + [M+H] ⁺		256,0	36	173,1	31	190,0	19	0,93	****
Prosulfocarb	52888-80-9	ESI + [M+H] ⁺		252,1	36	91,2	29	128,2	17	1,06	****
Prosulfuron	94125-34-5	ESI + [M+H] ⁺		420,1	56	141,1	27	167,0	25	0,79	***
Prothioconazole	178928-70-6	ESI + [M+H] ⁺		344,0	46	125,0	39	189,1	27	1,01	n.a.
Prothiofos	34643-46-4	ESI + [M+H] ⁺		344,9	41	240,9	27	133,0	69	1,21	**
Pymetrozine	123312-89-0	ESI + [M+H] ⁺		218,1	56	105,0	27	78,9	47	0,40	****
Pyraclofos	89784-60-1	ESI + [M+H] ⁺		361,1	76	138,1	49	111,0	79	1,03	****
Pyraclostrobin	175013-18-0	ESI + [M+H] ⁺		388,1	6	194,0	19	163,0	29	1,02	****
Pyraflufen-ethyl	129630-17-7	ESI + [M+H] ⁺		413,0	91	339,0	25	253,0	43	1,00	***
Pyrazophos	13457-18-6	ESI + [M+H] ⁺		374,1	61	222,1	29	194,1	43	1,04	****
Pyridaben	96489-71-3	ESI + [M+H] ⁺		365,1	26	309,1	19	147,2	31	1,24	****
Pyridaphenthion	119-12-0	ESI + [M+H] ⁺		341,0	46	189,0	29	205,1	27	0,94	****
Pyridate	55512-33-9	ESI + [M+H] ⁺		379,1	6	207,1	21	351,1	17	1,29	****
Pyridat-metabolite (6-Chlor-3-phenyl-pyridazin-4-ol)	40020-01-7	ESI + [M+H] ⁺		207,0	66	104,0	31	77,1	43	0,39	***
Pyrifenox	88283-41-4	ESI + [M+H] ⁺		295,0	16	93,1	31	263,1	25	0,98	***
Pyrimethanil	53112-28-0	ESI + [M+H] ⁺		200,1	51	82,0	35	106,9	33	0,91	****
Pyriproxyfen	95737-68-1	ESI + [M+H] ⁺		322,1	16	96,2	21	185,1	29	1,11	****
Quinalphos	13593-03-8	ESI + [M+H] ⁺		299,0	21	163,0	29	147,0	29	0,99	****
Quinmerac	90717-03-6	ESI + [M+H] ⁺		222,0	21	204,1	23	141,0	43	0,32	****
Quinoclamine	2797-51-5	ESI + [M+H] ⁺		208,0	26	77,0	49	105,2	33	0,74	**
Quinoxifen	124495-18-7	ESI + [M+H] ⁺		307,9	21	162,0	57	197,1	43	1,14	****

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Quizalofop	76578-12-6	ESI - [M-H]-		343,1	-86	270,9	-18	242,9	-34	0,89	n.a.
Quizalofop-ethyl	76578-14-8	ESI + [M+H]+		373,1	71	298,9	25	271,0	33	1,08	****
Quizalofop-P	94051-08-8	ESI + [M+H]+		345,1	81	299,1	25	162,9	49	0,89	**
Resmethrin	10453-86-8	ESI + [M+NH4]+		356,2	21	171,2	21	128,1	57	1,23	****
Rimsulfuron	122931-48-0	ESI + [M+H]+		432,1	46	182,0	29	324,9	21	0,54	****
Rotenone	83-79-4	ESI + [M+H]+		395,1	66	213,0	29	192,2	31	0,98	n.a.
Sebuthylazine	7286-69-3	ESI + [M+H]+		230,1	21	174,2	25	104,2	41	0,91	****
Sebuthylazine, desethyl-	not available	ESI + [M+H]+		202,1	51	145,9	25	104,0	35	0,78	****
Sethoxydim	74051-80-2	ESI + [M+H]+		328,1	16	282,2	17	178,1	25	0,97	**
Siduron	1982-49-6	ESI + [M+H]+		233,2	56	137,1	25	94,0	33	0,91	****
Silthiofam	175217-20-6	ESI + [M+H]+		268,1	36	139,1	25	73,0	39	0,98	****
Simazine	122-34-9	ESI + [M+H]+		202,1	26	124,2	25	132,2	27	0,75	***
Simazine, 2-hydroxy-	2599-11-3	ESI + [M+H]+		184,1	51	69,0	59	114,1	27	0,48	****
Simetryn	1014-70-6	ESI + [M+H]+		214,1	31	124,2	27	144,0	27	0,83	****
S-Metolachlor	51218-45-2	ESI + [M+H]+		284,1	16	251,9	19	176,1	35	0,97	****
Spinosyn A	131929-60-7	ESI + [M+H]+		732,5	51	142,1	37	98,3	75	1,22	****
Spinosyn D	131929-63-0	ESI + [M+H]+		746,5	66	142,2	39	98,1	79	1,29	***
Spiroxamine	118134-30-8	ESI + [M+H]+		298,3	41	144,2	27	100,1	41	0,99	****
Sulfentrazone	122836-35-5	ESI - [M-H]-		385,0	-56	307,1	-30	198,9	-44	0,77	***
Sulfometuron-methyl	74222-97-2	ESI + [M+H]+		365,1	51	150,1	23	107,1	29	0,60	****
Sulfosulfuron	141776-32-1	ESI + [M+H]+		471,1	11	261,0	23	211,1	21	0,62	****
Sulfotep	3689-24-5	ESI + [M+H]+		323,0	46	115,0	39	97,1	45	0,98	****
Sulprofos	35400-43-2	ESI + [M+H]+		323,0	26	247,1	17	219,0	21	1,13	n.a.
tau-Fluvalinate	102851-06-9	ESI + [M+NH4]+		520,1	31	208,1	23	181,1	41	1,23	***
Tebuconazole	107534-96-3	ESI + [M+H]+		308,1	21	70,0	39	124,9	47	0,99	****
Tebufenozide	112410-23-8	ESI + [M+H]+		353,2	41	296,9	15	133,0	23	0,97	****
Tebufenpyrad	119168-77-3	ESI + [M+H]+		334,2	51	117,0	47	145,0	37	1,08	****
Tebutam	35256-85-0	ESI + [M+H]+		234,2	21	91,0	29	192,2	19	0,96	****
Tebuthiuron	34014-18-1	ESI + [M+H]+		229,1	26	172,2	23	116,0	35	0,77	n.a.
Teflubenzuron	83121-18-0	ESI - [M-H]-		379,0	-6	338,9	-12	358,8	-8	1,12	***
Tefluthrin	79538-32-2	ESI + [M+NH4]+		436,1	31	177,1	29	127,0	79	1,19	n.a.
TEPP	107-49-3	ESI + [M+H]+		291,1	46	179,0	27	99,0	49	0,71	****
Tepraloxydim	149979-41-9	ESI + [M+H]+		342,1	56	250,1	19	166,1	29	0,75	****
Terbacil	5902-51-2	ESI + [M+H]+		217,0	21	161,0	17	144,0	35	0,76	**

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Terbufos	13071-79-9	ESI + [M+H] ⁺		289,0	31	57,1	35	103,3	15	1,09	****
Terbumeton	33693-04-8	ESI + [M+H] ⁺		226,1	16	170,2	23	114,0	33	0,91	****
Terbuthylazine	5915-41-3	ESI + [M+H] ⁺		230,1	41	174,1	23	104,2	43	0,90	****
Terbuthylazine, 2-hydroxy-	not available	ESI + [M+H] ⁺		212,1	31	156,0	23	114,0	35	0,79	****
Terbuthylazine, desethyl-	30125-63-4	ESI + [M+H] ⁺		202,1	26	145,9	23	104,2	37	0,78	****
Terbutryn	886-50-0	ESI + [M+H] ⁺		242,1	21	186,1	25	68,1	57	0,96	****
Tetrachlorvinphos	22248-79-9	ESI + [M+H] ⁺		366,9	46	127,1	21	241,0	27	0,97	****
Tetraconazole	112281-77-3	ESI + [M+H] ⁺		372,0	46	159,0	39	70,0	47	0,95	****
Tetramethrin [(1R)-isomers]	7696-12-0	ESI + [M+NH ₄] ⁺		349,2	26	164,2	29	135,2	23	1,09	***
Thiabendazol, 5-hydroxy-	948-71-0	ESI + [M+H] ⁺		218,0	71	190,9	35	146,9	43	0,59	n.a.
Thiabendazole	148-79-8	ESI + [M+H] ⁺		202,0	61	131,1	43	174,9	35	0,72	***
Thiacloprid	111988-49-9	ESI + [M+H] ⁺		253,0	81	126,0	29	186,0	19	0,65	****
Thiamethoxam	153719-23-4	ESI + [M+H] ⁺		292,0	51	211,0	17	181,0	31	0,30	****
Thifensulfuron-methyl	79277-27-3	ESI + [M+H] ⁺		388,0	36	167,0	21	204,9	33	0,51	****
Thiodicarb	59669-26-0	ESI + [M+H] ⁺		355,0	26	88,0	21	107,8	21	0,82	****
Thiofanox	39196-18-4	ESI + [M+H] ⁺		219,1	16	57,0	17	60,9	15	0,80	****
Thiofanox-sulfone	39184-59-3	ESI + [M+NH ₄] ⁺		268,1	16	57,0	29	75,9	17	0,50	***
Thiofanox-sulfoxide	39184-27-5	ESI + [M+NH ₄] ⁺		252,1	11	104,0	17	57,2	27	0,44	***
Thiometon	640-15-3	ESI + [M+H] ⁺		246,9	26	89,1	17	61,1	49	0,81	**
Thiophanate (-ethyl)	23564-06-9	ESI + [M+H] ⁺		371,1	41	151,0	27	325,1	17	0,86	n.a.
Thiophanate-methyl	23564-05-8	ESI + [M+H] ⁺		343,0	26	151,0	25	192,0	21	0,75	****
Tolclofos-methyl	57018-04-9	ESI + [M+H] ⁺		301,0	46	268,9	23	175,0	35	1,02	**
Tolyfluanid	731-27-1	ESI + [M+NH ₄] ⁺		364,0	6	237,9	19	137,1	37	0,98	****
Transfluthrin	118712-89-3	ESI + [M+NH ₄] ⁺		388,0	1	82,0	17	162,9	33	1,12	n.a.
Triadimefon	43121-43-3	ESI + [M+H] ⁺		294,0	36	197,2	21	225,1	19	0,93	***
Triadimenol	55219-65-3	ESI + [M+H] ⁺		296,1	11	70,1	19	227,2	15	0,94	***
Tri-allate	2303-17-5	ESI + [M+H] ⁺		304,0	41	142,9	35	86,1	23	1,10	***
Triasulfuron	82097-50-5	ESI + [M+H] ⁺		402,1	46	167,1	25	140,8	29	0,65	****
Triazamate	112143-82-5	ESI + [M+H] ⁺		315,1	26	72,1	33	226,0	17	0,95	***
Triazophos	24017-47-8	ESI + [M+H] ⁺		314,0	36	119,1	47	162,1	25	0,94	****
Tribenuron-methyl	101200-48-0	ESI + [M+H] ⁺		396,1	51	155,0	21	180,9	27	0,67	****
Trichlorfon	52-68-6	ESI + [M+NH ₄] ⁺		274,0	6	108,9	31	221,0	21	0,50	***
Triclopyr	55335-06-3	ESI - [M-H] ⁻		255,9	-16	197,9	-12	219,8	-6	0,75	***
Tricyclazole	41814-78-2	ESI + [M+H] ⁺		190,0	46	163,1	31	136,2	37	0,67	****

Pesticide (Metabolite)	CAS No.	Ionization	Quasi molecular ion	Q1 Mass (amu)	Declustering potential (V)	1 st SRM		2 nd SRM		Relative retention on endcapped RP phase ^a	Sensitivity of detection ^b
						Q3 Mass (amu)	Collision energy (V)	Q3 Mass (amu)	Collision energy (V)		
Tridemorph	24602-86-6	ESI + [M+H] ⁺		298,3	56	130,1	35	116,1	33	1,34	**
Trietazine	1912-26-1	ESI + [M+H] ⁺		230,1	26	99,1	33	132,1	29	0,95	****
Trifloxystrobin	141517-21-7	ESI + [M+H] ⁺		409,1	11	186,1	23	206,1	21	1,03	****
Triflumizole	68694-11-1	ESI + [M+H] ⁺		346,0	6	278,0	17	73,1	23	1,05	****
Triflumuron	64628-44-0	ESI - [M-H] ⁻		357,0	-16	154,0	-14	175,9	-22	1,03	****
Triflusulfuron-methyl	126535-15-7	ESI + [M+H] ⁺		493,1	46	264,0	29	238,0	29	0,83	***
Trinexapac-ethyl	95266-40-3	ESI + [M+H] ⁺		253,1	71	68,8	29	207,0	17	0,57	**
Tris-(1,3-dichloro-isopropyl)-phosphat	13674-87-8	ESI + [M+H] ⁺		430,9	66	98,9	31	209,0	21	n.a.	n.a.
Triticonazole	131983-72-7	ESI + [M+H] ⁺		318,1	36	70,2	33	125,2	41	0,95	****
Tritosulfuron	142469-14-5	ESI + [M+NH ₄] ⁺		463,1	1	195,0	29	145,0	59	0,76	**
Uniconazole	83657-22-1	ESI + [M+H] ⁺		292,1	31	70,1	37	125,1	37	0,98	***
Vamidothion	2275-23-2	ESI + [M+H] ⁺		288,0	16	145,9	17	117,9	31	0,55	****

^a Relative retention time is based on retention time of imazalil. If more than one peak exists, RRT of the most intense peak is given. In the case of "n.a." no data available.

^b Four stars means very sensitive detectable, one star means detection with lowest sensitivity. In the case of "n.a." no data available.

Bibliography

- [1] <http://www.bfr.bund.de/cd/5832>
- [2] Pesticide Manual, 13th edition 2004, C.D.S. Tomlin (Ed.), British Crop Protection Council, Farnham, Surrey, UK

BSI — British Standards Institution

BSI is the independent national body responsible for preparing British Standards. It presents the UK view on standards in Europe and at the international level. It is incorporated by Royal Charter.

Revisions

British Standards are updated by amendment or revision. Users of British Standards should make sure that they possess the latest amendments or editions.

It is the constant aim of BSI to improve the quality of our products and services. We would be grateful if anyone finding an inaccuracy or ambiguity while using this British Standard would inform the Secretary of the technical committee responsible, the identity of which can be found on the inside front cover.
Tel: +44 (0)20 8996 9000. Fax: +44 (0)20 8996 7400.

BSI offers members an individual updating service called PLUS which ensures that subscribers automatically receive the latest editions of standards.

Buying standards

Orders for all BSI, international and foreign standards publications should be addressed to Customer Services. Tel: +44 (0)20 8996 9001.
Fax: +44 (0)20 8996 7001. Email: orders@bsi-global.com. Standards are also available from the BSI website at <http://www.bsi-global.com>.

In response to orders for international standards, it is BSI policy to supply the BSI implementation of those that have been published as British Standards, unless otherwise requested.

Information on standards

BSI provides a wide range of information on national, European and international standards through its Library and its Technical Help to Exporters Service. Various BSI electronic information services are also available which give details on all its products and services. Contact the Information Centre.
Tel: +44 (0)20 8996 7111. Fax: +44 (0)20 8996 7048. Email: info@bsi-global.com.

Subscribing members of BSI are kept up to date with standards developments and receive substantial discounts on the purchase price of standards. For details of these and other benefits contact Membership Administration.
Tel: +44 (0)20 8996 7002. Fax: +44 (0)20 8996 7001.
Email: membership@bsi-global.com.

Information regarding online access to British Standards via British Standards Online can be found at <http://www.bsi-global.com/bsonline>.

Further information about BSI is available on the BSI website at <http://www.bsi-global.com>.

Copyright

Copyright subsists in all BSI publications. BSI also holds the copyright, in the UK, of the publications of the international standardization bodies. Except as permitted under the Copyright, Designs and Patents Act 1988 no extract may be reproduced, stored in a retrieval system or transmitted in any form or by any means – electronic, photocopying, recording or otherwise – without prior written permission from BSI.

This does not preclude the free use, in the course of implementing the standard, of necessary details such as symbols, and size, type or grade designations. If these details are to be used for any other purpose than implementation then the prior written permission of BSI must be obtained.

Details and advice can be obtained from the Copyright & Licensing Manager.
Tel: +44 (0)20 8996 7070. Fax: +44 (0)20 8996 7553.
Email: copyright@bsi-global.com.