# INTERNATIONAL STANDARD

ISO 3527

Second edition 2000-10-15

# Oil of parsley fruits (*Petroselinum sativum* Hoffm.)

Huile essentielle de fruits de persil (Petroselinum sativum Hoffm.)



Reference number ISO 3527:2000(E)

#### ISO 3527:2000(E)

#### **PDF** disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

#### © ISO 2000

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office Case postale 56 • CH-1211 Geneva 20 Tel. + 41 22 749 01 11 Fax + 41 22 749 09 47 E-mail copyright@iso.ch Web www.iso.ch

Printed in Switzerland

#### **Foreword**

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this International Standard may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

International Standard ISO 3527 was prepared by Technical Committee ISO/TC 54, Essential oils.

This second edition cancels and replaces the first edition (ISO 3527:1975), which has been technically revised.

Annexes A and B of this International Standard are for information only.

## Oil of parsley fruits (Petroselinum sativum Hoffm.)

#### 1 Scope

This International Standard specifies certain characteristics of the oil of parsley fruits (*Petroselinum sativum* Hoffm.), in order to facilitate assessment of its quality.

#### 2 Normative references

The following standards contain provisions which, through reference in this text, constitute provisions of this International Standard. For dated references, subsequent amendments to, or revisions of, any of these publications do not apply. However, parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the standards indicated below. For undated references, the latest edition of the standard referred to applies. Members of ISO and IEC maintain registers of currently valid International Standards.

ISO/TR 210, Essential oils — General rules for packaging, conditioning and storage.

ISO/TR 211, Essential oils — General rules for labelling and marking of containers.

ISO 212, Essential oils — Sampling.

ISO 279, Essential oils — Determination of relative density at 20 °C — Reference method.

ISO 280, Essential oils — Determination of refractive index.

ISO 592, Essential oils — Determination of optical rotation.

ISO 709, Essential oils — Determination of ester value.

ISO 875, Essential oils — Evaluation of miscibility in ethanol.

ISO 1242, Essential oils — Determination of acid value.

ISO 11024-1, Essential oils — General guidance on chromatographic profiles — Part 1: Preparation of chromatographic profiles for presentation in standards.

ISO 11024-2, Essential oils — General guidance on chromatographic profiles — Part 2: Utilization of chromatographic profiles of samples of essential oils.

#### 3 Term and definition

For the purposes of this International Standard, the following term and definition applies.

#### 3.1

#### oil of parsley fruits

essential oil obtained by steam distillation of the ripe fruits of cultivated parsley (*Petroselinum sativum* Hoffm.) of the Apiaceae family

NOTE 1 Oil of parsley fruits is commercially known as "oil of parsley seed".

NOTE 2 CAS number of oil of parsley fruits: 94334-31-3.

#### 4 Requirements

#### 4.1 Appearance

Clear liquid, sometimes crystallized.

#### 4.2 Colour

Almost colourless to amber yellow.

#### 4.3 Odour

Characteristic of the crushed fruit, but distinct from that of the green part of the plant.

#### 4.4 Relative density at 20 °C, $d_{20}^{20}$

Minimum: 1,043 Maximum: 1,083

#### Refractive index at 20 °C 4.5

1,5130 Minimum: 1,522 0 Maximum:

#### 4.6 Optical rotation at 20 °C

Between -10° and -4°.

#### Miscibility in 85 % ethanol (volume 4.7 fraction) at 20 °C

It shall not be necessary to use more than 6 volumes of 85 % ethanol (volume fraction) to obtain a clear solution with 1 volume of essential oil.

#### Acid value 4.8

Maximum:

#### 4.9 Ester value

Minimum: 1 Maximum: 10

#### 4.10 Chromatographic profile

Analysis of the essential oil shall be carried out by gas chromatography. In the chromatogram obtained, the representative and characteristic components shown in Table 1 shall be identified. The proportions of these components, indicated by the integrator, shall be as shown in Table 1. This constitutes the chromatographic profile of the essential oil.

Table 1 — Chromatographic profile

Component	Min.	Max.
	%	%
$\alpha$ -Pinene	10	22
β-Pinene	7	15
Myristicin	25	50
Apiol	5	35
1,2,3,4- Tetramethoxy-5- allylbenzene	1	12
Elemicin	1	12

NOTE The chromatographic profile is normative, contrary to typical chromatograms given for information in annex A.

#### 4.11 Flashpoint

Information on the flashpoint is given in annex B.

#### Sampling

See ISO 212.

Minimum volume of test sample: 25 ml.

This volume allows each of the tests specified in this International Standard to be carried out at least once.

#### 6 **Test methods**

### Relative density at 20 °C, $d_{20}^{20}$

See ISO 279.

#### 6.2 Refractive index at 20 °C

See ISO 280.

#### Optical rotation at 20 °C

See ISO 592.

#### 6.4 Miscibility in 85 % ethanol (volume fraction) at 20 °C

See ISO 875.

#### 6.5 Acid value

See ISO 1242.

#### Ester value

See ISO 709. Test sample: 2 g. Saponification time: 1 h.

#### Chromatographic profile

See ISO 11024-1 and ISO 11024-2.


### Packaging, labelling, marking and storage

See ISO/TR 210 and ISO/TR 211.

## Annex A

(informative)

## Typical chromatograms of the analysis by gas chromatography of the essential oil of parsley fruits (*Petroselinum sativum* Hoffm.)



#### **Peak identification**

- 1 α-Pinene
- 2 β-Pinene
- 3 β-Phellandrene + limonene
- 4 Myristicin
- 5 Elemicin
- 6 1,2,3,4-Tetramethoxy-5allylbenzene
- 7 Apiol

#### **Operating conditions**

Column: capillary, silica; length 50 m; internal diameter 0,25 mm

Stationary phase: polydimethyl siloxane (OV 101)

Film thickness: 0,25 μm


Oven temperature: programmed from 60 °C to 200 °C at a rate of 2,5 °C/min

Injector temperature: 270 °C Detector temperature: 280 °C Detector: flame ionization type Carrier gas: hydrogen Volume injected: 0,05 μl

Carrier gas flow rate: 2 ml/min

Split ratio: 1/60

Figure A.1 — Typical chromatogram taken on an apolar column



#### **Peak identification**

- 1 α-Pinene
- 2 β-Pinene
- 3 β-Phellandrene
- 4 1,2,3,4-Tetramethoxy-5allylbenzene
- 5 Elemicin
- 6 Myristicin
- 7 Apiol

#### **Operating conditions**

Column: capillary, silica; length 20 m; internal diameter 0,1 mm

Stationary phase: polyethylene glycol 20 000

Film thickness: 0,20 μm

Oven temperature: 50 °C for 1 min, then programmed from 50 °C to 200 °C at a rate

of 10 °C/min

Injector temperature: 250 °C Detector temperature: 250°C Detector: flame ionization type Carrier gas: hydrogen Volume injected: 0,2 μl

Carrier gas flow rate: 0,3 ml/min

Split ratio: 1/350

Pressure programming: 220,7 kPa<sup>1)</sup> for 20 s, then 34,5 kPa/min up to 310,3 kPa,

then 310,3 kPa for 20 min

Figure A.2 — Typical chromatogram taken on a polar column

<sup>1)</sup> 1 kPa = 0,145 psi

## Annex B (informative)

## **Flashpoint**

#### **B.1 General information**

For safety reasons, transport companies, insurance companies, and people in charge of safety services require information on the flashpoints of essential oils, which in most cases are flammable products.

A comparative study on the relevant methods of analysis (see ISO/TR 11018<sup>2</sup>)) concluded that it was difficult to recommend a single apparatus for standardization purposes, given that:

- there is wide variation in the chemical composition of essential oils;
- the volume of the sample needed for certain requirements would be too costly for high-priced essential oils;
- as there are several different types of equipment which can be used for the determination, users cannot be expected to use one specified type only.

Consequently, it was decided to give a mean value for the flashpoint in an informative annex in each International Standard, for information, in order to meet the requirements of the interested parties.

The equipment with which this value was obtained should be specified.

For further information see ISO/TR 11018<sup>2</sup>).

#### **B.2** Flashpoint of oil of parsley fruits

The mean value is + 50 °C.

NOTE Obtained with "Luchaire" equipment.

<sup>2)</sup> ISO/TR 11018, Essential oils — General guidance on the determination of flashpoint.

ISO 3527:2000(E)

ICS 71.100.60

Price based on 5 pages

 $\hfill \hfill \square$  ISO 2000 – All rights reserved