INTERNATIONAL STANDARD



4315

H-13-15

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION •МЕЖДУНАРОДНАЯ ОРГАНИЗАЦИЯ ПО СТАНДАРТИЗАЦИИ •ORGANISATION INTERNATIONALE DE NORMALISATION

Surface active agents — Determination of alkalinity — Titrimetric method

Agents de surface — Détermination de l'alcalinité — Méthode titrimétrique

First edition - 1977-02-01

UDC 661.185:543.241

Ref. No. ISO 4315-1977 (E)

Descriptors: surfactants, chemical analysis, determination of content, alkalinity, volumetric analysis.

Price based on 2 pages

SO 4315-1977 (E

FOREWORD

ISO (the International Organization for Standardization) is a worldwide federation of national standards institutes (ISO member bodies). The work of developing International Standards is carried out through ISO technical committees. Every member body interested in a subject for which a technical committee has been set up has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work.

Draft International Standards adopted by the technical committees are circulated to the member bodies for approval before their acceptance as International Standards by the ISO Council.

International Standard ISO 4315 was drawn up by Technical Committee ISO/TC 91, Surface active agents, and was circulated to the member bodies in August 1975.

It has been approved by the member bodies of the following countries:

Romania

Australia India Austria Iran Belgium Italy Brazil Japan Canada Netherlands Egypt, Arab Rep. of New Zealand France Poland Portugal Germany

South Africa, Rep. of Spain Switzerland Thailand Turkey

United Kingdom U.S.A. U.S.S.R. Yugoslavia

No member body expressed disapproval of the document.

© International Organization for Standardization, 1977 ●

Printed in Switzerland

Hungary

Surface active agents — Determination of alkalinity — Titrimetric method

0 INTRODUCTION

Solutions, pastes and powders of organic surface active substances, in particular those which contain hydrolysable products, can become weakly alkaline as a result of various breakdown reactions.

The method specified in this International Standard is based on the determination of the alkalis in the form of carbonates, bicarbonates, alkali hydroxides or free organic bases such as triethanolamine.

1 SCOPE AND FIELD OF APPLICATION

This International Standard specifies a titrimetric method for the determination of the alkalinity of surface active agents in general.

This method is applicable only if so indicated in the specific standard for each product; in particular it is not suitable for products containing soaps.

2 REFERENCE

ISO 607, Surface active agents — Detergents — Methods of sample division. 1)

3 PRINCIPLE

Titration of a clear solution of the surface active agent with a standard volumetric hydrochloric acid solution, in the presence of methyl orange or bromophenol blue as as indicator.

4 REAGENTS

During the analysis, use only reagents of recognized analytical grade and only distilled water or water of equivalent purity.

- 4.1 Hydrochloric acid, 0,1 N standard volumetric solution.
- 4.2 Methyl orange or bromophenol blue, 1 g/l solution.
- 1) In preparation. (Revision of ISO/R 607.)

5 APPARATUS

Ordinary laboratory apparatus, and in particular:

- **5.1 Conical flask**, capacity 250 ml, complying with the requirements of ISO 1773.
- **5.2** Burette, capacity 10 ml, complying with the requirements of class A of ISO 385.

6 SAMPLING

The laboratory sample of the surface active agent shall be prepared and stored according to the instructions given in ISO 607.

7 PROCEDURE

7.1 Test portion

Weigh, to the nearest 0,001 g, about 5 g of the laboratory sample into the conical flask (5.1).

7.2 Determination

Add 50 ml of water to the test portion (7.1) and dissolve or homogenize if necessary. Then add 2 drops of the indicator solution (4.2) and titrate with the hydrochloric acid solution (4.1) until the end-point is reached.

NOTE — To facilitate the colour comparison, prepare by the same procedure a solution of about 5 g of the laboratory sample in 50 ml of water and add 2 drops of the indicator solution (4.2).

8 EXPRESSION OF RESULTS

The alkalinity of the product, expressed as a percentage by mass of sodium oxide (Na_2O), is given by the formula

$$\frac{V \times T \times 3,1}{m}$$

where

V is the volume, in millilitres, of the standard volumetric hydrochloric acid solution (4.1) used;

T is the exact normality of the standard volumetric hydrochloric acid solution (4.1) used;

m is the mass, in grams, of the test portion.

9 TEST REPORT

The test report shall include the following particulars:

- a) all information necessary for the complete identification of the sample;
- b) the reference of the method used;

- c) the results obtained and the form in which they are expressed;
- d) the test conditions;
- e) any operation not included in this International Standard or regarded as optional, as well as any incidents which may have affected the results.