

---

---

**Essential oil of geranium**  
**(*Pelargonium* × *ssp.*)**

*Huile essentielle de géranium (Pelargonium × ssp.)*



Reference number  
ISO 4731:2012(E)

© ISO 2012

.....



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2012

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 4731 was prepared by Technical Committee ISO/TC 54, *Essential oils*.

This third edition cancels and replaces the second edition (ISO 4731:2006), which has been technically revised.



# Essential oil of geranium (*Pelargonium* × *ssp.*)

## 1 Scope

This International Standard specifies certain characteristics of essential oil of geranium (*Pelargonium* × *ssp.*) coming from different *Pelargonium* species commonly known as *Pelargonium graveolens*, in order to facilitate assessment of its quality.

## 2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO/TR 210, *Essential oils — General rules for packaging, conditioning and storage*

ISO/TR 211, *Essential oils — General rules for labelling and marking of containers*

ISO 212, *Essential oils — Sampling*

ISO 279, *Essential oils — Determination of relative density at 20 °C — Reference method*

ISO 280, *Essential oils — Determination of refractive index*

ISO 592, *Essential oils — Determination of optical rotation*

ISO 875, *Essential oils — Evaluation of miscibility in ethanol*

ISO 1242, *Essential oils — Determination of acid value*

ISO 11024 (all parts), *Essential oils — General guidance on chromatographic profiles*

## 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

### 3.1

#### **essential oil of geranium**

essential oil obtained by steam distillation of the fresh or slightly withered herbaceous parts of *Pelargonium* × *spp.* of the Geraniaceae family, which have given rise to differing ecotypes according to geographical regions

Note 1 to entry: For information on the CAS number, see ISO/TR 21092.[2]

## 4 Requirements

### 4.1 Appearance

Clear, mobile liquid.

### 4.2 Colour

Various shades of amber yellow to greenish yellow.

For the Bourbon type<sup>1)</sup>: yellowish green to brownish green.

### 4.3 Odour

Rose-like, with a varying minty note.

### 4.4 Relative density at 20 °C, $d_{20}^{20}$

Value	North Africa	China	Bourbon type <sup>1</sup>	Madagascar
Minimum	0,885	0,882	0,885	0,887
Maximum	0,905	0,899	0,897	0,897

### 4.5 Refractive index at 20 °C

Value	North Africa	China	Bourbon type <sup>1</sup>	Madagascar
Minimum	1,461	1,460	1,460	1,462
Maximum	1,475	1,472	1,470	1,471

### 4.6 Acid value

North Africa	China	Bourbon type <sup>1</sup>	Madagascar
<10	<10	<10	<10

### 4.7 Optical rotation at 20 °C

Value	North Africa	China	Bourbon type <sup>1</sup>	Madagascar
Maximum	-14°	-14°	-17°	-17°
Minimum	-8°	-7°	-9°	-9°

### 4.8 Miscibility in ethanol 70 % volume fraction at 20 °C

It shall not be necessary to use more than 3 volumes of ethanol 70 % volume fraction to obtain a clear solution with 1 volume of essential oil.

### 4.9 Chromatographic profile

Carry out the analysis of the essential oil by gas chromatography. Identify in the chromatogram obtained the representative and characteristic components shown in Table 1. The proportions of these components, indicated by the integrator, shall be as shown in Table 1. This constitutes the chromatographic profile of the essential oil.

1) Bourbon type includes Reunion Island origin (Bourbon Island is the former name of Reunion Island).

Table 1 — Chromatographic profile

Component	North Africa		China		Bourbon type <sup>a</sup>		Madagascar	
	min. %	max. %	min. %	max. %	min. %	max. %	min. %	max. %
(Z)-rose oxide	0,7	1,5	1,3	3,5	0,3	1,1	0,4	1,4
(E)-rose oxide	0,3	0,6	0,5	1,5	0,1	0,5	0,1	0,6
Menthone	n.d. <sup>b</sup>	2,1	n.d. <sup>b</sup>	2,5	n.d. <sup>b</sup>	2,0	n.d. <sup>b</sup>	2,0
Isomenthone	4,0	8,0	4,0	7,0	5,0	10,0	5,0	10,0
Linalool	4,0	8,5	2,0	4,5	8,0	11,0	4,0	10,0
Guaia-6,9-diene	n.d. <sup>b</sup>	0,5	4,0	7,0	5,0	8,5	5,0	9,0
Citronellyl formate	4,0	8,0	7,0	12,0	6,5	11,0	6,5	11,0
$\alpha$ -Terpineol	0,3	0,6	0,1	0,5	0,3	1,2	0,3	1,0
Geranyl formate	2,0	7,0	1,0	3,0	4,0	8,0	3,8	7,0
Citronellol	25,0	36,0	32,0	43,0	18,0	26,0	18,0	26,0
Geraniol	10,0	18,0	5,0	12,0	12,0	20,0	10,0	20,0
Geranyl butyrate	0,7	2,0	0,4	1,0	0,7	2,0	0,7	1,7
10- <i>epi</i> - $\gamma$ -Eudesmol	3,0	6,2	n.d. <sup>b</sup>	n.d. <sup>b</sup>	n.d. <sup>b</sup>	n.d. <sup>b</sup>	n.d. <sup>b</sup>	n.d. <sup>b</sup>
Geranyl tiglate	0,9	2,0	1,0	1,6	0,7	2,0	0,7	2,0
$\beta$ -Phenylethyl tiglate	0,5	1,2	0,4	1,0	0,4	1,0	0,4	1,0

NOTE The chromatographic profile is normative, contrary to typical chromatograms given for information in [Annex A](#).

<sup>a</sup> Bourbon type includes Reunion Island origin (Bourbon Island is the former name of Reunion Island).

<sup>b</sup> Not detectable.

#### 4.10 Flashpoint

Information on the flashpoint is given in [Annex B](#).

### 5 Sampling

Sampling shall be performed in accordance with ISO 212.

Minimum volume of test sample: 50 ml.

NOTE This volume allows each of the tests specified in this International Standard to be carried out at least once.

### 6 Test methods

#### 6.1 Relative density at 20 °C, $d_{20}^{20}$

Determine the relative density in accordance with ISO 279.

#### 6.2 Refractive index at 20 °C

Determine the refractive index in accordance with ISO 280.

#### 6.3 Optical rotation at 20 °C

Determine the optical rotation in accordance with ISO 592.

**6.4 Acid value**

Determine the acid value in accordance with ISO 1242.

**6.5 Miscibility in ethanol 70 % volume fraction at 20 °C**

Determine the miscibility in accordance with ISO 875.

**6.6 Chromatographic profile**

Determine the chromatographic profile in accordance with ISO 11024.


**7 Packaging, labelling, marking and storage**

These items shall be in accordance with ISO/TR 210 and ISO/TR 211.



**Annex A**  
(informative)

**Typical chromatograms of the analysis by gas chromatography of  
essential oil of geranium (*Pelargonium* × spp.)**

**Key****Peak identification**


- 1 Linalool
- 2 (*Z*)-rose oxide
- 3 (*E*)-rose oxide
- 4 Menthone
- 5 Isomenthone
- 6  $\alpha$ -Terpineol
- 7 Citronellol
- 8 Geraniol
- 9 Citronellyl formate
- 10 Geranyl formate
- 11 Guaia-6,9-diene
- 12 Geranyl butyrate
- 13 Phenylethyl tiglate
- 14 10-*epi*- $\gamma$ -Eudesmol
- 15 Geranyl tiglate

**Operating conditions**

- Column: capillary, length 50 m, internal diameter 0,2 mm  
 Stationary phase: poly dimethylsiloxane  
 Film thickness: 0,25  $\mu$ m  
 Oven temperature: temperature programming from 65 °C to 230 °C,  
 temperature gradient 2 °C/min  
 Injector temperature: 230 °C  
 Detector temperature: 250 °C  
 Detector: flame ionization type  
 Carrier gas: hydrogen  
 Volume injected: 0,2  $\mu$ l  
 Carrier gas flow rate: 1,1 ml/min  
 Split ratio: 1/100

*t* time

**Figure A.1 — Typical chromatogram of the analysis of North Africa oil of geranium taken on an apolar column**

**Key****Peak identification**

- |    |                                     |
|----|-------------------------------------|
| 1  | ( <i>Z</i> )-rose oxide             |
| 2  | ( <i>E</i> )-rose oxide             |
| 3  | Menthone                            |
| 4  | Isomenthone                         |
| 5  | Linalool                            |
| 6  | Guaia-6,9-diene                     |
| 7  | Citronellyl formate                 |
| 8  | $\alpha$ -Terpineol                 |
| 9  | Geranyl formate                     |
| 10 | Citronellol                         |
| 11 | Geraniol                            |
| 12 | Geranyl butyrate                    |
| 13 | 10- <i>epi</i> - $\gamma$ -Eudesmol |
| 14 | Geranyl tiglate                     |
| 15 | Phenylethyl tiglate                 |

**Operating conditions**

Column: capillary, length 50 m, internal diameter 0,2 mm	
Stationary phase: poly(ethylene glycol) (Carbowax 20 M <sup>a</sup> )	
Film thickness: 0,25 $\mu$ m	
Oven temperature: temperature programming from 65 °C to 230 °C, temperature gradient 2 °C/min	
Injector temperature: 230 °C	
Detector temperature: 250 °C	<i>t</i> time
Detector: flame ionization type	
Carrier gas: hydrogen	
Volume injected: 0,2 $\mu$ l	
Carrier gas flow rate: 1,1 ml/min	
Split ratio: 1/100	

<sup>a</sup> Carbowax 20M is an example of a suitable product available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of this product.

**Figure A.2 — Typical chromatogram of the analysis of North Africa oil of geranium taken on a polar column**

## Annex B (informative)

### Flashpoint

#### B.1 General information

For safety reasons, transport companies, insurance companies, and people in charge of safety services require information on the flashpoints of essential oils, which in most cases are flammable products.

A comparative study on the relevant methods of analysis (see ISO/TR 11018<sup>[1]</sup>) concluded that it was difficult to recommend a single apparatus for standardization purposes, given that:

there is a wide variation in the chemical composition of essential oils;

the volume of the sample needed in certain requirements would be too costly for high-priced essential oils;

as there are several different types of equipment which can be used for the determination, users cannot be expected to use one specified type only.

Consequently, it was decided to give a mean value for the flashpoint annexed to each International Standard, for information, in order to meet the requirements of the interested parties.

The equipment with which this value was obtained has to be specified.

For further information, see ISO/TR 11018.<sup>[1]</sup>

#### B.2 Flashpoint of the essential oil of geranium

The mean values are indicated in Table B.1.

**Table B.1 — Mean flashpoint according to origin**

	North Africa	China	Bourbon type <sup>a</sup>	Madagascar
<b>Mean flashpoint, °C</b>	86	84	83	83
<sup>a</sup> Bourbon type includes Reunion Island origin (Bourbon Island is the former name of Reunion Island).				

NOTE Obtained with Luchoire<sup>2)</sup> equipment.

2) Equipment available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of this product.

## Bibliography

- [1] ISO/TR 11018, *Essential oils — General guidance on the determination of flashpoint*
- [2] ISO/TR 21092, *Essentials oils — Characterization*

© ISO 2012. All rights reserved.

