
**Foodstuffs — Determination of
ochratoxin A in cereals and cereal
products —**

Part 1:

High performance liquid chromatographic
method with silica gel clean up

*Produits alimentaires — Dosage de l'ochratoxine A dans les céréales et
produits dérivés —*

*Partie 1: Méthode par chromatographie liquide haute performance
comprenant une étape d'extraction par chromatographie sur gel de silice*



Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

International Standard ISO 15141-1 was prepared by the European Committee for Standardization (CEN) in collaboration with ISO Technical Committee TC 34, *Agricultural food products*, Subcommittee SC 4, *Cereals and pulses*, in accordance with the Agreement on technical cooperation between ISO and CEN (Vienna Agreement).

Throughout the text of this standard, read "...this European Standard..." to mean "...this International Standard...".

ISO 15141 consists of the following parts, under the general title *Foodstuffs — Determination of ochratoxin A in cereals and cereal products*:

- *Part 1: High performance liquid chromatographic method with silica gel clean up*
- *Part 2: High performance liquid chromatographic method with bicarbonate clean up*

Annexes A and B of this part of ISO 15141 are for information only.

© ISO 1998

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the publisher.

International Organization for Standardization
Case postale 56 • CH-1211 Genève 20 • Switzerland
Internet iso@iso.ch

Printed in Switzerland

Contents

Foreword	iii
1 Scope	1
2 Normative references	1
3 Principle	1
4 Reagents	1
5 Apparatus and equipment	3
6 Procedure	4
7 Calculation	6
8 Precision	7
9 Test report	7
Annex A (informative) Precision data	8
Annex B (informative) Bibliography.....	9

Foreword

The text of EN ISO 15141-1:1998 has been prepared by Technical Committee CEN/TC 275 "Food analysis - Horizontal methods", the secretariat of which is held by DIN, in collaboration with Technical Committee ISO/TC 34 "Agricultural food products".

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by April 1999, and conflicting national standards shall be withdrawn at the latest by April 1999.

This European Standard „Foodstuffs - Determination of ochratoxin A in cereal and cereal products“ consists of two parts:

Part 1: High performance liquid chromatographic method with silica gel clean up

Part 2: High performance liquid chromatographic method with bicarbonate clean up

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Czech Republic, Denmark, Finland, France, Germany, Greece, Iceland, Ireland, Italy, Luxembourg, Netherlands, Norway, Portugal, Spain, Sweden, Switzerland and the United Kingdom.

4.7 *n*-Hexane**4.8 Dichloromethane****4.9 Acetone****4.10 Methanol**

4.11 Solvent mixture I: toluene (4.6) and glacial acetic acid (4.2) 99+1 parts per volume (V+V)

4.12 Solvent mixture II: acetone (4.9) and toluene (4.6) 5+95 (V+V)

4.13 Solvent mixture III: toluene (4.6) and glacial acetic acid (4.2) 90+10 (V+V)

4.14 Mobile phase

Mix 99 volume parts of acetonitrile (4.5) with 99 volume parts of water and 2 volume parts of glacial acetic acid (4.2) and degas this solution before use.

4.15 Boron trifluoride

4.16 Boron trifluoride in methanol solution, $\rho(\text{BF}_3) = 14 \text{ g}/100 \text{ ml}$

WARNING: Use a well maintained fume hood. Avoid contact with skin, eyes, and respiratory tract.

4.17 Ochratoxin A, in crystal form or as a film in ampoules

4.18 Ochratoxin A stock solution

Dissolve 1 mg of the ochratoxin A (crystals) (4.17) or the contents of 1 ampoule (if ochratoxin A has been obtained as a film) in solvent mixture I (4.11) to give a solution containing approximately 20 $\mu\text{g}/\text{ml}$ to 30 $\mu\text{g}/\text{ml}$ of ochratoxin A.

To determine the exact concentration, record the absorption curve between a wavelength of 300 nm and 370 nm in 5 nm steps in a 1 cm quartz cell (5.5) with solvent mixture I (4.11) as reference. Identify the wavelength for maximum absorption by recording in 1 nm steps around the maximum as reference. Calculate the mass concentration of ochratoxin A, ρ_{OTA} , in micrograms per millilitre of solution using equation 1:

$$\rho_{\text{OTA}} = A_{\text{max}} \times \frac{M \times 100}{\kappa \times \delta} \quad (1)$$

where

A_{max} is the absorption determined at the maximum of the absorption curve (here: at 333 nm);

M is the relative molecular mass of ochratoxin A ($M = 403 \text{ g}/\text{mol}$);

κ is the molar absorption coefficient of ochratoxin A, in solvent mixture I (here: $544 \text{ m}^2/\text{mol}$);

δ is the path length of the cell in centimetres.

4.19 Ochratoxin A standard solution $\rho_{\text{OTA}} = 1 \mu\text{g}/\text{ml}$

Evaporate under a nitrogen flow 1 ml of the stock solution (4.18) or the aliquot portion which is

equivalent to an absolute amount of 100 µg of ochratoxin A to dryness and dilute to 100 ml with the mobile phase (4.14).

This solution can be stored in a refrigerator at 4 °C. Stability shall be checked.

4.20 Ochratoxin A calibration solutions

Pipette suitable volumes of ochratoxin A standard solution (4.19), e.g. 1 ml, 2,5 ml, 4 ml and 5 ml into e.g. a 100 ml volumetric flask (5.12) and dilute to the mark with the mobile phase (4.14). The amount of ochratoxin A in the calibration solutions should cover the range of 0,2 ng to 1,0 ng per 20-µl-injection volume.

4.21 Sodium hypochlorite solution, $\rho(\text{NaOCl}) = 4 \text{ g}/100 \text{ ml}$

5 Apparatus and equipment

Usual laboratory equipment and, in particular, the following:

5.1 Laboratory mill, suitable to grind to 1 mm

5.2 Rotary evaporator, with a water bath capable of being controlled between 20 °C and 50 °C

5.3 Mechanical shaker

5.4 Spectrometer, suitable for measurement at wavelengths of 300 nm up to 370 nm, having a spectral band width of not more than $\pm 2 \text{ nm}$

5.5 Quartz cells, with 1 cm optical path length and no significant absorption between wavelengths of 300 nm and 370 nm

5.6 Centrifuge tubes, e.g. of capacity 250 ml, plastic made of high density polyethylene (HDPE), with screw cap

5.7 Cooling centrifuge, preferably a refrigerated centrifuge, capable of producing a gravitational force of at least 3500 g at the base of the centrifuge tubes (5.6)

5.8 Solid phase extraction columns, e.g. SEP-PAK®¹⁾ disposable silica gel

After the pack has been opened, condition at 105 °C for 2 h and store over activated silica gel with moisture indicator. Before use, wash with 10 ml of toluene (4.6). Check the recovery with each new batch. In the case of use of SEP-PAK columns, the cartridges have the following specification:

- mean mass of the packing material: 690 mg
- pore size: 12,5 nm
- particle size: 55 µm to 105 µm

in a 3 ml polypropylene tube.

5.9 Solvent containers, such as syringes, e.g. of 50 ml capacity with central opening and stop-cock

¹⁾ SEP PAK® is an example of a suitable product available commercially. This information is given for the convenience of users of this Standard and does not constitute an endorsement by CEN of these products.

5.10 Pear-shaped flasks, 50 ml, with ground glass joint

5.11 Separating funnel, 50 ml

5.12 Volumetric flask, 100 ml

5.13 Membrane filter for aqueous solutions, made of polytetrafluoroethylene (PTFE), with a diameter of 4 mm and a pore size of 0,45 µm

5.14 Sieve, with an aperture size of not more than 1 mm

5.15 Vials with crimped caps or screw cap vials

5.16 Microsyringe, of capacity 500 µl

5.17 HPLC apparatus, comprising the following

5.17.1 High performance liquid chromatograph, eluent reservoir, a pump, an injection system, a fluorescence detector with variable wavelength setting and a data processing, e.g. an integrator with plotter.

5.17.2 Analytical reversed phase HPLC separating column, C₁₈, e.g. Lichrospher® 100 RP 18²⁾ which ensures a baseline resolved resolution of the ochratoxin A peak from all other peaks.

- length: 250 mm
- internal diameter: 4 mm
- spherical particles of size: 5 µm

NOTE: Shorter columns can also be used (e.g. a column with a length of 120 mm to 150 mm)

5.17.3 Precolumn, C₁₈

- length: 40 mm
- internal diameter: 4 mm
- spherical particles of size: 5 µm

6 Procedure

6.1 General

The whole analytical procedure should be performed in one working day. If several samples are processed at the same time all samples should be analysed during the following night using an automatic sample injector.

6.2 Preparation of the test samples

Grind the laboratory sample using a laboratory mill (5.1) until it passes through the sieve (5.14) and mix it thoroughly.

NOTE: Grinding is not necessary for wheat flour with a maximum size of 250 µm.

²⁾ Lichrospher® 100 RP 18 is an example of a suitable product available commercially. This information is given for the convenience of users of this Standard and does not constitute an endorsement by CEN of these products.

6.3 Extraction of ochratoxin A from the sample

Place 20 g (m_0), weighed to the nearest 0,1 g, of the sample prepared as in 6.2 in a centrifuge tube (5.6). For ochratoxin A contents of more than 5,0 µg/kg repeat the analysis using a test portion of 10 g, otherwise the risk of reduced recovery has to be taken into account. Successively add 30 ml of hydrochloric acid solution (4.3), 50 ml of magnesium chloride solution (4.4), stir with a glass rod, and add 100 ml of toluene (4.6) (V_1).

Shake for 60 min and subsequently centrifuge the suspension. The centrifugation time depends on the efficiency of the centrifuge, while cooling prevents loss of toluene. Remove 50 ml (= toluene aliquot portion V_2) from the upper toluene layer and load it onto the solid phase mini disposable column which has been prepared as in 5.8 and to which the syringe (5.9) is attached as solvent reservoir.

NOTE 1: Care should be taken not to overload the column.

Wash the column twice with 10 ml of n-hexane (4.7) and again, twice with 10 ml of solvent mixture II (4.12). Subsequently wash with 5 ml of toluene. Discard all the washings.

Elute ochratoxin A with two 15 ml portions of solvent mixture III (4.13) into a 50 ml pear-shaped flask (5.10). Evaporate the eluate under reduced pressure to dryness cautiously without exceeding 40 °C. Take up the residue by pipetting 1 ml (V_3) of the mobile phase (4.14) into the pear-shaped flask and filter through a membrane (5.13) into a vial (5.15) (= sample test solution).

NOTE 2: Elution of ochratoxin A and the subsequent steps in the procedure described in this clause can depend on the type of solid phase extraction columns that is used. The elution volume for example should be checked to be appropriate for the type of column that is used.

NOTE 3: The size and/or shape of the flask can have a negative influence on the recovery.

6.4 HPLC operating conditions

When the column according to 5.17.2 and the mobile phase according to 4.14 were used the following settings were found to be appropriate.

Flow rate:	1 ml/min
Fluorescence detection:	Excitation wavelength: 330 nm
	Emission wavelength: 460 nm
Injection volume:	20 µl (V_4)

6.5 Calibration graph

Prepare a calibration graph at the beginning of the analysis and whenever the chromatographic conditions change.

Inject at least four calibration solutions of different suitable concentrations (see 4.20).

Plot the fluorescence values of the ochratoxin A calibration solutions (4.20) against the ochratoxin A mass concentrations in nanograms.

Ensure that the linearity check is carried out [10].

6.6 Identification

Identify ochratoxin A by comparing the retention time of the sample with that of the standard substance.

Sometimes it can be necessary to identify the ochratoxin A peak by simultaneous injection of sample test solution and standard solution.

6.7 Determination

Immediately chromatograph the sample. To carry out the determination by the external standard method, integrate the peak area or determine the peak height, and compare the results with the corresponding values for the standard substance with the nearest peak area/height, or use a calibration graph. In the case of a calibration graph, additional solutions with concentrations within the linear range may be prepared for the calibration graph.

Inject equal volumes of sample test solution and standard solution used for the calibration graph.

Read off the mass of ochratoxin A, (m_1), in nanograms, corresponding to the fluorescence of the sample test solution from the calibration graph.

If the ochratoxin A response of the sample is outside the calibration graph, adjust the amount of sample injected by concentrating or diluting the sample test solution.

6.8 Confirmation

If necessary confirm the identity by disappearance of the peak at the retention time for ochratoxin A and appearance of a new peak at the same retention time as that of standard methyl ester of ochratoxin A.

Take 500 μ l of the extract prepared as in 6.3, transfer into a pear-shaped flask and evaporate to dryness in a rotary evaporator (5.2). Take up the residue in 1 ml of dichloromethane (4.8), and add 2 ml of boron trifluoride methanol solution (4.16).

Stopper the flask tightly and heat it in a water bath at 50 °C to 60 °C for 15 min. After cooling, transfer the solution into a 50 ml separating funnel containing 30 ml of water, shake 3 times with 10 ml of dichloromethane each time for 30 s. Combine the organic phases in a second 50 ml separating funnel, add 20 ml of water for washing and shake for 30 s.

Subsequently filter the dichloromethane phase through sodium sulfate (4.1) into a pear-shaped flask, evaporate to dryness, take up in 500 μ l of mobile phase (4.14) and subject this solution to chromatographic separation under the conditions as described in 6.4. The completeness of derivatization can be checked from the chromatograms. It is possible with this procedure to verify mass fractions of ochratoxin A of not less than 0,4 μ g/kg.

An adequate standard solution (4.19) should be treated separately to check the retention times of the ochratoxin A methyl ester and the completeness of the derivatization.

7 Calculation

Calculate the mass fraction w_{OTA} of ochratoxin A in micrograms per kilogram using equation (2) (external standard method):

$$w_{OTA} = \frac{V_1 \times V_3 \times m_1}{V_2 \times V_4 \times m_0} \quad (2)$$

where

V_1 is the volume of the solvent used for extraction (6.2), in millilitres, here: 100 ml;

V_2 is the volume of the centrifugate (toluene aliquot portion), in millilitres, here: 50 ml;

V_3 is the total volume of the sample test solution, in millilitres, here: 1 ml;

V_4 is the injection volume, in millilitres;

m_1 is the mass of ochratoxin A corresponding to the measured peak area or peak height read off the calibration graph, in nanograms;

m_0 is the mass of the test portion, in grams.

Report the result according to current legislation and after rounding to two decimal places.

Indicate whether or not a correction for recovery has been applied.

8 Precision

8.1 General

Details of the interlaboratory test of the precision of the method according to ISO 5725:1986 [1] are summarized in annex A. The values derived from the interlaboratory tests may not be applicable to analyte concentration ranges and matrices other than given in annex A.

8.2 Repeatability

The absolute difference between two single test results found on identical test material by one operator using the same apparatus within the shortest feasible time interval will exceed the repeatability limit r in not more than 5 % of the cases.

The values for wheat whole flour are:

$\bar{x} = 0,41 \mu\text{g/kg}$	$r = 0,18 \mu\text{g/kg}$
$\bar{x} = 1,23 \mu\text{g/kg}$	$r = 0,70 \mu\text{g/kg}$

8.3 Reproducibility

The absolute difference between two single test results on identical test material reported by two laboratories will exceed the reproducibility limit R in not more than 5 % of the cases.

The values for wheat whole flour are:

$\bar{x} = 0,41 \mu\text{g/kg}$	$R = 0,30 \mu\text{g/kg}$
$\bar{x} = 1,23 \mu\text{g/kg}$	$R = 1,10 \mu\text{g/kg}$

9 Test report

The test report shall contain at least the following data:

- all information necessary for the identification of the sample;
- a reference to this draft European Standard or to the method used;
- the results and the units in which the results have been expressed;
- date and type of sampling (if known);
- date of receipt of the laboratory sample;
- date of test;
- any particular points observed in the course of the test;
- any operations not specified in the method or regarded as optional which might have affected the results.

Annex A (informative) Precision data

The following data were obtained in interlaboratory tests according to ISO 5725 : 1986 [1] conducted by the Max-von-Pettenkofer-Institute of the Federal Health Office, Foodchemistry Department, Berlin, Germany on wheat whole flour [5], [6].

Table A.1

Sample	wheat whole flour	wheat whole flour
Year of inter-laboratory test	1993	1991
Number of laboratories	13	13
Number of samples	1	1
Number of laboratories retained after eliminating outliers	13	13
Number of outliers	0	0
Number of accepted results	65	65
Mean value \bar{x} ($\mu\text{g}/\text{kg}$)	0,407	1,227
Repeatability standard deviation s_r ($\mu\text{g}/\text{kg}$)	0,062	0,248
Repeatability relative standard deviation RSD_r	15,32 %	20,21%
Repeatability limit r ($\mu\text{g}/\text{kg}$)	0,176	0,702
Reproducibility standard deviation s_R ($\mu\text{g}/\text{kg}$)	0,105	0,388
Reproducibility relative standard deviation RSD_R	25,80 %	31,62 %
Reproducibility limit R ($\mu\text{g}/\text{kg}$)	0,298	1,097
Recovery	90 % \pm 15 %	80 % \pm 15 %

Annex B (informative) Bibliography

- [1] ISO 5725:1986 Precision of test methods - Determination of repeatability and reproducibility for a standard test method by inter-laboratory tests.
- [2] Majerus, P., Cutka, I., Dreyer, A., El-Dessouki, S., Eyrich, W., Reusch, H., Schurer, B., and Waiblinger, H.U.: Zur Belastungssituation von Ochratoxin A in Lebensmitteln pflanzlichen Ursprungs. In: Dt. Lebensm. Rundsch., 89, Vol 4 (1993) pp 112 ff.
- [3] Jiao, Y., Blaas, W., Rühl, Ch., and Weber, R.: Ochratoxin A in Lebensmitteln pflanzlicher Herkunft. In: Dt. Lebensm. Rundsch., 90, Vol 10 (1994) pp 318 ff.
- [4] Jiao, Y., Blaas, W., Rühl, Ch., and Weber, R.: Identification of ochratoxin A in food samples by chemical derivatization and gas chromatography - mass spectrometry. In: J. Chromat. 595 (1992) pp. 364 - 367.
- [5] Untersuchung von Lebensmitteln: Bestimmung von Ochratoxin A: L 15.00-1 1992-12 (Food Analysis: Determination of Ochratoxin A in cereals and cereal products L 15.00-1 1992-12) in: Amtliche Sammlung von Untersuchungsverfahren nach § 35 LMBG: Verfahren zur Probenahme und Untersuchung von Lebensmitteln, Tabakerzeugnissen, kosmetischen Mitteln und Bedarfsgegenständen/Bundesgesundheitsamt (In: Collection of official methods under article 35 of the German Federal Foods Act; Methods of sampling and analysis of foods, tobacco products, cosmetics and commodity goods/Federal Health Office) Loseblattausgabe, Stand Aug. 1993 Bd. 1(Loose leaf edition, as of 1993 - 08 Vol. I.) Berlin, Köln, Beuth Verlag GmbH.
- [6] Majerus, P., Weber, R., and Wolff, J.: Nachweis und Bestimmung von Ochratoxin A in Getreide und Getreideprodukten (Detection and determination of Ochratoxin A in cereals and cereal products) In: Bundesgesundheitsblatt (Journal of the Federal Health Office) 37, Nov. 1994, no 11, pp.454 - 458.
- [7] Tauchmann, F.; Mintzlaff, H.-J.; Leistner, L.: Schutzmaßnahmen beim Arbeiten mit Mykotoxinen (Protective measures for working with mycotoxins) Alimenta 1972, 11, 85.
- [8] Castegnaro, M., Hunt, D.C., Sansone, E.B., Schuller, P.L., Siriwardana, M.G., Telling, G.M., van Egmond, H.P., and Walker, E.A.: Laboratory decontamination and destruction of aflatoxins B₁, B₂, G₁ and G₂ in laboratory wastes. In: IARC Scientific publication no 37, International Agency for Research on Cancer (WHO), Lyon, France; 1980, 59p.
- [9] Castegnaro, M., Berek, J., Fremy, J.M., Lafontaine, M., Miraglia, M., Sansone, E.B., and Telling, G.M.: Laboratory decontamination and destruction of carcinogens in laboratory wastes. In: IARC Scientific publication no 113, International Agency for Research on Cancer (WHO), Lyon, France; 1991, 63p.
- [10] van Trijp, J.M.P. and Roos, A.H.: Model for the calculation of calibration curves, RIKILT Report 91.02, January 1991.

ICS 67.060

Price based on 9 pages
